

Appendix References. General plus specific (including key, supplemental bibliographic) sources of input to Appendix Table A-2 and Appendix Table A-3, ordered alphabetically by scientific family name

General

Kells V, Rocha LA, Allen LG (2016) A Field Guide to Coastal Fishes from Alaska to California. Johns Hopkins Univ Press, Baltimore.

Love MS, Yoklavich M, Thorsteinson L (2002) The Rockfishes of the Northeast Pacific. Univ Calif Press, Berkeley & Los Angeles, CA. 404 pp

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA. 650 pp

Shanks AL, Eckert GL. (2005) Population persistence of California current fishes and benthic crustaceans: a marine drift paradox. Ecol Monogr 75, 505-524

Achiridae

Warburton (1979) Growth and production of some important species of fish in a Mexican coastal lagoon system. J Fish Biol 14, 449-464

Acipenseridae

Brennan JS (1988) Age determination and verification of California white sturgeon, *Acipenser transmontanus*: a comparative analysis. MSc thesis, San Jose State Univ

Brennan JS, Cailliet GM (1989) Comparative age-determination techniques for white sturgeon in California. Trans Am Fish Soc 118, 296-310

Chapman FA, Van Eenennaam JP, Doroshov SI (1996) The reproductive condition of white sturgeon, *Acipenser transmontanus*, in San Francisco Bay, California. Fish Bull 94, 628-634

Deng DF, Van Eenennaam JP, Doroshov SI (2002) Comparison of early life stages and growth of green and white sturgeon. Pages 237–248 in (W Van Winkle, PJ Anders, DH Secor, DA Dixon, eds), Biology, Management, and Protection of North American sturgeon. Am Fish Soc Symp 28, Bethesda, MD

Kynard B, Parker E, Parker T (2005) Behavior of early life intervals of Klamath River green sturgeon, *Acipenser medirostris*, with a note on body color. Environ Biol Fish 72, 85-97

Rien TA, Beamesderfer RC (1994) Accuracy and precision of white sturgeon age estimates from pectoral fin rays. Trans Am Fish Soc 123, 255-265

Agonidae

Arbour JH, Avendaño P, Hutchings JA (2010) Aspects of the ecology and life history of alligatorfish *Aspidophoroides monopterygius*. Environ Biol Fish 87:353–362

Busby MS, Ambrose DA (1993) Development of larval and early juvenile pygmy poacher *Odontopyxis trispinosa*, and blacktip poacher *Xeneretmus latifrons* (Scorpaeniformes: Agonidae). FishBull_91, 397-413

Glubukov AI, Orlov AM (2009) Data on distribution and biology of poachers Agonidae from the northwestern part of the Bering Sea. J Ichthyol_48, 426-442

Jensen GC (2005) A unique feeding method by a teleost fish, the fourhorn poacher *Hypsagonus quadricornis* (Agonidae). Biol Bull_09, 165-167

Smith RL, Gillispie JG (1983) Notes on the biology of the Bering Poacher, *Occella dodecaedron* (Tilesius), and the sturgeon poacher, *Agonus acipenserinus* Tilesius in the Southeast Bering Sea. Copeia 1983, 454-460

Tokranov AM (1993) Sexual dimorphism in sea poachers (Agonidae) off Kamchatka. J Ichthyol_33, 113-122

Tokranov AM (2000) Distribution and some features of the biology of *Bathyagonus nigripinnis* (Agonidae) in Pacific waters of Southeast Kamchatka and the northern Kurils. J Ichthyol 40, 585

Tokranov AM, Orlov AM (2013) Specific features of distribution and ecology of sawback poacher *Sarritor frenatus* (Agonidae) in Pacific waters of the northern Kuril Islands and southeast Kamchatka. J Ichthyol_53, 209-219

Albulidae

Donovan MK, Friedlander AM, Harding KK, Schemmel EM, Filous A, Kamikawa K, Torkelson N (2015) Ecology and niche specialization of two bonefish species in Hawaii. Environ Biol Fish 98, 2159–2171

Filous A, Lennox RJ, Eveson P, Raveino R, Clua EEG, Cooke SJ, Danylchuk AJ (2019a) Population dynamics of roundjaw bonefish *Albula glossodonta* at a remote coralline Atoll inform community-based management in an artisanal fishery. Fish Manag Ecol. 2019, 1–15.

Filous A, | Lennox RJ, Coleman RR, Friedlander AM, Clua EEG, Danylchuk AJ (2019b) Life-history characteristics of an exploited bonefish *Albula glossodonta* population in a remote South Pacific atoll. J Fish Biol 95, 562-574

Kamikawa KT, Friedlander AM, Harding KK, Filous A, Donovan MK, Schemmel E (2015) Bonefishes in Hawai'i and the importance of angler-based data to inform fisheries management. Environ Biol Fish 98, 2147–2157

Pfieler E, Padron R, Crabtree RE (2000). Growth rate, age and size of bonefish from the Gulf of California. J Fish Biol 56, 448-453

Pfieler E, Watts T, Pugh J, van der Heiden AM (2008) Speciation and demographic history of the Cortez bonefish, *Albula* sp. A (Albuliformes: Albulidae), in the Gulf of California inferred from mitochondrial DNA. J Fish Biol 73, 382-394

Wallace EM(2015) High intraspecific connectivity in the Indo-Pacific bonefishes: implications for conservation and management. Environ Biol Fish 98, 2173–2186

Alopiidae

Cailliet GM, Bedford DW (1983) The biology of three pelagic sharks from California waters, and their emerging fisheries: a review. Calif Coop Ocean Fish Invest Rep 24, 57-69

Camhi MD -Pikitch EK, Babcock EA, eds (2008) Sharks of the Open Ocean: Biology, Fisheries and Conservation. Blackwell

Cartamil DP (2009) Movement patterns, habitat preferences, and fisheries biology of the common thresher shark (*Alopias vulpinus*) in the Southern California Bight. PhD thesis, Univ Calif, San Diego

Cartamil DP, Sepulveda CH, Wegner NC, Aalbers SA, Baquero A, Graham JB (2011) Archival tagging of subadult and adult common thresher sharks (*Alopias vulpinus*) off the coast of southern California. Mar Biol 158, 935-944

Chen C-T, Liu K-M, Chang Y-C (1997) Reproductive biology of the bigeye thresher shark. Ichthyol Res 44, 227-2350

Drew M,, White WT, Dharmadi, Harry AV, Huveneers C (2015) Age, growth and maturity of the pelagic thresher *Alopias pelagicus* and the scalloped hammerhead *Sphyrna lewini*. J Fish Biol 86, 333-354

Kohler NE, Casey JG, Turner PA (1995) Length-weight relationships for 13 species of sharks from the western North Pacific..Fish Bull 93, 412-418

Liu K-M, Chiang C-T, Chen C-T (1998). Age and growth estimates of the bigeye thresher shark, *Alopias superciliosus*, in northeastern Taiwan waters. Fish Bull 96, 482-491

Liu K-M, Chen C-T, Liao H-S, Joung S-J. (1999) Age, growth, and reproduction of the pelagic thresher shark, *Alopias pelagicus* in the Northwestern Pacific. Copeia 1999(1),68-74

Mollet HF, Cailliet GM (2002) Comparative population demography of elasmobranchs using life history tables, Leslie matrices and stage-based matrix models. Mar Freshwat Res 53,503-516

Smith SE, Rasmussen RC, Ramon DA, Cailliet GM (2008) The biology and ecology of thresher sharks (Alopiidae). Pp 60-68 in (Camhi MD, Pikitch EK, Babcock EA, eds) Sharks of the Open Ocean: Biology, Fisheries and Conservation. Blackwell

Ammodytidae

Robards MD; Willson MF, Armstrong RH, Piatt JF (eds) (1999). Sand lance: a review of biology and predator relations and annotated bibliography. Res. Pap. PNW-RP-521. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 327 p

Robards MD, Rose GA, Piat JF (2002) Growth and abundance of Pacific sand lance, *Ammodytes hexapterus*, under differing oceanographic regimes. Environ Biol Fish 64, 429–441

Anarrhichadidae

Marliave JB (1987) The life history and captive reproduction of the wolf-eel *Anarrhichthys ocellatus* at the Vancouver Public Aquarium. International Zoo Yearbook 26, 70-81

Anoplomatidae

Echave KB, Hanselman DH, Adkison MD, Sigler MF (2012) Interdecadal change in growth of sablefish (*Anoplopoma fimbria*) in the northeast Pacific Ocean. Fish Bull 110, 361-374

Fujiwara S,-Hankin DG (1988) Sex ratio, spawning period, and size and age at maturity of sablefish *Anoplopoma fimbria* off northern California. Nipp Suis Gakk 54,1333-1338

Guzmaan JAM, Luckenbach JA, Middleton MA, Massee KC, Jensen C, Goetz FW, Jasonowicz AJ, Swanson P. (2017) Reproductive life history of sablefish (*Anoplopoma fimbria*) from the U.S. Washington coast. PLoS ONE 12 (9),e0184413, 24pp

Hanselman DH, Heifetz J, Echave KB, Dressel SC (2015) Move it or lose it: movement and mortality of sablefish tagged in Alaska. *Can J Fish Aquat Sci* 72, 238-251

Head M, Keller AA, Bradburn M (2014) Maturity and growth of sablefish, *Anoplopoma fimbria*, along the U.S. west coast. *Fish Res* 159, 156-167

Howard KB (2008) Inter-decadal change in sablefish, *Anoplopoma fimbria*, growth and maturity in the Northeast Pacific Ocean. MSc thesis, Univ Alaska, Fairbanks

Kapur M, Haltuch M, B. Connors B, Rogers L, Berger A, , E. Koontz E, Cope J, Echave K, Fenske K, Hanselman D, Punt AE (2020) Oceanographic features delineate growth zonation in Northeast Pacific sablefish. *Fish Res* 222, 105414

Maloney NE, Sigler MF (2008) Age-specific movement patterns of sablefish (*Anoplopoma fimbria*) in Alaska. *Fish Bull* 106, 305-316

Morita SH, Morita K, Nishimura A (2012) Sex-biased dispersal and growth in sablefish (*Anoplopoma fimbria*) in the northeastern Pacific Ocean. *Environ Biol Fish* 94, 505-511

Rodgveller CJ, Stark JW, Echave KB, Hulson P-JF (2016) Age at maturity, skipped spawning, and fecundity of female sablefish (*Anoplopoma fimbria*) during the spawning season. *Fish Bull* 114, 89-102

Sogard SM, Berkeley SA (2017) Patterns of movement, growth, and survival of adult sablefish (*Anoplopoma fimbria*) at contrasting depths in slope waters off Oregon. *Fish Bull* 115, 233-251

Tolimieri N, Haltuch MA, Lee Q, Jacox MG, Bograd SJ (2018) Oceanographic drivers of sablefish recruitment in the California Current. *Fish Oceanogr* DOI: 10.1111/fog.12266

Zolotov OG, Spirin IY (2012) On the fishery and abundance of skillfish (*Erilepis zonifer*) at the Emperor Seamounts (North-West Pacific). *Trans Pac Res Fish Centre (TINRO)* 168, 79-88

Zolotov OG, Spirin IY, Zudina SM (2014) New data on the range, biology, and abundance of skillfish *Erilepis zonifer* (Anoplopomatidae). *J Ichthyol* 54, 251-265

Apogonidae

Lea RN, Rosenblatt RH (2000) Observations on fishes associated with the 1997-98 El Nino off California. *Calif Coop Ocean Fish Invest Rep* 41, 117-129

Atherinopsidae

Clark FN (1925) The life history of *Leuresthes tenuis*, an atherine fish with tide-controlled spawning habits. Calif Fish Game Comm, Fish Bull 10, 51p

Clark FN (1929) The life history of the California Jack Smelt, *Atherinopsis californiensis*. Calif Fish Game Div, Fish Bull 16, 23p

Santos AJ, Frederick AR, Higgins BA, Carrillo A, Carter AL, Dickson KA,. German DP, Horn MH (2018) The beach-spawning California grunion *Leuresthes tenuis* eats and digests conspecific eggs. J Fish Biol; 93, xx-xx (Orcid ID: 0000-0002-6971-4197)

Schultz LP (1933) The age and growth of *Atherinops affinis oregonia* Jordan and Snyder and of other subspecies of baysmelt along the Pacific coast of the United States. Univ Washington Press.

Bathymasteridae

Glubokov AI (2009) New data on the blue-eyed searcher *Bathymaster signatus* (Bathymasteridae, Perciformes) from the Northwestern part of the Bering Sea. J Ichthyol 49 154-161

Batrachoididae

Bass AH, Horvath BJ, Brothers EB (1996) Nonsequential developmental trajectories lead to dimorphic vocal circuitry for males with alternative reproductive tactics. Develop Neurobiol 30, 493-504

Bose APH (2017) Offspring cannibalism and parental care in the plainfin midshipman fish..PhD thesis, McMaster University. 270p

Bose APH, Adragna JB, Balshine S (2017) Otolith morphology varies between populations, sexes and male alternative reproductive tactics in a vocal toadfish *Porichthys notatus*. J Fish Biol 90, 311–325

Brantley RK, Tseng J, Bass AH (1993) The ontogeny of inter-and intrasexual vocal muscle dimorphisms in a sound producing fish. Brain Behav Evol _42, 331-349

DeMartini EE (1990) Annual variations in fecundity, egg size and condition of the plainfin midshipman (*Porichthys notatus*). Copeia 1990(3), 850-855

Hubbs CL (1920) The bionomics of *Porichthys notatus* (Girard). Am Nat 54, 380-384

LeMoine CMR, Bucking C, Craig PM, Walsh PJ (2014) Divergent hypoxia tolerance in adult males and females of the plainfin midshipman (*Porichthys notatus*) *Physiol Biochem Zool*, 87, 325-

Miller JS, Mazzoldi C, Rasotto MB, Balshine S (2019) Differential investment in male accessory glands: lessons from a marine fish with alternative reproductive tactics. *Mar Biol* 166, 37_s00227-019-3474-8

Sak BP (1990) Age, growth and reproductive characteristics of the plainfin midshipman (*Porichthys notatus*) in Monterey Bay, with notes on seasonal movements. MSc thesis, San Francisco State Univ

Sisneros JA, Alderks PW, Leon K, Sniffen B (2009) Morphometric changes associated with the reproductive cycle and behaviour of the intertidal-nesting, male plainfin midshipman *Porichthys notatus*. *J Fish Biol* 74, 18-36

Blenniidae

Labelle M, Nursall JR (1985) Some aspects of the early life history of the redlip blenny, *Ophioblennius atlanticus* (Teleostei: Blenniidae). *Copeia* 1985(1), 39-49

Muss A, Robertson DR, Stepien CA, Wirtz P, Bowen BW (2001) Phylogeography of *Ophioblennius*: the role of ocean currents and geography in reef fish evolution. *Evolution* 55, 561-572

Riginos C, Victor BC (2001) Larval spatial distributions and other early life-history characteristics predict genetic differentiation in Eastern Pacific blennioid fishes. *Proc Roy Soc Lond B* 268, 1931-1936

Stephens JS Jr, Johnson RK, Key GS, McCosker JE (1970) The comparative ecology of three sympatric species of California blennies of the genus *Hypsoblennius* Gill (Teleostomi, Blenniidae). *Ecol Monogr* 40, 213-233

Stevens EG, Moser HG (2002) Observations on the realy life history of the mussel blenny blenny, *Hypsoblennius jenkinsi* and the bay blenny, *Hypsoblennius gentilis*, from specimens reared in the laboratory. *Calif Coop Ocean Fish Invest Rep* 23, 269-275

Bothidae

Tapia-Garcia M, García-Abad MC, Ladrón de Guevara GC (2000) Reproduction, distribution and abundance of *Bothus constellatus* (Pisces: Bothidae), in the Gulf of Tehuantepec, Mexico. *Rev Biol Trop* 48, 205-213

Bramidae

Pearcy WG, Fisher JP, Yoklavich MM (1992) Biology of the Pacific pomfret (*Brama japonica*) in the North Pacific Ocean. Can J Fish Aquat Sci 50, 2608-2625

Polovina JJ, Abecassis M, Howell EA, Woodworth P (2009) Increases in the relative abundance of mid-trophic level fishes concurrent with declines in apex predators in the subtropical North Pacific, 1996–2006. Fis Bull 107,:523–531

Savinykh VF, Vlasova LV (1994) The length-age structure and growth rate of the Pacific pomfret, *Brama japonica* (Bramidae). J Ichthyol 34, 494-500

Seki MP, Mundy BC (1991) Some notes on the early life stages of the Pacific pomfret, *Brama japonica*, and Other Bramidae from the Central North Pacific Ocean Jpn J Ichthyol 38, 63-68

Carangidae

Ben-Aderet NJ (2017) The biology of yellowtail (*Seriola lalandi*) in the Southern California Bight::spatial insights from recreational catch records, tagging and life-history characteristics. PhD thesis, Univ Calif, San Diego, 124 p.

Ben-Aderet NJ, Johnston EM, Cravey R, Sandin SA, (2020) Revisiting the life history of yellowtail jack (*Seriola dorsalis*) in the Southern California Bight: new evidence for ontogenetic habitat shifts and regional differences in a changing environment. Fish Bull 188, 157-170

Baxter JL (1960) A study of the yellowtail *Seriola dorsalis* (Gill). Calif Dep Fish Game, Fish Bull 110, 96 p

Espino Barr E, Gallardo Cabello EM, Cabral Solís EG, Garcia Boa A, Puente Gómez M (2008) Growth of the Pacific green jack *Caranx caninus* (Pisces: Carangidae). Rev Biol Trop 56, 171-179

Gillanders BM,-Ferrell DJ, Andrew NL (1999a) Aging methods for yellowtail kingfish, *Seriola lalandi*, and results from age- and size-based growth models. Fish Bull 97, 812-827

Gillanders BM, Ferrell DJ, Andrew NL (1999b) Size at maturity and seasonal changes in gonad activity of yellowtail kingfish (*Seriola lalandi*; Carangidae) in New South Wales, Australia, N ZI J Mar Freshwat Res 333, 457-468

Madigan DJ, Snodgrass OE, Fisher NS (2018) From migrants to mossbacks: tracer- and tag-inferred habitat shifts in the California yellowtail *Seriola dorsalis*. Mar Ecol Prog Ser 597: 221–230

Martinez-Takeshita N, Purcell CM, Chabot CL, Craig MT, Paterson CN, et al (2015) A tale of three tails: cryptic speciation in a globally distributed marine fish of the genus *Seriola*. *Copeia* 103, 357-368

Maricchiolo MV, Genovese L (1999) The reproductive biology of the amberjack, *Seriola dumerilii* (Risso 1810). I. oocyte development in captivity. *Aquacult Res*, 30, 349-355

Neabenzahl DA (1997) Age, growth, and population structure of jack mackerel (*Trachurus symmetricus*) from the Northeastern Pacific Ocean. MSc thesis, San Francisco State Univ

Renones O, Morales-Nin B, Deuderos S (1999) Population size structure, age and growth of *Naucrates ductor* in the small scale FADs fishery of Mallorca Island (western Mediterranean). *Sci Mar* 63, 355-366

Shiraishi T, Ohshimo S, Yukami R (2010) Age, growth and reproductive characteristics of gold striped amberjack *Seriola lalandi* in the waters off western Kyushu, Japan. *N Z J Mar Freshwat Res* 44, 117-127

Shiraishi T, Tanaka H, Ohshimo S, Ishida H, Morinaga N (2010) Age, growth and reproduction of two species of scad, *Decapterus macrosoma* and *D. macarellus* in the waters off Southern Kyushu. *Jpn Agric Res Q* 44(2), 197-206

Vassilopoulou V, Siapatis A, Christides G, Bekas P (2004) The biology and ecology of juvenile pilotfish (*Naucrates ductor*) associated with fish aggregating devices (FADs) in eastern Mediterranean waters. *Medit Mar Sci* 5, 61-70

Widodo J (1991) Maturity and spawning of shortfin scad (*Decapterus macrosoma* (Carangidae) of the Java Sea. *Asian Fish Sci* 4, 245-252

Carcharhinidae

Bonfil R, Clarke S, Nakano H (2008) The biology and ecology of the oceanic whitetip shark, *Carcharhinus longimanus*. Pp 128-139 in (Camhi MD, Pikitch EK, Babcock EA, eds), *Sharks of the Open Ocean: Biology, Fisheries and Conservation*. Blackwell

Cailliet GM, Bedford DW (1983) The biology of three pelagic sharks from California waters, and their emerging fisheries: a review. *Calif Coop Ocean Fish Invest Rep* 24, 57-69

Cailliet GM, Musick JA, Simpfendorfer CA, Stevens JD (2005) Ecology and life history characteristics of chondrichthyan fish. Chapter 3, pp 12-18 in (SL Fowler, RD Cavanagh, M Camhi, GH Burgess, G Cailliet, SV Fordham [eds]), *Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes, Status Survey*. IUCN/Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK. 461 p

Cruz-Martinez A, Chiappa-Carrara X, Arenas-Fuentes V (2004) Age and growth of the bull shark, *Carcharhinus leucas*, from southern Gulf of Mexico. J NW Atl Fish Sci 35, 367-374

Drew M, Rogers P, Huveneers C (2017) Slow life-history traits of a neritic predator, the bronze whaler (*Carcharhinus brachyurus*). Mar Freshwat Res 68, 461-472

Goldman KJ, Cailliet GM, Andrews AH, Natanson LJ (2012) Assessing the age and growth of chondrichthyan fishes. Ch14, pp 423- 451 in (Zzzzz, eds) Biology of Sharks and their Relatives. Taylor & Francis Group

Joung S-J2, Jun-Hsu Chen J-H, Chien-Pang Chin C-P, Liu K-M (2015) Age and growth of the dusky shark, *Carcharhinus obscurus*, in the western North Pacific Ocean. Terr Atmos Ocean Sci 26(2), Pt 1, 153-160

Allen H. Andrews Æ W. Hunt Howell (2008) Using bomb radiocarbon analyses to validate age and growth estimates for the tiger shark, *Galeocerdo cuvier*, in the western North Atlantic. Mar Biol 154:423–434

Lucifora LO, Menni RC, Escalante AH (2005) Reproduction and seasonal occurrence of the copper shark, *Carcharhinus brachyurus*, from north Patagonia, Argentina. ICES J Mar Sci 62, 107-115

Meyer CG, O'Malley JM, Papastamatiou YP, Dale JJ, Hutchinson MR, Anderson JM, Royer MA, Holland KM (014) Growth and maximum size of tiger sharks (*Galeocerdo cuvieri*) in Hawaii. PLoS ONE 9(issue 1), e84799

Nakano H, Stevens JD (2008) The biology and ecology of the blue shark, *Prionace glauca*. Pp 140-151 in (Camhi MD, Pikitch EK, Babcock EA, eds) Sharks of the Open Ocean: Biology, Fisheries and Conservation. Blackwell

Natanson LJ, Gervelis BJ, Winton MV, Hamady LL, Gulak SJB, Carlson JK (2014) Validated age and growth estimates for *Carcharhinus obscurus* in the northwestern Atlantic Ocean, with pre- and post management growth comparisons. Environ Biol Fish 97, 881-896

Pratt HL Jr (1978) Reproduction in the blue shark, *Prionace glauca*. Fish Bull 77, 445-470

Queiroz- N, Humphries NE, Couto A, Vedor M, da Costa I, Sequeira AMM et al (2020) Global spatial risk assessment of sharks under the footprint of fisheries. Nature Res 572, 461-4xx <https://doi.org/10.1038/s41586-019-1444-4xx>

Simpfendorfer CA, Rory B. McAuley RB, Chidlow J, Unsworth P (2002) Validated age and growth of the dusky shark, *Carcharhinus obscurus*, from Western Australian waters. Mar Freshwat Res 53, 567-573

Wintner SP, Dudley SFJ, Kistnasamy AN, Everett B, ,etal (2002) Age and growth estimates for the Zambezi shark, *Carcharhinus leucas*, from the east coast of South Africa. Mar Freshwat Res 53, 557-566

Zhu A, Dai X, Xu L, Chen X, Chen Y (2011) Reproductive biology of female blue shark *Prionace glauca* in the southeastern Pacific Ocean. Environ Biol Fish 91, 95-102

Centrolophidae

Horn MH (1977) Observations on feeding, growth, locomotor behavior, and buoyancy of a pelagic stromateid fish, *Ichthyos lockingtoni*. FishBull_75, 453-456

Okamoto M, Ida H, Sugisaki H, Kurita Y (2002) Larval and juvenile development of the medusafish, *Ichthyos lockingtoni* (Perciformes: Centrolophidae), from the Northwest Pacific, and a comparison with Northeast Pacific larval and juvenile forms. Ichthyol Res 49, 93-96

Cetorhinidae

Natanson LJ, Wintner SP, Johansson F, Piercy A, Campbell P, De Maddalena A, Gulak SJB, Human B, Fulgosi FC, Ebert DA, Hemida F, Mollen FH, Vanni S, Burgess GH, Compagno LJV (2008) Ontogenetic vertebral growth patterns in the basking shark *Cetorhinus maximus*. Mar Ecol Progr Ser 361, 267-278

Pauly D (2002) Growth and mortality of the basking shark *Cetorhinus maximus* and their implications for management of whale sharks *Rhincodon typus*. Occasional Paper of the IUCN Species Survival Commission, Shark Specialist Group, 199-208

Chaenopsidae

Thresher RE (1977) Field observations on the behavior and ecology of the orangethroat pikeblenny, *Chaenopsis alepidota*, in the Sea of Cortez. Bull Mar Sci 60, 1195-1204

Chimaeridae

Barnett LAK (2008) Life history, abundance, and distribution of the spotted ratfish, *Hydrolagus coliei*. MSc thesis, Calif State Univ, Monterey Bay

Barnett LAK, Earley RL, Ebert DA, Cailliet GM (2009a) Maturity, fecundity, and reproductive cycle of the spotted ratfish, *Hydrolagus coliei*. *MarBiol* 156, 301-

Barnett LAK, Ebert DA, Cailliet GM (2009) Assessment of the dorsal fin spine for chimaeroid (Holocephali: Chimaeriformes) age estimation. *J Fish Biol* 75, 1258-1270

Barnett LAK, Ebert DA, Cailliet GM (2012) Evidence of stability in a chondrichthyan population: case study of the spotted ratfish *Hydrolagus coliei* (Chondrichthyes: Chimaeridae). *J Fish Biol* 80, 1765-1788

Johnson AG, Horton HF (1972) Length-weight relationship, food habits, parasites, and sex and age determination of the ratfish, *Hydrolagus coliei* (Lay and Bennett). *Fish Bull* 70, 421-429

King JR, McPhie RP (2015) Preliminary age, growth and maturity estimates of spotted ratfish (*Hydrolagus coliei*) in British Columbia. *Deep Sea Res II* 115, 55-63

Clupeidae

Brooks ML, Fleishman E, Brown LR, Lehman PW, Werner I, Scholz N, Mitchelmore C, Lovvorn JR, Johnson ML, Schlenk D, van Drunick S, Drever JI, Stoms DM, Parker AE, Dugdale R (2012) Life histories, salinity zones, and sublethal contributions of contaminants to pelagic fish declines illustrated with a case study of San Francisco Estuary, California, USA. *Ests Coasts* 35, 603-621

Butler JZ, et al (1996) Age composition, growth, and maturation of the Pacific sardine (*Sardinops sagax*) during 1994. *Calif Coop Oceanic Fish Invest Rep* 37, 152-159

Carmona R, Alexandres F (1994) Growth determination of *Opisthonema libertate* (Clupeiformes: Clupeidae) by means of otolith reading. *Rev Biol Trop* 42(1-2), 233-238

Dawson WA (1986) The interpretation of otolith structure for the assessment of age and growth of some pelagic fisheries from the coast of Ecuador: mackerel (*Scomber japonicus* L.), Pacific sardine (*Sardinops sagax*), Pacific thread herrings (*Opisthonema medirastres*, *O. bulleri*), and round herring (*Etrumeus teres*). *Boletin cientifico tecnico, Instit National Pesca (Ecuador)* 24

Feyrer F, Sommer T, Slater SB (2009) Old school vs. new school: status of threadfin shad (*Dorosoma petenense*) five decades after its introduction to the Sacramento–San Joaquin delta. *San Fran Est Watershed Sci* 7, 1-17

Gunderson DR, Dygert PH (1988) Reproductive effort as a predictor of natural mortality rate. *J Conseil int Explor Mer* 44, 200-209

HAY DE, Rose KA, Schweigert J, Megrey BA (2008) Geographic variation in North Pacific herring populations: Pan-Pacific comparisons and implications for climate change impacts. *Progr Oceanogr* 77, 233-240

Houde ED, Zastrow CE (1993) Ecosystem- and taxon-specific dynamic and energetic properties of larval fish assemblages. *Bull Mar Sci* 53, 290-335

Ishida M, Mitani T, Uehara S, Honda H (2004) Stock assessment of the Pacific stock of the round herring *Etrumeus teres*. In: Stock assessment of the fisheries resources around Japan. Tech Rep, Fish Agency and Fish Res Agency, Tokyo, Japan, 471–481

Johnson JE (1970) Age, growth, and population dynamics of threadfin shad, *Dorosoma petenense* (Guenther), in central Arizona reservoirs. *Trans Am Fish Soc* 99, 739-753

Johnson, JE (1971) Maturity and fecundity of threadfin shad, *Dorosoma petenense* (Guenther), in central Arizona reservoirs. *Trans Am Fish Soc* 100, 74-85

Leggett WE, Carscadden JC (1978) Latitudinal variation in reproductive characteristics of American Shad (*Alosa sapidissima*): evidence for population specific life history strategies in fish. *J Fish Res Bd Can* 35, 1469-1478

Lo NCH, Macewicz BJ, Griffith DA (2010) Biomass and reproduction of Pacific sardine (*Sardinops sagax*) off the Pacific northwestern United States, 2003–2005. *Fish Bull* 108, 174-192

Nyuji M, Takasuka A (2017) Spawning cycle and fecundity of a multiple spawner round herring *Etrumeus teres* off southern Japan: oocyte growth and maturation analysis *J Sea Res* 122, 11-18

Olney JE, Denny SC, Hoenig JM (2001) Criteria for determining maturity stage in female American shad, *Alosa sapidissima*, and a proposed reproductive cycle. *Bull Fran Peche Piscicult* 363, 881-901

Oozeki Y, Takasuka A, Kubota H, Barange M (2007) Characterizing spawning habitats of Japanese sardine (*Sardinops melanostictus*), Japanese anchovy (*Engraulis japonicus*), and Pacific round herring (*Etrumeus teres*) in the Northwestern Pacific. *Calif Coop Oceanic Fish Invest Rep* 48, 191-203

Oshimo S, Gotoh T, Otsuka T, Gejima K (2011) Age, growth and reproductive characteristics of round herring *Etrumeus teres* in the East China Sea. *Nippon Suisan Gaikashi* 77, 15-22 [inJpn with Engl abstract]

Palance DG, Macewicz BJ, Stierhoff K, Demer DA, Zwolinski JP (2019) Length conversions and mass–length relationships of five forage-fish species in the California current ecosystem *J Fish Biol* 95, 1116-1124

Plaza G, Honda H, -Sakaji H, Nashida K (2006) Patterns of growth in the early life history of the round herring *Etrumeus teres*. J Fish Biol 68,1421-1435

Plaza G, Sakaji H, Honda H, Hirota Y, Nashida K (2007) Spawning pattern and type of fecundity in relation to ovarian allometry in the round herring *Etrumeus teres*. Mar Biol 152, 1051-1064

Rodriguez-Murillo JA, Soto-Rojas RL (1999) Population dynamics of *Opisthonema medirastre* (Pisces: Clupeidae) in the Pacific coast of Costa Rica. Uniciencia,Heredia_15-16, 61-64

Rodriguez JA, -Palacias JA, Chavarria AL (1989) Sexual maturity and fecundity of the *Opisthonema medirastre* and *O. bulleri* (Pisces: Clupeidae) on the Pacific coast of Costa Rica. Rev Trop Biol 37, 49-54

Ruiz-Dominguez M, Quinonez-Velazquez C (2018) Age, growth, and mortality of *Opisthonema libertate* on the coasts of northwestern Mexico. Ciencias Marinas 44, 235-250

Spratt JD (1981) Status of the Pacific herring, *Clupea harengus pallasii*, resource in California 1972 to 1980. Calif Dep Fish Game, Fish Bull 171,107 p

Surma S, Pitcher TJ, Pakhomov EA (2021) Trade-offs and uncertainties in Northeast Pacific herring fisheries: ecosystem modelling and management strategy evaluation. ICES J Mar Sci fsab125

Vega-Corrales LA (2010) Evaluacion poblacional del stock explotable del complejo *Opisthonema* (Pisces: Clupeidae) en el Golfo de Nicoya, Costa Rica. Rev Mar Cost 2, 83-94 [in Spanish with Engl abstract]

Coryphaenidae

Chang SK, DiNardo G, Farley J, Brodziak J, Yuan ZL. (2013) Possible stock structure of dolphinfish (*Coryphaena hippurus*) in Taiwan coastal waters and globally based on reviews of growth parameters. Fisheries Research 147, 127-13

Uchiyama JH, Burch RK,-Kraal SA Jr (1986) Growth of dolphins, *Coryphaena hippurus* and *C. equiselis* in Hawaiian waters as determined by daily increments on otoliths. Fish Bull 84, 186-191

Zúñiga-Flores MS, Ortega-García S, Rodríguez-Jaramillo M D C, López-Martínez J (2011) Reproductive dynamics of the common dolphinfish *Coryphaena hippurus* in the southern Gulf of California. Mar Biol Res 7, 677-689

Cottidae

Alofs KM, Polivka KM (2004) Microhabitat-scale influences of resources and refuge on habitat selection by an estuarine opportunist fish. *Mar Ecol Progr Ser* 271, 297-306

Awata S, Sasak H, Goto T, Koya Y, Takeshima H, Yamazaki A, Munehara H (2019) Host selection and ovipositor length in eight sympatric species of sculpins that deposit their eggs into tunicates or sponges. *Mar Biol* 166, 59 <https://doi.org/10.1007/s00227-019-3506-4>

Brogan JD, Anderl DM (2012) Great sculpin (*Myoxocephalus polyacanthocephalus*). NMFS, NOAA, AK Fish Sci Cen Rep 18283_DS1

Burge RT, Schultz SA (1973) The marine environment in the vicinity of Diablo Cove with special reference to abalones and bony fishes. *Calif Dep Fish Game, Mar Res Tech Rep* 19, 433p

Buser TJ, Burns MD, Lopez JA (2017) Littorally adaptive? testing the link between habitat, morphology, and reproduction in the intertidal sculpin subfamily Oligocottinae (Pisces: Cottoidea). *PeerJ* 5, e3634

Cartwright RL (2009) Description of early life history stages of the northern sculpin (*Icelinus borealis* Gilbert) (Teleostei: Cottidae). *Fish Bull* 107, 175-185

Chadwick EMP (1976) A comparison of growth and abundance for tidal pool fishes in California and British Columbia. *J Fish Biol* 8, 27-34

COPE JM, Berger AM, Whitman AD, Budrick JE, Bosley KM, Tsou T-S, Niles CB, Privitera-Johnson K, Hillier LK, Hinton KE, Wilson MN (2019) Assessing cabezon (*Scorpaenichthys marmoratus*) stocks in waters off of California and Oregon, with catch limit estimation for Washington State. *Pac Fish Manag Council, Portland, OR. SAFE Rep*, 439 p

Craik GJS (1978). A further investigation of the homing behaviour of the intertidal cottid, *Oligocottus maculosus* Girard. PhD thesis, Univ Brit Columbia

Datsky AV (2017) Biological features of the common fish species in Olyutorsky-Navarin region and the adjacent waters of the Bering Sea: 4. Family sculpins (Cottidae). *J Ichthyol* 57, 341-353

Davis JLD (2000) Spatial and seasonal patterns of habitat partitioning in a guild of southern California tidepool fishes. *Mar Ecol Prog Ser* 196, 253-268

Davis JLD, Levin LA (2002) Importance of pre-recruitment life-history stages to population dynamics of the woolly sculpin *Clinocottus analis*. *Mar Ecol Progr Ser* 234, 229-246

Feeney RF (1987) Development of the eggs and larvae of the yellowchin sculpin, *Icelinus quadriseriatus* (Pisces: Cottidae). *Fish Bull* 85, 201-212

Feeney RF (1992) Post-yolksac larval development of two southern California sculpins, *Clinocottus analis* and *Orthonopias triacis* (Pisces: Cottidae). *Fish Bull* 90, 454-468

Freeman MC, Neally N, Grossman GD (1985) Aspects of the life history of the fluffy sculpin, *Oligocottus snyderi*. Fish Bull 83, 645-655

Grebel JM, Cailliet GM (2010) Age, growth, and maturity of cabezon (*Scorpaenichthys marmoratus*) in California. Calif Fish Game 96, 36-52

Grossman GD, DeVlaming V (1984) Reproductive ecology of female *Oligocottus snyderi* Greeley: a North American intertidal sculpin. J Fish Biol 25, 231-240

Hannah RW, Blume MTO, Thompson JE (2011) Length and age at maturity of female yelloweye rockfish (*Sebastes rubberimus*) and cabezon (*Scorpaenichthys marmoratus*) from Oregon waters based on histological evaluation of maturity. OR Dep Fish Wldlf, Fish Div, Info Rep, number 2009-04

Hoff GR (2000) Biology and ecology of threaded sculpin, *Gymnocanthus pistilliger*, in the eastern Bering Sea. Fish Bull 98, 711-722

Hutchinson CE, TenBrink TT (2011) Age determination of the yellow Irish lord: management implications as a result of new estimates of maximum age. No Am J Fish Manag 31, 1116-1122

Jones AC (1962) The biology of the euryhaline fish *Leptocottus armatus* Girard (Cottidae). Univ Calif Publ Zool 67, 321-368

Lauth RR (1988) Seasonal spawning cycle, spawning frequency, and batch fecundity of the cabezon, *Scorpaenichthys marmoratus*, in Puget Sound, Washington. Fish Bull 87, 145-154

Luksenburg JA, Pedersen T (2002) Sexual and geographical variation in life history parameters of the shorthorn sculpin. J Fish Biol 61, 1453-1464

Luksenburg JA, Pedersen T, Falk-Petersen IB (2004) Reproduction of the shorthorn sculpin *Myoxocephalus scorpius* in northern Norway. J Sea Res 51, 157-166

Matarese AC, Kendall AW Jr, Blood DM, Vinter BM (1989) Laboratory guide to early life history stages of Northeast Pacific fishes. NOAA Tech Rep NMFS 80

Mgaya YD (1992) Density and production of *Clinocottus globiceps* and *Oligocottus maculosus* (Cottidae) in tidepools at Helby Island, British Columbia. Mar Ecol Progr Ser 85, 219-225

Mgaya YD (1995) Age and growth analysis of the mosshead sculpin *Clinocottus globiceps* Girard 1857 (Pisces: Cottidae) from Helby Island, British Columbia J appl Ichthyol 11, 50-59

Mireles C, Nakamura R, Wendt DE (2012) A collaborative approach to investigate site fidelity, home range, and homing behavior of cabezon (*Scorpaenichthys marmoratus*). Fish Res 113, 133-142

Misitano DA (1978) Description of laboratory-reared larvae of the buffalo sculpin, *Enophrys bison* (Pisces:Cottidae). Copeia 1978(4), 35-642

OConnell CP (1953 The life history of the cabezon *Scorpaenichthys marmoratus* (Ayres)). Calif Dep Fish Game, Fish Bull 93, 54pp

Panchenko VV (2001) Reproduction peculiarities of plain sculpin *Myoxocephalus jaok* in Peter the Great Bay, Sea of Japan. Russ J Mar Biol 27, 111-112

Panchenko VV (2002) Age and growth of sculpins of the genus *Myoxocephalus* (Cottidae) in Peter the Great Bay (the Sea of Japan). J Ichthyol_42, 516-522

Panchenko VV (2010) Age and growth of Far Eastern staghorn sculpin *Gymnocanthus herzensteini* Cottidae in Peter the Great Bay (Sea of Japan). J Ichthyology 50, 295-300

Panchenko VV (2012) Growth and Age of Staghorn Sculpins of the Genus *Gymnocanthus* (Cottidae) in Peter the Great Bay and Adjacent Waters of Primorye (Sea of Japan). J Ichthyol 52, 226-238

Petersen CW, Mazzoldi C, Zarrella KA, Hale RE(2005) Fertilization mode, sperm characteristics, mate choice and parental care patterns in *Artedius* spp. (Cottidae) J Fish Biol_67, 239-254

Petersen CW, Zarrella KA, Ruben ZZ, Mazzoldi C (2004) Reproductive biology of the rosy lip sculpin, an intertidal spawner. J Fish Biol 64, 863-875

Pfister CA (1997) Demographic consequences of within-year variation in recruitment. Mar Ecol Progr Ser 153, 229-238

Pierce BE, Pierson KB (1990) Growth and reproduction of the tidepool sculpin *Oligocottus maculosus*. Jpn J Ichthyol 36,4, 410-418

Shelekhov VA, Panchenko VV (2007) Age and growth of threaded sculpin *Gymnocanthus pistilliger* (Cottidae) in the Southern Primorye (Sea of Japan). J Ichthyol_47, 175-183

Reuter RF, TenBrink TT (2007) Bering Sea and Aleutian Islands sculpins 2007. SAFE Rep, 1117-1143. No Pac Fish Manag Council, Portland, OR

Reuter RF, TenBrink TT (2008) NOAA Assessment of sculpin stocks in the Bering Sea/Aleutian Islands. SAFE Rep, 1409-1448. No Pac Fish Manag Council, Portland, OR

Shanks AL, Pfister CA (2009) Annual recruitment of three species of tide-pool fishes is driven by variation in springtime coastal hydrodynamics. Limnol Oceanogr 54, 1481-1487

Shelekhov VA, Panchenko VV (2007) Age and growth of threaded sculpin *Gymnocanthus pistilliger* (Cottidae) in the Southern Primorye (Sea of Japan). J Ichthyol_47, 175-183

Snook R (1997) Aspects of the life history of *Orthonopias triacis* in Monterey Bay, California. MSc thesis, San Jose State Univ

Spies ID, Nichol K, Aydin K, TenBrink TT (2017) Assessment of the sculpin stock complex in the Gulf of Alaska. NMFS, NOAA, GOA SAFE Rep, 27 p. No Pac Fish Manage Council, Portland, OR

Spies I, Ormseth OA, Martin M, TenBrink TT (2011) Assessment of the sculpin complex in the Gulf of Alaska. _No Pac Fish Manag Council, SAFE Rep GOA Sculpin, 1377-1392

Spies I, Ormseth OA, TenBrink TT (2011) Bering Sea and Aleutian Islands sculpins. No Pac Fish Manag Council, SAFE_Rep, BSAI Sculpins, 1735-1770

Stein R (1973) Description of Laboratory-Reared Larvae of *Oligocottus maculosus* Girard (Pisces: Cottidae). Copeia 1973, 373-377

TenBrink TT, Aydin KY (2009) Life history traits of sculpins in the eastern Bering Sea and Aleutian Islands. No Pac Res Bd, Final Rep Project 628, 215p

TenBrink TT, Buckley TW (2013) Life history aspects of the yellow Irish lord (*Hemilepidotus jordani*) Eastern Bering Sea and Aleutian Islands. NW Nat 94, 126-136

Timola O, Luotonen H (1986) Age distribution and growth of the fourhorn sculpin, *Myoxocephalus quadricornis*, in the NE Bothnian Bay and the Archipelago Sea. Bothnian Bay Rep 4, 25-32

Tokranov AM (1984) On the reproduction of the great sculpin *Myoxocephalus polyacanthocephalus* (Pallas) (Cottidae) off Kamchatka. Voprosy Ikhtiologii 24, 601-608

Tokranov AM (1995) Size-sex structure of the genus *Triglops* sculpins (Cottidae) in the coastal Kamchatka waters. Voprosy Ikhtiologii 35, 134-136

Tokranov AM (2014) Some features of biology of butterfly sculpin *Melletes papilio* (Cottidae) in the Pre-Kamchatka waters of the Sea of Okhotsk. J Ichthyol 54, 558-565

Tokranov AM, Orlov AM (2007) Some biological features of rare and poorly studied sculpins (Cottidae, Hemitriptidae, Psychrolutidae) in the Pacific waters off the Northern Kuril Islands and Southeastern Kamchatka, Russian Federation. Raffles Bull Zool, Suppl No 14, 187-198

Washington BB (1982) Identification and systematics of larvae of *Artedius*, *Clinocottus*, and *Oligocottus* (Scorpaeniformes:Cottidae). MSc thesis, Dep Fish Wildlf, OR State Univ

Washington BB (1986) Systematic relationships and ontogeny of the sculpins *Artedius*, *Clinocottus*, and *Oligocottus* (Cottidae: Scorpaeniformes). Proc Calif Acad Sci 44, 157-224

Weiss EF Jr (1962) The age, growth and food habits of the marine cottid *Leptocottus armatus*. MSc thesis, Fish Wildlife Manag, Montana State Univ, 18 p

Weiss EF Jr (1969) The age and growth of the marine cottid *Leptocottus armatus*. Proc Montana Acad Sci 29, 63-71

Wells AW (1986) Aspects of ecology and life history of the wooly sculpin, *Clinocottus analis*, from Southern California. Calif Fish Game 72, 213-226

Yamazaki A, Ogino A, Munehara H (2020) Dispersion and settlement of two sympatric sculpins of the genus *Gymnocanthus*. J Fish Biol Feb2020_jfb.14291

Cryptacanthodidae

Antonenko DV, Solomatov SF, Panchenko VV (2011) On occurrence and biology of Berg Wrymouth *Cryptacanthodes bergi* (Cryptacanthodidae) in the north-western part of the Sea of Japan. J Ichthyol 51, 173-177

Cyclopteridae

Arita GS (1969) Sexual dimorphism in the cyclopterid fish *Eumicrotremus orbis*. J Fish Res Bd Can 26, 3262-3265

Orlov AM, Tokranov AM (2008) Specific features of distribution, some features of biology, and the dynamics of catches of smooth lumpsucker *Aptocyclus ventricosus* (Cyclopteridae) in waters of the Pacific Ocean off the Kuril Islands and Kamchatka. J Ichthyol 4, 81-95

Solomatov SF, Orlov AM (2018). Smooth lumpsucker *Aptocyclus ventricosus* in the northwestern Sea of Japan: distribution and some life history traits. Arch Pol Fish, Fish Aquat Life 26, 5-20

Yoshida H, Yamaguchi H (1985) Distribution and feeding habits of the pelagic smooth lumpsucker *Aptocyclus ventricosus* in the Aleutian Basin. Bull Fac Fish Hokk Univ 36, 200-209

Zhukova K, Maznikova OA, Orlov AM (2018) Gonadal maturation of the female smooth lumpsucker (*Aptocyclus ventricosus*). Fish Bull 116, 153-160

Cyprinodontidae

Bernardi G, Talley D (2000) Genetic evidence for limited dispersal in the coastal California killifish, *Fundulus parvipinnis* . J Exp Mar Biol Ecol 255, 187-199

Fritz ES (1975) The life history of the California killifish *Fundulus parvipinnis* Girard, in Anaheim Bay, California. Calif Dep Fish Game, Fish Bull 165, 91-106

PerezEspana H, Galvan-Magana F,-AbitiaCardenas LA (1998) Growth, consumption, and productivity of the California killifish in Ojo de Liebre Lagoon, Mexico. J Fish Biol 52, 1068-1077

Dalatiidae

Kiraly SJ, Moore JA, Jasinski PH (2003) Deepwater and other sharks of the U.S. Atlantic Ocean Exclusive Economic Zone. Mar Fish Rev 65, 1-63

Dasyatididae

Mollet HF, Cailliet GM (2002) Comparative population demography of elasmobranchs using life history tables, Leslie matrices and stage-based matrix models. Mar Freshwat Res 53, 503-516

Mollet HF (2002) Distribution of the pelagic stingray, *Dasyatis violacea* (Bonaparte, 1832), off California, Central America, and worldwide. Mar Freshwat Res 53, 525-530

Mollet HF, Ezcurra JM, O'Sullivan JB (2002) Captive biology of the pelagic stingray, *Dasyatis violacea* (Bonaparte, 1832). Mar Freshwat Res 52, 531-541

Neer ZZ (2008)

Smith WD (2005) Life history aspects and population dynamics of a commercially exploited stingray, *Dasyatis dipterura* . MSc thesis in Mar Biol, San Francisco State Univ

Smith WD, Cailliet GM, Cortes ET (2008) Demography and elasticity of the diamond stingray, *Dasyatis dipterura*: parameter uncertainty and resilience to fishing pressure. Mar Freshwat Res 59, 575-586

Smith WD, Cailliet GM, Mellendez EM (2007) Maturity and growth characteristics of a commercially exploited stingray, *Dasyatis dipterura*. Mar Freshwat Res 58, 54-66

Snelson FF Jr., Roman BL, Burgess GH (2008) The reproductive biology of pelagic elasmobranchs. Pp 24-45 in (Camhi MD, Pikitch EK, Babcock EA, eds) Sharks of the Open Ocean: Biology, Fisheries and Conservation. Blackwell

Diodontidae

Lucano-Ramirez G, et al. (2011) Reproduction of the spiny puffer, *Diodon holocanthus* (Pisces: Diodontidae) in the continental shelf of Mexican Central Pacific. Rev Biol Trop 59, 217-232

Echineidae

Battaglia P, Potoschi A, Valastro M, Andaloro F, Romeo T (2016) Age, growth, biological and ecological aspects of *Remora osteochir* (Echineidae) in the Mediterranean Sea. J Mar Biol Assoc UK 96, 639-645

Echinorhinidae

Dawson CL, Starr RM (2000) Movements of subadult prickly sharks *Echinorhinus cookei* in the Monterey Canyon. Mar Ecol Progr Ser 386, 253-262

Eleotridae

Nordlie FG (2012) Life-history characteristics of eleotrid fishes of the western hemisphere, and perils of life in a vanishing environment. Rev Fish Biol Fish_22, 189-224

Elopidae

Adams AJ, Horodysky AZ, McBride RS, Guindon K, JShenker J, MacDonald TC, Harwel HD, Ward R, Carpenter K (2014) Global conservation status and research needs for tarpons (Megalopidae), ladyfishes (Elopidae) and bonefishes (Albulidae). Fish & Fish 15, 280-311

Blake C, Blake BF (1981) Age determination in six species of fish from a Mexican Pacific coastal lagoon. J Fish Biol 18, 471-478

Levesque JC (2015) Age, growth, and recruitment patterns of juvenile ladyfish (*Elops* sp) from the east coast of Florida (USA). Peer J, DOI_10.7717/peerj.1392

Embiotocidae

- Anderson RD, Bryan CF (1970) Age and growth of three surfperches (Embiotocidae) from Humboldt Bay, California. *Trans Am Fish Soc* 99, 475-482
- Anderson TW (2001) Predator responses, prey refuges, and density-dependent mortality of a marine fish. *Ecology* 82, 245-257
- Anderson TW, Sabado BD (1995) Correspondence between food availability and growth of a planktivorous temperate reef fish. *J Exp Mar Biol Ecol* 189, 65-76
- Anderson TW, Sabado BD (1999) Effects of kinship on growth of a live-bearing reef fish. *Mar Biol* 133, 115-121
- Baltz DM (1984) Life history variation among female surfperches (Perciformes : Embiotocidae). *Environ Biol Fish* 10, p159-171
- Baltz DM, Knight EE (1983) Age, growth, reproductive characteristics, and seasonal depth distribution of the spotfin surfperch, *Hyperprosopon anale*. *Calif Fish & Game* 69, 97-104
- Bane G, Robinson M (1970) Studies on the shiner perch, *Cymatogaster aggregata* Gibbons, in Upper Newport Bay, California. *Wasmann J Biol* 28, 259-268
- Banerjee T (1966) Survey of the fishes of Tomales Bay with notes on the life history of the white seaperch, *Phanerodon furcatus* Girard. MSc thesis, Univ Pacific
- Banerjee T (1971) Maturity and fecundity of the white seaperch (*Phanerodon furcatus*) from Tomales Bay, California. *J Fish Res Bd Can* 28, 1051-1053
- Banerjee T (1973) Age and rate of growth of the white seaperch (*Phanerodon furcatus*) from Tomales Bay, California. *Trans Am Fish Soc* 102, 448-450
- Bayer RD (1985) Shiner perch and Pacific staghorn sculpins in Yaquina Estuary, Oregon. *Northwest Sci* 59, 230-240
- Behrens DW (1977) Fecundity and reproduction of the viviparous perches *Hypsurus caryi* (Agassiz) and *Embiotoca jacksoni* Agassiz. *Calif Fish Game* 63, 234-252
- Bennett DE, Wydowski RS (1977) Biology of the redbtail surfperch (*Amphistichus rhodoterus*) from the central Oregon coast. *US Fish Wildl Serv, Tech Pap* 90, 23 p
- Bernardi G (2005) Phylogeography and demography of sympatric sister surfperch species, *Embiotoca jacksoni* and *E. lateralis* along the California coast: historical versus ecological factors. *Evolution* 59, 386-394
- Bernardi G, Bucciarelli G (1999) Molecular phylogeny and speciation of the surfperches (Embiotocidae, Perciformes). *Molec Phylog Evol* 13, 77-81

Brookins CG (1995) Population dynamics and reproductive ecology of the redbelt surfperch *Amphistichus rhodotus* (Embiotocidae). PhD in Fish Sci, OR State Univ

Bryant M (1978) Ecological studies of the striped perch *Embiotoca lateralis* and the kelp greenling *Hexagrammos decagrammus* in Carmel Bay, California. MSc thesis, Calif State Univ, Fresno

Carlisle JG Jr, Schott JW, Abramson NJ (1960) The barred surfperch (*Amphistichus argenteus*) in southern California. Calif Dept Fish & Game, Fish Bull 109, 79 p

Chu CT (1989) Functional design and prey capture dynamics in an ecologically generalized surfperch (Embiotocidae). J Zool 217, 417-440

Coleman SW, Jones AG (2011) Patterns of multiple paternity and maternity in fishes. Biol J Linn Soc 103, 735-760

Darling JDS, Noble ML, Shaw E (1980) Reproductive strategies in the surfperches. I. Multiple insemination in natural populations of the shiner perch, *Cymatogaster aggregata*. Evolution 34, 271-277

DeMartini EE, Moore TO, Plummer KM (1983) Reproductive and growth dynamics of *Hyperprosopon argenteum* (Embiotocidae) near San Diego, California. Env Biol Fish 8, 29-38

Eckmayer WJ (1979) Age and growth of four surfperches (Embiotocidae) from the Outer Harbor of Anaheim Bay, California. Calif Fish & Game 65, 265-272

Froeschke B, Allen LG, Pondella DJ II (2007) Life history and courtship behavior of black perch, *Embiotoca jacksoni* (Teleostomi: Embiotocidae), from southern California. Pac Sci 61, 521-531

Gnose CE (1967) Ecology of the striped seaperch, *Embiotoca lateralis*, in Yaquina Bay, Oregon. MSc thesis, Oregon State Univ, 53 pp

Goldberg SR (1978) Reproductive cycle of the white seaperch, *Phanerodon furcatus* (Embiotocidae). Copeia 1978(2), 335-338

Gordon CD (1965) Aspects of the age and growth of *Cymatogaster aggregata* Gibbons. MSc thesis, Univ British Columbia, 90 pp

Hubbs, CL (1921) The ecology and life history of *Amphigonopterus aurora* and other viviparous perches of California. Biol Bull 40, 181-209

rHubbs CL, Hubbs LC (1954) Data on the life history, variation, ecology and relationships of the kelp perch, *Brachyistius frenatus*, an embiotocid fish of the Californias. Calif Fish Game 40, 183-198

Isaacson PA, Isaacson DM (1966) Notes non the life history of the black perch, *Embiotoca jacksoni* Agassiz. *Tran Am Fish Socn* 95, 107-109

Izumiya M, Westphal MF, Crow, KD (2020) (1966) In the surf zone: reproductive strategy of the calico surfperch (*Amphistichus koelzi*) in a comparative context. *J Fish Biol* xx, yyy-yyy in press

Johnson DW, Freiwald J, Bernardi G (2015) Genetic diversity affects the strength of population regulation in a marine fish. *Ecology* 97, 627-639

Johnson DW, Stirling BS, Paz J, Satterfield DR (2019) Geographic variation in demography of black perch (*Embiotoca jacksoni*): Effects of density, food availability, predation, and fishing. *J Exp Mar Biol Ecol* 516, 16-24

LaPlante LH (2006) Length-specific brood size and winter parturition in pink seaperch (*Zalemibus rosaceus*) (Perciformes: Embiotocidae). *Fish Bull* 104, 623-625

LeBreque JR, Alva-Campbell YR, Archambeault S, Crow KD (2014) Multiple paternity is a shared reproductive strategy in the live-bearing surfperches (Embiotocidae) that may be associated with female fitness. *Ecol Evol* 4, 2316-2329

Liu J-X, Avise JC (2011) High degree of multiple paternity in the viviparous shiner perch, *Cymatogaster aggregata*, a fish with long-term female sperm storage. *Mar Biol* 158, 893-901

Longo GC (2016) Molecular evolution in embiotocid surfperches. PhD in Ecol Evol Biol, Univ Calif, Santa Cruz

Longo GC, Bernardi G, Lea RN (2015) Taxonomic revisions within Embiotocidae (Teleostei, Perciformes) based on molecular phylogenetics. *Zootaxa* 4482(3), 591–596

Ngoile MA (1978) Biology of the redbtail surfperch, *Amphistichus rhodoterus* in northern California. MSc thesis, Humboldt State Univ

Odenweller DB (1975) The life history of the shiner surfperch *Cymatogaster aggregata* Gibbons, in Anaheim Bay, California. *Calif Dep Fish Game, Fish Bull* 177, 107-115

Quaranta K (2011) Feeding morphology and kinematics in surfperches (Embiotocidae: Perciformes): evolutionary and functional consequences. MSc thesis in Mar Sci, Calif State Univ, Stanislaus

Reisser CMO, Beldade R, Bernardi G (2009) Multiple paternity and competition in sympatric congeneric reef fishes, *Embiotoca jacksoni* and *E. lateralis*. *Molec Ecol* 18, 1504-1510

Satterfield DR, Steele MA (2019) Effects of size and sex on the courting success and foraging behaviour of *Embiotoca jacksoni*. *J Fish Biol* 95, 472-479

- Schultz ET (1990) Daily otolith increments and the early life history of a viviparous fish, *Micrometrus minimus* (Embiotocidae). *Copeia* 1990(1), 59-67
- Schultz ET (1991) The effect of energy reserves on breeding schedule: is there a saturation point? *Func Ecol* 5, 819-824
- Schultz ET (1993a) Sexual size dimorphism at birth in *Micrometrus minimus* (Embiotocidae): A prenatal cost of reproduction. *Copeia* 1993(2), 456-462
- Schultz ET (1993b) The Effect of Birth Date on Fitness of Female Dwarf Perch, *Micrometrus minimus* (Perciformes: Embiotocidae). *Evolution* 47, nmr2, p520-539
- Schultz ET (2008) A sex difference in seasonal timing of birth in a livebearing fish. *Copeia* 2008(3), 673-679
- Schultz ET, Clifton LM, Warner RR (1991) Energetic constraints and size-based tactics: the adaptive significance of breeding-schedule variation in a marine fish (Embiotocidae: *Micrometrus minimus*) *Am Nat* 138, 1408-1430
- Schultz ET, Rountos PC (2001) Analysis of daily growth patterns in young-of-year male dwarf surfperch (Embiotocidae: *Micrometrus minimus*) suggests alternative tactics: breed or grow after birth. *Copeia* 2001(1), 14-24
- Shaw E, Allen J, (1977) Reproductive behaviour in the female shiner perch *Cymatogaster aggregata*. *Mar Biol* 40, 81-86
- Shaw E, Allen J, Stone R (1974) Notes on collection of shiner perch, *Cymatogaster aggregata*, in Bodega Harbor, California. *Calif Fish Game* 60, 15-22
- Sivalingham S (1953) Age and growth of *Taeniotoca lateralis*. MSc thesis, Univ Wash, 53 p
- Suomela AJ (1931) The age and growth of *Cymatogaster aggregata* Gibbons collected in Puget Sound, Washington. MSc thesis, Univ hWash, 43 p
- Swedberg SE (1965) Age-fecundity relationships in the striped seaperch *Embiotoca lateralis* from Yaquina Bay Oregon. MSc thesis, Oregon State Univ, 41 p
- Tarp FH (1952) A revision of the family Embiotocidae.(the surfperches). *Calif Dep Fish Game, Fish Bull* 88, 99 p
- Tofts AMS, Johnson DW, Carter AJR (2016) Strong nonlinear selection against fluctuating asymmetry in wild populations of a marine fish. *Evolution* 70, 2899-2908
- Tootell JS, Steele MA (2012) Factors affecting courtship success and behavior of a temperate reef fish, *Brachyistius frenatus*. *Bull So Calif Acad Sci* 111, 132-140

Wares PG (1971) Biology of the pile perch, *Rhacochilus vacca*, in Yaquina Bay, Oregon. US Bur Sport Fish Wildlf, Tech Pap 57

Warner RR, Harlan RK (1982) Sperm competition and sperm storage as determinants of sexual dimorphism in the dwarf surfperch, *Micrometrus minimus*. Evolution 36, 44-55

Westphal MF (2020) unpubl data—pers comm to EE DeMartini, April 20, 2020

Westphal MF, Izumiyama M, Crow KD (2020, unpubl draft ms) Fecundity of the calico surfperch *Amphistichus koelzi* (Teleostei: Embiotocidae). Northwest Sci

Westphal MF, Morey SR, Uyeda JC, Morgan TJ (2011) Molecular phylogeny of the subfamily Amphistichinae (Teleostei: Embiotocidae) reveals parallel divergent evolution of red pigmentation in two rapidly evolving lineages of sand-dwelling surfperch. J Fish Biol 79, 313-330

Wilson DC, Millemann RE (1969) Relationships of female age and size to embryo number and size in the shiner perch, *Cymatogaster aggregata*. J Fish Res Bd Can 26, 2339-2344

Woods PJ (2007) Habitat-dependent geographical variation in ontogenetic allometry of the shiner perch *Cymatogaster aggregata* Gibbons (Teleostei: Embiotocidae). J Evol Biol, 1783-1798

Wydoski RS, Bennett DE (1978) Contributions to the life history of the silver surfperch (*Hyperprosopon ellipticum*) from the Oregon coast. Calif Fish Game 59, 178-190

Engraulidae

Bayliff WH (1967) Growth, mortality, and exploitation of the Engraulidae, with special reference to the anchoveta, *Cetengraulis mysticetus*, and the colorado, *Anchoa naso*, in the Eastern Pacific Ocean. Bull Inter-Amer Trop Tuna Comm, 12(5), 365-432

Chiappa-Carrera X, Gallardo-Cabello M (1992) (1992) Tallas y otolitos en la determinación de la edad de la anchoveta *Engraulis mordax* (Pisces: Engraulidae). Rev Biol Trop, 40 (1), 1-5

Heath, KL (1980) Comparative life histories of two species of anchovies, *Anchoa delicatissima* and *A. compressa* (F. Engraulidae), from Newport Bay, California. MA thesis, Cal State Univ Fullerton. 71p

Howard GV, Landa A (1958) A study of the age, growth, sexual maturity, and spawning of the anchoveta (*Cetengraulis mysticetus*) in the Gulf of Panama. Bull Inter-Amer Trop Tuna Comm, 11(9), 365-432

Hunter JR, Macewicz BJ (1980) Sexual maturity, batch fecundity, spawning frequency, and temporal pattern of spawning for the northern anchovy, *Engraulis mordax*, during the 1979 spawning season. Calif Coop Oceanic Fish Invest Rep, 21, 139-149

Litz NC, Emmett RL, Heppell SS, Brodeur RD (2008) Ecology and distribution of the northern subpopulation of northern anchovy (*Engraulis mordax*) off the U.S. west coast. Calif Coop Oceanic Fish Invest Rep, 49, 167-182

MacCall AD (1973) The mortality rate of *Engraulis mordax* in Southern California. Calif Dep Fish Game, Tech Rep 1973(4), 23 p

Parrish RH, Mallicoate DL, Klingbeil RA (1986) Age-dependent fecundity, number of spawnings per year, sex ratio, and maturation stages of northern anchovy, *Engraulis mordax*. Fish Bull, 84, 503-

Parrish RH, Mallicoate DL, Mais KF (1985) Regional variations in the growth and age composition of northern anchovy, *Engraulis mordax*. Fish Bull, 83, 483-496

Pike GC (1951) Age, growth and maturity studies of the Pacific anchovy (*Engraulis mordax*) from the coast of British Columbia. MA thesis, Dep Zool, Univ Brit Columbia.

Spratt JD (1972) The use of otoliths to separate groups of northern anchovies. Calif Dep Fish Game, Tech Rep, 1972(1), 25 p

Epinephelidae

Balart EF, Gonzalez-Cabello A, Romero-Ponce RC, Zayas-Alvarez A, Calderon-Parra M, Campos-Davila L, Findley LT (2006) Length–weight relationships of cryptic reef fishes from the southwestern Gulf of California, Mexico. J appl Ichthyol 22, 316-318

Craig MT, Pondella DJ Jr, Hafner JC (1999) Analysis of age and growth in two Eastern Pacific groupers (Serranidae: Epinephelinae). Bull Mar Sci 65, 807-814

Morales-Nin B (1994) Growth of demersal fish species of the Mexican Pacific Ocean. Mar Biol 121, 211-217

Rodriguez-Romero S, Palacios-Salgado JDS, Lopez-Martinez J, Hernandez Vazquez S, Velazquez-Abunader JI (2009) The length–weight relationship parameters of demersal fish species off the western coast of Baja California Sur, Mexico. J appl Ichthyol 25, 114–116

Saenz-Arroyo ZZ, Roberts CM, Torre J, Carino-Olvera M (2005) Using fishers' anecdotes, naturalists' observations and grey literature to reassess marine species at risk: the case of the Gulf grouper in the Gulf of California, Mexico. Fish Fish 6, 121-133

Saenz-Arroyo A, Roberts CM, Torre J, Carino-Olvera M, Enriquez-Andrade RR (2005) Rapidly shifting environmental baselines among fishers of the Gulf of California. *Proc Roy Soc B* 272, 1957-1962

Exocoetidae

Ichimaru T, Nakazono A (1999) Ovarian development and spawning of flying fish, *Cypselurus heterurus doederleini*, in the northwest coastal water of Kyusyu. *Bull Jpn Soc Sci Fish* 10, 680-688

Gadidae

Andrews AH (2016) Lead–radium dating of Pacific cod (*Gadus macrocephalus*) validation of the young-fish scenario. *Mar Freshwat Res* 67, 1982-1986

Barbeaux S, Aydin K, Fissel B, Holsman K, Laurel B, Palsson W, et al (2019) Chapter 2: Assessment of the Pacific cod stock in the Gulf of Alaska. GOA SAFE Rep, No Pac Fish Manag Council

Bouchard C, Mollard S, Suzuki K, Robert D, Fortier L (2016) Contrasting the early life histories of sympatric Arctic gadids *Boreogadus saida* and *Arctogadus glacialis* in the Canadian Beaufort Sea). *Polar Biol* 39,1005-1022

Brown MS, Busby ML, Mier KL (2001) Walleye pollock *Theragra chalcogramma* during transformation from the larval to juvenile stage: otolith and osteological development. *Mar Biol* 139, 845-851

Datsky AV (2016) Biological features of the common fish species in Olyutorsky-Navarin region and the adjacent waters of the Bering Sea: 1. Gadidae (Cods) family. *J Ichthyol* 56, 868-889

Dorn MW, Deary AL, Fissel BE, Jones DT, Lauffenburger NE, Palsson WA, et al (2019) Chapter 1: Assessment of the walleye pollock stock in the Gulf of Alaska. GOA SAFE Rep, No Pac Fish Manag Council

Duffy-Anderson JT, Barbeaux SJ, Farley E, Heintz R, Horne JK, Parker-Stetter SL, Petrik C, Siddon EC, Smart T (2016) The critical first year of life of walleye pollock (*Gadus chalcogrammus*) in the eastern Bering Sea: implications for recruitment and future research. *Deep-Sea Res II, Topical Stud Oceanogr* 134, 283–301

Dunn JR, Matarese AC (1986) A review of the early life history of Northeast Pacific gadoid fishes. *Fish Res* 5, 163-184.

- Graham M, Hop H (1995) Aspects of reproduction and larval biology of Arctic cod (*Boreogadus saida*). Arctic 48(2), 130-135
- Helser TE, Colman JR, Anderl DM, Kestelle CR (2017) Growth dynamics of saffron cod (*Eleginus gracilis*) and Arctic cod (*Boreogadus saida*) in the Northern Bering and Chukchi Seas. Deep-Sea Research II 135, 66–77
- Hinckley S (1987) The reproductive biology of walleye pollock, *Theragra chalcogramma*, in the Bering Sea, with reference to spawning stock structure. Fish Bull 85, 481-498
- Hop H, Welch HE, Crawford RE (1997) Population structure and feeding ecology of Arctic cod schools in the Canadian High Arctic. Am Fish Soc Symp 19, 68-80
- Kestelle CR, Helser TE, McKay JL, Johnston CG, Anderl DM, Matta ME, Nicol DG (2017) Age validation of Pacific cod (*Gadus macrocephalus*) using high-resolution stable oxygen isotope (^{18}O) chronologies in otoliths. Fish Res 185, 43-53
- Laurel-BJ, Spencer M, Iseri P, Copeman LA (2016) Temperature-dependent growth and behavior of juvenile Arctic cod (*Boreogadus saida*) and co-occurring North Pacific gadids. Polar Biol 39, 1127-1135
- McQueen K, Eveson JP, Dolk B, Lorenz T, Mohr T, Schade FM, Krumme U (2019) Growth of cod (*Gadus morhua*) in the western Baltic Sea: estimating improved growth parameters from tag-recapture data. Can J Fish. Aquat. Sci. 76: 1326–1337
- Mueter FJ, Nahrgang J, Nelson RJ, Berge J (2016) The ecology of gadid fishes in the circumpolar Arctic with a special emphasis on the polar cod (*Boreogadus saida*). Polar Biol 39, 961-967
- Munk KM (2001) Walleye pollock otolith aging: comparison of the techniques used by two agencies. AK Dep Fish Game, Reg Info Rep No 5J01-06
- Nahrgang J, Storhaug E, Murzina SA, Delmas O, Nemova NN, Berge J (2016) Aspects of reproductive biology of wild-caught polar cod (*Boreogadus saida*) from Svalbard waters. Polar Biol 39, 1155-1164
- Narimatsu Y, Ueda Y, Okuda T, Hattori T, Fujiwara K, Ito M (2010) The effect of temporal changes in life-history traits on reproductive potential in an exploited population of Pacific cod, *Gadus macrocephalus*. ICES J Mar Sci 67, 1659-1666
- Neidetcher-SK, Hurst TP, Ciannelli L, Logerwell EA (2014) Spawning phenology and geography of Aleutian Islands and eastern Bering Sea Pacific cod (*Gadus macrocephalus*). Deep-Sea Res II 109, 204-214
- Nicol DG, Kotwicki S, Zimmermann M (2013) Diel vertical migration of adult Pacific cod *Gadus macrocephalus* in Alaska. J Fish Biol 83, 170-189

Ormseth OA, Norcross BL (2009) Causes and consequences of life-history variation in North American stocks of Pacific cod. ICES J Mar Sci 66, 349-357

Roberson NE, Kimura DK, Gunderson DR, Shimada AM (2005) Indirect validation of the age-reading method for Pacific cod (*Gadus macrocephalus*) using otoliths from marked and recaptured fish. Fish Bull 103, 153-160

Safronov SN (1986) Peculiarities of reproduction and principles of change in the fecundity of Pacific navaga, *Eleginus gracilis* (Gadidae). J Ichthyol 26, 59-86

Smirnova MA, Orlova SY, Orlov SM (2019) The population genetic organization of Pacific cod *Gadus macrocephalus* in the North Pacific based on microsatellite analysis. J Ichthyol 59, 555-565

Stahl JP, Kruse GH (2008) Spatial and temporal variability in size at maturity of walleye pollock in the Eastern Bering Sea. Trans Am Fish Soc 137, 1543-1557

Stark JW (2007) Geographic and seasonal variations in maturation and growth of female Pacific cod (*Gadus macrocephalus*) in the Gulf of Alaska and Bering Sea. Fish Bull 105, 396-407

Thompson GG, Spies IB, Palsson WA (2019) 2A. Assessment of the Pacific Cod Stock in the Aleutian Islands. NPFMC Bering Sea and Aleutian Islands. SAFE Rep. No Pac Fish Manag Council, Portland, OR

Thompson GG, Thorson JT (2019) 2. Assessment of the Pacific Cod Stock in the Eastern Bering Sea. NPFMC Bering Sea and Aleutian Islands. SAFE Rep. No Pac Fish Manag Council, Portland, OR

Williams BC-Kruse GH, Dorn MW (2016) Interannual and spatial variability in maturity of walleye pollock *Gadus chalcogrammus* and implications for spawning stock biomass estimates in the Gulf of Alaska. PLoS ONE 11(10), e0164797

Yasupov RR, Rakitina MV (2017) Reproduction, embryonic and early post-embryonic development of saffron cod *Eleginus gracilis* (Gadidae) from the Tauyskaya Guba Bay (northern Okhotsk Sea). Trans Pac Res Instit Fish Oceanogr 189, 52-66

Gasterosteidae

Baker JA, Foster SA, Heins DC, Bell MA, King RW (1998) Variation in female life-history traits among Alaskan populations of the threespine stickleback, *Gasterosteus aculeatus* L. (Pisces: Gasterosteidae). Biol J Linn Soc 63, 141-159

Baker JA, Heins DC, Foster SA, King RW (2008) An overview of life-history variation in female threespine stickleback. Behaviour 145, 579-602

DeFaveri J, Merila J (2013) Variation in age and size in Fennoscandian three-spined sticklebacks (*Gasterosteus aculeatus*). PLoS ONE 8(11), e80866

DeFaveri J, Shikano T, Merila J (2014) Geographic variation in age structure and longevity in the nine-spined stickleback (*Pungitius pungitius*). J Fish Biol 9,(7), e102660

Dufresne F, FitzGerald GJ, Lachance S (1990) Age and size-related differences in reproductive success and reproductive costs in threespine sticklebacks (*Gasterosteus aculeatus*). Behav Ecol 1, 140-147

Greenbank J, Nelson PR (1959) Life history of the threespine stickleback *Gasterosteus aculeatus* Linnaeus in Karluk Lake and Bare Lake, Kodiak Island, Alaska. Fish Bull 59, 537-559

Jones JW, Hynes HBN (1950) The age and growth of *Gasterosteus aculeatus*, *Pygosteus pungitius* and *Spinachia vulgaris*, as shown by their otoliths. J Anim Ecol 19, 59-73

Lefebvre R, Larsson S, Bystrom P (2011) A temperature-dependent growth model for the three-spined stickleback *Gasterosteus aculeatus*. J Fish Biol 79, 1815-1827

McPhail JD (1977) Inherited interpopulation differences in size at first reproduction in threespine stickleback, *Gasterosteus aculeatus* L. Heredity 38, 53-60

Rollins JL, Cghiang P, Waite JN, VonHippel FA, Bell MA (2017) Jacks and jills: alternative life-history phenotypes and skewed sex ratio in anadromous threespine stickleback (*Gasterosteus aculeatus*). Evol Ecol Res 18, 363-382

Ruiz-Campos G, Contreras-Banderas S, De Lourdes Lozano-Vilano M, Gonzalez-Guzman S, Alaniz-Garcia J (2000) Ecological and distributional status of the continental fishes of northwestern Baja California, Mexico. Bull So Calif Acad Sci 99, 59-90

Ruiz-Campos-G. Gonzalez Acosta AF, De La Cruz Aguero J (2006) Length–weight and length–length relationships for some continental fishes of northwestern Baja California, Mexico. J appl Ichthyol 22, 314-316

Singam AR, MacColl ADC (2018) Otolith development in wild populations of stickleback: Jones & Hynes method does not apply to most populations. J Fish Biol 93, 272-281

Snyder RJ (1991a) Quantitative genetic analysis of life histories in two freshwater populations of the threespine stickleback. Copeia 1991(2), 526-529

Snyder RJ (1991b) Migration and life histories of the threespine stickleback: evidence for adaptive variation in growth rate between populations. Environ Biol Fish 31, 381-388

von_Hippel FA, Smayda LC, Zimmerman CE, Bell MA (2013) Validation of daily growth increments in otoliths to age threespine stickleback (*Gasterosteus aculeatus*). Evol Ecol Res 15, 947-957

Wright PJ, Huntingford FA (1993) Daily growth increments in the otoliths of the three-spined stickleback, *Gasterosteus aculeatus* L. J Fish Biol 42, 65-77

Wright HA, Wootton RJ, Barber I (2004) Interpopulation variation in early growth of threespine sticklebacks (*Gasterosteus aculeatus*) under laboratory conditions. Can J Fish Aquat Sci 61, 1832-1838

Yurtseva A, Anastasia Yurtseva^{1,2} | Kristina Noreikiene K, Lajus D, Li Z, Alapassi T, Ivanova T, Ivanov M, Golovin P, Vesala S, Merilä J (2019) Aging three-spined sticklebacks *Gasterosteus aculeatus*: comparison of estimates from three structures. J Fish Biol 95, 802-811

Gempylidae

Keller HR, Kerstetter DW (2014) Length–length and length–weight relationships of oilfish (*Ruvettus pretiosus*), escolar (*Lepidocybium flavobrunneum*), snake mackerel (*Gempylus serpens*), and longnose lancetfish (*Alepisaurus ferox*) from the Gulf of Mexico and the western North Atlantic Ocean. J appl Ichthyol 30, 242-243

Maximov VP (1970) Some data on the biology of *Lepidocybium bruneum*. Voprosy Iktiologii 10, 50-57

Nakamura I, Parin NV (1993) Snake mackerels and cutlassfishes of the World. FAO Species Catalogue 125, 1–129

Gerreidae

Cota-Gomez VM, DeLaCruz-Aguero G, Garcia-Rodriguez FJ, DeLaCruz-Aguero J (2013) Length-weight relations for 14 fish species of La Paz Bay, Baja California Sur, Mexico. Acta Ichthyol Piscat 43, 297-298

Gonzalez-Acosta AF, DeLaCruz-Aguero G, DeLaCruz-Aguero J (2004) Length–weight relationships of fish species caught in a mangrove swamp in the Gulf of California (Mexico). J appl Ichthyol 20, 154-155

Lopez-Martinez-et al (2011) Population parameters of the Pacific flagfin mojarra *Eucinostomus currani* (Perciformes: Gerreidae) captured by shrimp trawling fishery in the Gulf of California. Rev Biol Trop 59, 887-897

Gobiidae

Behrems KC(1983) The comparative ecology and interactions between two sympatric gobies (*Lythrypnus dalli* and *Lythrypnus zebra*). PhD thesis, Univ Calif, Los Angeles, 343p

Forrester GE, Fredericks BI, Gerdeman D, Evans B, Steele MA, Zayed K, Schweitzer LE, Suffet IH Ric Vance RR, Ambrose RF (2003) Growth of estuarine fish is associated with the combined concentration of sediment contaminants and shows no adaptation or acclimation to past conditions. *Mar Environ Res* 56, 423-442

GonzalezAcosta AF, DeLaCruzAguero G, DeLaCruzAguero J (2004) Length–weight relationships of fish species caught in a mangrove swamp in the Gulf of California (Mexico). *J appl Ichthyol* 20, 154-155

Gregor CA, Anderson TW (2016) Relative importance of habitat attributes to predation risk in a temperate reef fish. *Environ Biol Fish* 99, 539-556

Grossman GD (1979) Demographic characteristics of an intertidal bay goby (*Lepidogobius lepidus*). *Environ Biol Fish*, 4, 207-218

Hoshino N, Kinoshita T, Kanno Y (1993) Age, growth and ecological characteristics of goby, *Acanthogobius flavimanus* in Hakodate Bay Hokkaido Japan. *Bull Fac Fish Hokkaido Univ* 44, 147-157

Kroon FJ, deGraf M, Liley NR (2000) Social organisation and competition for refuges and nest sites in *Coryphopterus nicholsii* (Gobiidae), a temperate protogynous reef fish. *Environ Biol Fish*, 57, 401-411

McGourty K, Kinzinger A, Goldsmith G (2008) Spawning time, fecundity, habitat utilization, and parasites of a Northern California population of tidewater goby, *Eucyclogobius newberryi*. *California Fish and Game*, 94, 18-32

Prasad RR(1959) Reproduction in *Clevelandia ios* (JordanRDAL and Gilbert), with an account of the embryonic and larval development. *Proc Nat Inst Sci India* 25B(1), 13-20

Spies BT, Tarango BC, Steele MA (2014) Larval duration, settlement, and larval growth rates of the endangered tidewater goby (*Eucyclogobius newberryi*) and the arrow goby (*Clevelandia ios*) (Pisces, Teleostei). *Bull So Calif Acad Sci*, 113, 165-175

Steele MA(1997) The relative importance of processes affecting recruitment of two temperate reef fishes. *Ecology* 78, 129-145

Swenson RO(1999) The ecology, behavior, and conservation of the tidewater goby, *Eucyclogobius newberryi*. *Environ Biol Fish* 55, 99-114

Yong L, Grober MS (2014) Sex differences in the energetic content of reproductive tissues in the blackeye goby, *Rhinogobiops nicholsii*. *Environ Biol Fish* 97, 321-328

Gonostomatidae

Andrianov DP, Bekker VE (1989) Reproductive biology of mesopelagic fishes of the eastern Pacific. J Ichthyol 29, 43-48

DeLaCruz-Aguero J, Cota-Gomez VM (2006) Length–weight relationships of 10 deep-sea fish species from the Mexican Pacific Ocean. J appl Ichthyol 22, 319-321

Gymnuridae

Bizzarro JJ (2005) Fishery biology and feeding ecology of rays in Bahia Almejas, Mexico MSci thesis, Mar Sci, San Francisco State Univ, 468 p

Hemitripteridae

Burge RT, Schultz SA (1973) The Marine Environment in the Vicinity of Diablo Cove with Special Reference to Abalones and Bony Fishes. Calif Dep Fish Game, Mar Res Tech Rep No 19, 433 p

Kolpakov NV, Dolganova NT (2006) On the biology of *Blepsias cirrhosis* (Hemitripteridae) from coastal waters of Northern Primorye. J Ichthyol 46, 454-459

Spies I, Nichol D, Aydin K, TenBrink TT (2017) Assessment of the sculpin stock complex in the Gulf of Alaska NMFS, NOAA, Gulf of Alaska SAFE Rep 28 p

Spies I, Nichol D, TenBrink TT (2016) Assessment of the sculpin stock complex in the Gulf of Alaska. No Pac Fish Manag Council, Gulf of Alaska SAFE Rep, 6 p

TenBrink TT, Aydin KY (2009) Life history traits of sculpins in the eastern Bering Sea and Aleutian Islands. No Pac Res Bd Finl Rep 628, 215 p

Hemiramphidae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Heterodontidae

Garcia-Rojas MG, Villavicencio-Garayzar C (2003) Reproductive biology of the horned shark *Heterodontus francisci*, Girard, 1854 (Chondrichthyes:Heterodontidae), in Lagoon San Ignacio, Baja California Sur, Mexico. MARUBA 6, Congreso de Ciencias del Mar, Havana Cuba

Nelson DR, Johnson RH (1970) Diel activity rhythms in the nocturnal, bottom-dwelling sharks, *Heterodontus francisci* and *Cephaloscyllium ventriosum*. Copeia 1970(4), 732-739

Hexagrammidae

Antonenko DV (1999) Reproduction of masked greenling *Hexagrammos octogrammus* in Peter the Great Bay, Japan Sea. Russ J Mar Biol 25, 90-91

Barker MW (1980) Population and fishery dynamics of recreationally exploited marine bottomfish of northern Puget Sound. PhD thesis, Univ Washington, 134 pp

Beaudreau AH, Essington TE (2011) Use of pelagic prey subsidies by demersal predators in rocky reefs insight from movement patterns of lingcod. Mar Biol 158, 471-483

Berger AM, Arnold L, Rodomsky BT (2015) Status of kelp greenling (*Hexagrammos decagrammus*) along the Oregon coast in 2015. ID312, Final Product, Kelp Greenling Assessment, OR Dep Fish Wildl

Bryant M (1978) Ecological studies of the striped perch *Embiotoca lateralis* and the kelp greenling *Hexagrammos decagrammus* in Carmel Bay, California. MSc thesis, Calif State Univ, Fresno, 62 p

Canino MF, Spies IB, Guthridge JL, Hollowed MM (2010) Genetic assessment of the mating system and patterns of egg cannibalism in atka mackerel. Mar Coast Fish Dynam Manag Ecosys Sci 2, 388-398

Cooper DW, McDermott SF, Ianelli JN (2010) Spatial and temporal variability in atka mackerel female maturity at length and age. Mar Coast Fish Dynam Manag Ecosys Sci 2, 329-338

Cope JM, MacCall AD (2005) Status of kelp greenling (*Hexagrammos decagrammus*) in Oregon and California waters as assessed in 2005, Final Status Rep KG05. No Pac Fish Manag Council, Portland OR

Crow KD (1995) The reproductive biology of kelp greenling *Hexagrammos decagrammus*. MSc thesis, San Francisco State Univ

Crow KD, Munehara H, Bernardi G (2010) Sympatric speciation in a genus of marine reef fishes. Molec Ecol 19, 2089-2105

Crow-KD, Powers DA, Bernardi G (1997) Evidence for multiple maternal contributors in nests of kelp greenling (*Hexagrammos decagrammus* (Hexagrammidae). Copeia 1997(1) 9-15

Frid A, Marliave J, Heithaus MR (2012) Interspecific variation in life history relates to antipredator decisions by marine mesopredators on temperate reefs. PLoS ONE 7(6), e0040083 doi10.1371/journal.pone.0040083

Gomelyuk VE, Leunov VP (1999) Changes in age-specific behaviour and spatial structure of masked greenling *Hexagrammos octogrammus*, (Hexagrammidae). Environ Biol Fish 54, 313-323

Greenley A, Green K, Starr RM (2016) Seasonal and ontogenic movements of lingcod (*Ophiodon elongatus*) in Central California, with implications for marine protected area management. Calif Coop Oceanic Fish Invest Rep 57, 71-88

Haltuch MA, Wallace J, Allen C, Akselrud CA, Nowlis J, Barnett LAK, Valero JL, Tsou T-S, Lam L (2018) 2017 Lingcod Stock Assessment. No Pac Fish Manag Council, Portland, OR

Hanan DA, Curry BE (2012) Long-term movement patterns and habitat use of nearshore groundfish: tag-recapture in Central and Southern California waters. Open Fish Sci J 5, 30-43

Jones MK, Mulligan T (2014) Juvenile rockfish recruitment in Trinidad Bay, California. Trans Am Fish Soc 143, 543-551

Kendall AW Jr, Vinter B (1984) Development of hexagrammids (Pisces: Scorpaeniformes) in the Northeastern Pacific Ocean. NOAA Tech Rep NMFS 2, 44 p

Lam LS (2019) Geographic and habitat-based variation in lingcod (*Ophiodon elongatus*) demography and life history along the U.S. west coast. MSc thesis, Calif State Univ, Monterey Bay

Lam LS, Basnett BL, Haltuch MA, Cope J, Andrews K, Nichols KM, Longo GC, Samhour JF, Hamilton SL (2021) Geographic variability in lingcod *Ophiodon elongatus* life history and demography along the US West Coast: oceanographic drivers and management implications. Mar Ecol Progr Ser 670, 203-222

Lauth RR, Guthridge JL, Cooper DWZ, McEntire SW (2010) Behavioral ecology of color patterns in atka mackerel. Mar Coast Fish Dynam Manag Ecosys Sci 2, 399-411

Lauth RR, Guthridge J, Nicol D, McEntire SW, Hillgruber N (2007) Timing and duration of mating and brooding periods of Atka mackerel (*Pleurogrammus monopterygius*) in the North Pacific Ocean. Fish Bull 105, 560-570

Longo GC, Lam L, Basnett B, Samhour J, Hamilton S, Andrews K, Williams G, Goetz G, McClure M, Nichols KM (2020) Strong population differentiation in lingcod (*Ophiodon*

elongatus) is driven by a small portion of the genome. *Evol Applic*, <https://doi.org/10.1111/eva.130377>

Markevitch AI (2004) Parental behavior of the male greenlings *Hexagrammos otakii* and *H. octogrammus* (Hexagrammidae). *J Ichthyol* 44, 521

Markevitch AI (2011) Utilization of the “nesting” site territory and relationships with conspecific Individuals during parental care in the males of masked greenling *Hexagrammos octogrammus* (Hexagrammidae, Scorpaeniformes). *J Ichthyol* 51, 543-550

Matthews KR (1992) A telemetric study of the home ranges and homing routes of lingcod *Ophiodon elongatus* on shallow rocky reefs off Vancouver Island, British Columbia. *Fish Bull* 90, 784-790

Matyushin VM (1992) Growth and production of masked greenlings in seagrass in Vityaz Bay (Sea of Japan). *Russ J Mar Biol* 17, 33-37

McDermott SF, Cooper DW, Guthridge JL, Spies IB, Canino MF, Woods P, Hillgruber N (2011) Effects of maternal growth on fecundity and egg quality of wild and captive Atka mackerel. *Mar Coast Fish Dynam Manag Ecosys Sci* 3, 321-335

McDermott SF, Fritz LW, Haist V (2005) Estimating movement and abundance of Atka mackerel (*Pleurogrammus monopterygius*) with tag–release–recapture data. *Fish Oceanogr* 14, suppl 1, 113-130

McDermott SF, Lowe SA (1997) The reproductive cycle and sexual maturity of Atka mackerel, *Pleurogrammus monopterygius*, in Alaska waters. *Fish Bull* 95, 321-333

McDermott SF, Maslenikov KP, Gunderson DR (2007) Annual fecundity, batch fecundity, and oocyte atresia of Atka mackerel (*Pleurogrammus monopterygius*) in Alaskan waters. *Fish Bull* 105,19-29

Morita SH, Morita K, Nishimura A (2012) Sex-biased dispersal and growth in sablefish (*Anoplopoma fimbria*) in the northeastern Pacific Ocean. *Environ Biol Fish* 94, 505-511

Munehara H, Kanamoto Z, Miura T (2000) Spawning behavior and interspecific breeding in three Japanese greenlings (Hexagrammidae). *Ichthyol Res* 47, 287-292

Munehara H, Shimazaki K (1989) Annual maturation changes in the ovaries of masked greenling *Hexagrammos octogrammus*. *Nipp Suisan Gakkaishi* 55, 423-429

Munk KM (2001) Maximum ages of groundfishes in waters off Alaska and British Columbia and considerations of age determination. *AK Fish Res Bull* 8, 12-21

Orlov AM, Zolotov OG (2010) Distribution and some features of biology of rock greenling *Hexagrammos lagocephalus* in Pacific waters off the Northern Kuril Islands and Southeastern Kamchatka. J Ichthyol 50, 174-187

Rand KM, Beauchamp DA, Lowe SA (2010) Longitudinal growth differences and the influence of diet quality on Atka mackerel of the Aleutian Islands, Alaska: using a bioenergetics model to explore underlying mechanisms. Mar Coast Fish Dynam Manag Ecosys Sci 2, 362-374

Rodonsky BT, Kautzi LA, Hannah RW, Good CD (2015) Kelp greenling (*Hexagrammos decagrammus*) growth, spawning seasonality, and female length at maturity based on histological evaluation of ovaries from Oregon waters. OR Dep Fish Wildlf, Info Rep, 2015-02

Shestakov AV, Nazarkin MV (2006) On the biology of the white-spotted greenling *Hexagrammos stelleri* and the masked greenling *H. octogrammus* (Hexagrammidae) from Taii Bay of the Sea of Okhotsk. J Ichthyol 46, 677-680

Silberberg KR, Laidig TE, Adams PB, Albin D (2001) Analysis of maturity in lingcod, *Ophiodon elongatus*. Calif Fish Game 87,139-152

Simenstad CA, Mayer RM, Nakatani RE (1978) Nearshore fish and macroinvertebrate communities of Attu Island, Alaska. Ann Rep, Jan-Dec 1977, for Fish Wildlf Serv, Nat Fish Wildlf Lab, by Fish Res Instit, Coll Fish, Univ Wash, Seattle, Wash

Starr RM, OConnell V, Ralston S (2004) Movements of lingcod (*Ophiodon elongatus*) in southeast Alaska: potential for increased conservation and yield from marine reserves. Can J Fish Aquat Sci 61, 1083-1094

Tolimieri N, Andrews K, Williams G, Katz S, Levin PS (2009) Home range size and patterns of space use by lingcod, copper rockfish and quillback rockfish in relation to diel and tidal cycles. Mar Ecol Progr Ser 380, 229-243

Vdovin AN, Antonenko DV (1998) Growth and age of *Hexagrammos octogrammus* of Peter the Great Bay (the Sea of Japan). J Ichthyol 38,81-85

VonBiela VR, Kruse GH, Mueter FJ, Black BA, Douglas DC, Helser TE, Zimmerman CE (2015) Evidence of bottom-up limitations in nearshore marine systems based on otolith proxies of fish growth. Mar Biol 162, 1019-1031

Wonsettler AL, Webb JF (1997) Morphology and development of the multiple lateral line canals on the trunk in two species of *Hexagrammos* (Scorpaeniformes, Hexagrammidae). J Morph 233, 195-214

Hexanchidae

Ebert DA (1984) Aspects of the life history of California's two cowshark species, *Notorhynchus cepedianus* and *Hexanchus griseus*. MSc thesis in Biol, San Jose State Univ

Ebert DA (1986a) Biological aspects of the sixgill shark, *Hexanchus griseus*. Copeia 1986(1) 151-155

Ebert DA (1986b) Aspects on the biology of hexanchid sharks along the California coast. 2nd Indo-Pac Fish Conf, 437-449

Ebert DA (1989) Life history of the sevengill shark, *Notorhynchus cepedianus* Peron, in two Northern California bays. Calif Fish Game 75, 102-112

Ebert DA (1996) Biology of the sevengill shark *Notorynchus cepedianus* (Peron, 1807) in the temperate coastal waters of southern Africa So Afr J Mar Sci 17, 93-103

Ebert DA (2002) Some observations on the reproductive biology of the sixgill shark *Hexanchus griseus* (Bonnaterre, 1788) from southern African waters. So Afr J Mar Sci 24, 359-363

King JR, Surry M (2017) Seasonal and daily movements of the bluntnose sixgill shark (*Hexanchus griseus*) in the strait of Georgia from satellite tag data Environ Biol Fish 100, 1543-1559

Smith SE, Au DW, Show C (1998) Intrinsic rebound potentials of 26 species of Pacific sharks. Mar Freshwat Res 49, 663-678

Van Dykhuizen G, Mollett HF (1992) Growth, Age Estimation and Feeding of Captive Sevengill Sharks, *Notorynchus cepedianus*, at the Monterey Bay Aquarium. Aust J Mar Freshwat Res 43, 297-318

Williams GD, Andrews KS, Farrer DA, Bargmann GG, Levin PS (2011) Occurrence and biological characteristics of broadnose sevengill sharks (*Notorynchus cepedianus*) in Pacific Northwest coastal estuaries. Environ Biol Fish 91, 379-3880

Icosteidae

Allen GH (2001) The ragfish, *Icosteus aenigmaticus* Lockington, 1880: a synthesis of historical and recent records from the North Pacific Ocean and the Bering Sea. Mar Fish Rev 63, 1-3

Istiophoridae

Alvarado-Castillo R, Felix-Araga R (1998) Crecimiento de *Istiophorus platypterus* (Pisces: Istiophoridae) en la boca del Golfo de California. Rev Biol Trop 46, 115-119

Kopf RK, Davie PS, Bromhead D, Pepperell JG (2011) Age and growth of striped marlin (*Kajikia audax*) in the Southwest Pacific Ocean ICES J Mar Sci 68, 1884-1895

Kume S, Joseph J (1969) Size composition and sexual maturity of billfish caught by the Japanese longline fishery in the Pacific Ocean E of 130° degrees W. Bull Far Seas Res Lab, No 2, 115-162

Lee HH, Piner KR, Hinton MG, Chang Y-J (2014) Sex-structured population dynamics of blue marlin *Makaira nigricans* in the Pacific Ocean. Fish Sci 80, 869-878

Mello-Barrera FZ et al (2003) Growth and length-weight relationship of the striped marlin, *Tetrapturus audax* (Pisces: Istiophoridae), in Cabo San Lucas, Baja California Sur, Mexico. Ciencias Marinas 29, 305-313

Ramirez-Perez ZZ, Casimiro Quiñonez-Velázquez C, Abitia-Cardenas LA, etc (xxxx) Age and growth of sailfish *Istiophorus platypterus* (Shaw in Shaw and Nodder, 1792) from Mazatlan, Sinaloa, Mexico.

Felipe N. Melo-Barrera FN (2011) Age and growth of sailfish *Istiophorus platypterus* (Shaw in Shaw and Nodder, 1792) from Mazatlan, Sinaloa, Mexico. Environ Biol Fish 92, 187-196

Skillman RA, Yong MYY (1976) Von Bertalanffy growth curves for striped marlin, *Tetrapturus audax*, and blue marlin, *Makaira nigricans*, in the central North Pacific Ocean. Fish Bull 79, 553-566

Sun C-L, Chang Y-J, Tszeng C-C, Yeh S-Z, Su NJ (2009) Reproductive biology of blue marlin (*Makaira nigricans*) in the western Pacific Ocean. Fish Bull 107, 420-432

Sun C-L, Chang S-Y, Chang YJ, Yeh S-Z (2011) Reproductive biology of female striped marlin (*Kajikia audax*) in the waters off Taiwan (preliminary). ISC/11/BILLWG-1/11. Interim Scientific Committee (ISC) Billfish Working Group Workshop, 19-27 January 2011, Honolulu, Hawaii

Su N-J, Sun C-L, Punt AE, Yeh S-Z, Chiang WC, Chang Y-J, Chang H-Y (2013) Effects of sexual dimorphism on population parameters and exploitation ratios of blue marlin (*Makaira nigricans*) in the northwest Pacific Ocean. Aquat Liv Res 26, 19-24

Sun C-L, Liu C-H, Yeh S-Z (2013) Age and growth of black marlin (*Makaira indica*) in the waters off eastern Taiwan. WCPFC-SC3-BI SWG/WP-2. Western Central Pac Fish Comm, Sci Comm Third Reg Sess, 13-24 August 2007, Honolulu, HI, USA

Sun C-L, Su N-J, Yeh S-Z, Chang Y-J (2013) Sex-specific growth parameters and natural mortality rates for blue marlin (*Makaira nigricans*) in the Northwest Pacific Ocean. ISC/13/BILLWG-1/10. ISC Billfish Working Group Workshop, 16-23 January 2013, Honolulu, HI, USA

Thomas LR, Humphries RL Jr, Chang Y-J, Sun C-L (2011) Review of life history parameters for blue marlin *Makaira nigricans*. ISC/13/BILLWG-1/12. Interim Scientific Committee (ISC) Billfish Working Group Workshop, 19-27 January 2011, Honolulu, Hawaii

Kyphosidae

Bredvik JJ, Boerger C, Allen LG (2011) Age and growth of two herbivorous, kelp forest fishes, the opaleye (*Girella nigricans*) and halfmoon (*Medialuna californiensis*). Bull. So Calif Acad. Sci 110, 25–34

Stevens EG, Watson W, Moser HG (1989) Development and distribution of larvae and pelagic juveniles of three kyphosid fishes (*Girella nigricans*, *Medialuna californiensis*, and *Hermosilla azurea*) off California and Baja California. Fish Bull 87, 745-768

Terry A, -Bucciarelli G, -Bernardi G (2000) Restricted gene flow and incipient speciation in disjunct Pacific Ocean and Sea of Cortez populations of a reef fish species, *Girella nigricans*. Evolution 52, 652-659

White GE, Brown C (2013) Site fidelity and homing behaviour in intertidal fishes. Mar Biol 160, 1365-1372

Labridae

Adreani MS, Allen LG (2008) Mating system and reproductive biology of a temperate wrasse, *Halichoeres semicinctus*. Copeia 2008(2), 467-475

Adreani MS, -Erisman BE, Warner RR (2004) Courtship and spawning behavior in the California sheephead, *Semicossyphus pulcher* (Pisces: Labridae) Environ Biol Fish 71, 13-19

Caselle JZ, Hamilton SL, Schroeder DM, Love MS, Standish JD, Rosales-Casian JA, Sosa-Nishizaki O (2011) Geographic variation in density, demography, and life history traits of a harvested, sex-changing, temperate reef fish. Can J Fish Aquat Sci 68, 288-303

Cowen RK (1985) Large scale pattern of recruitment by the labrid, *Semicossyphus pulcher*: causes and implications. J Mar Res 43, 719-742

Cowen RK (1990) Sex change and life history patterns of the labrid, *Semicossyphus pulcher*, across an environmental gradient. Copeia 1990(3), 787-795

Dunn RP (2016) Tool use by a temperate wrasse, California sheephead *Semicossyphus pulcher*. J Fish Biol 88, 805-810

Hamilton SL, Caselle JE, Standish JD, Schroeder DM, Love MS, Rosales-Casian JA, Sosa-Nishizaki O (2007) Size-selective harvesting alters life-histories of a temperatye sex-changing fish. *Ecol Applic* 17, 2268-2280

Loke-Smith KA, Floyd AJ, Lowe CG, Hamilton SL, Caselle JE, Young KA (2012) Reassessment of the fecundity of California sheephead. *Mar Coast Fish Dynam Manag Ecosys Sci* 4, 599-604

Victor BC (1986) Duration of the planktonic larval stage of one hundred species of Pacific and Atlantic wrasses (family Labridae). *MarBiol* 90, 317-326

Walters K (2002) A geographic comparison of life histories in two species of fish. MSc thesis, San Francisco State Univ

Labrisomidae

Galvan-Villa CM, Hastings PA (2018) Mating behavior and rapid concealing of nuptial coloration in males of the fishgod blenny *Malacoctenus ebisui* (Teleostei: Blenniiformes). *Environ Biol Fish* 101, 735-739

Lamnidae

Bruce BD (2008) The biology and ecology of the white shark, *Carcharodon carcharias*. Pp 69-81 in (Camhi MD, Pikitch EK, Babcock EA, eds) *Sharks of the Open Ocean: Biology, Fisheries and Conservation*. Blackwell

Cailliet GM, Bedford DW (1983) The biology of three pelagic sharks from California waters and their emerging fisheries: a review. *Calif Coop Oceanic Fish Invest Rep* 24, 57-69

Goldman KJ, Musick JA (2008) The biology and ecology of the salmon shark, *Lamna ditropis*. Pp 95-104 in (Camhi MD, Pikitch EK, Babcock EA, eds) *Sharks of the Open Ocean: Biology, Fisheries and Conservation*. Blackwell

Ribot-Carballal MC, Galvan-Magana F, Quinonez-Velazquez, C (2005) Age and growth of the shortfin mako shark, *Isurus oxyrinchus*, from the western coast of Baja California Sur, Mexico. *Fisheries Research* 76, 14-21

Semba Y, Nakano H, Aoki I (2009) Age and growth analysis of the shortfin mako, *Isurus oxyrinchus*, in the western and central North Pacific Ocean. *Environ Biol Fish* 84, 377-391

Stevens JD (2008) The biology and ecology of the shortfin mako shark, *Isurus oxyrinchus*. Pp 87-94 in (Camhi MD, Pikitch EK, Babcock EA, eds) *Sharks of the Open Ocean: Biology, Fisheries and Conservation*. Blackwell

Lampridae

Hyde, J., Underkoffler K, Sundberg M (2014) DNA barcoding provides support for a cryptic species complex within the globally distributed and fishery important opah (*Lampris guttatus*). *Molec Ecol Resources* 14, 1239–1247

Underkoffler K, Luers MA, -Hyde JR, Craig MT (2018) A taxonomic review of *Lampris guttatus* (Brünnich 1788) (Lampridiformes; Lampridae) with descriptions of three new species. *Zootaxa* 4437(4), 551-565

Liparidae

Glubokov AI (2010) The data on *Careproctus furcellus* and *C. rastrinus* (Liparidae) from the Olyutorskii Gulf of the Bering Sea: size composition, indices of organs, and diet. *J Ichthyol* 50, 52-64

Johnson CR (1969) Contributions to the biology of the showy snailfish, *Liparis pulchellus* (Liparidae). *Copeia*, 1969, 830-835

Mori T, Fukuda K, Ohtsuka S, Yamauchi S, Yoshinaga T (2022) Reproductive behavior and alternative reproductive strategy in the deep-sea snailfish, *Careproctus pellucidus*. *Mar Biol* 169,42, <https://doi.org/10.1007/s00227-022-04028-9> xx-xx

Orlov A, Binohlan C (2009) Length–weight relationships of deep-sea fishes from the western Bering Sea. *J appl Ichthyol* 25, 223-227

Orlov AM, Tokranov AM (2011) Some rare and insufficiently studied snailfish (Liparidae, Scorpaeniformes, Pisces) in the Pacific waters off the Northern Kuril Islands and Southeastern Kamchatka, Russia. *Internat Scholar Res Network (ISRN), Zool*, article ID341640, 12p

Stein(1980) Aspects of reproduction of liparid fishes from the Continental Slope and Abyssal Plain off Oregon, with notes on growth. *Copeia* 1980, 687-699

Stein DL, Drazen JC, Schlining KL, Barry JP, Kuntz L (2006) Snailfishes of the central California coast: video, photographic and morphological observations. *JFishBiol* 69, 970-986

Tokranov AM (2000) Length-age composition of liparid fishes (Liparidae) in the Pacific waters of the south-eastern Kamchatka and the North Kuril Islands. *Voprosy iktiologii* 40, 347-352

Lutjanidae

Garcia-Contreras OE, Quinonez-Velazquez C, Moran-Angulo RE, Valdez-Pineda MC (2009) Age, growth, and age structure of Amarillo snapper off the coast of Mazatlan, Sinaloa, Mexico. *No Am J Fish Manag* 29, 223-230

Gonzalez-Acosta AF, DeLaCruz-Aguero G, DeLaCruz-Aguero J (2004) Length–weight relationships of fish species caught in a mangrove swamp in the Gulf of California (Mexico). *J appl Ichthyol* 20, 154-155

Hernandez-AlvarezC, Natalia J. Bayona-Vasquez NJ, Dominguez-Dominguez O, Uribe-Alcocer M, Diaz-Jaimes_P (2020) Phylogeography of the Pacific red snapper (*Lutjanus peru*) and spotted rose snapper (*Lutjanus guttatus*) in the inshore tropical Eastern Pacific. *Copeia* 108(1), 61-71

Lucano-Ramirez G, Ruiz-Ramírez S, González-Sansón G, Ceballos-Vázquez BP (2014) Reproductive biology of the yellow snapper, *Lutjanus argentiventris* (Pisces, Lutjanidae), from the Mexican central Pacific. *Ciencias Marinas* 40, 33-44

Pinon A, Amezcua F, Duncan N (2009) Reproductive cycle of female yellow snapper *Lutjanus argentiventris* (Pisces, Actinopterygii, Lutjanidae) in the SW Gulf of California: gonadic stages, spawning seasonality and length at sexual maturity *J_appl_Ichthyol_25*,p18-25

Rocha-Olivares A (1998) Age, growth, mortality, and population characteristics of the Pacific red snapper, *Lutjanus peru*, off the southeast coast of Baja California, Mexico. *Fish Bull* 96, 562-574

Rocha-Olivares A, Gomez-Munoz VM (2003) Validation of otolith age determination of the Pacific red snapper *Lutjanus peru* (Perciformes: Lutjanidae) in La Paz Bay and adjacent waters, B.C.S., Mexico. *Ciencias Marinas* 19, 321-331

Rojo-Vazquez JA, .Lucano-Ramirez G, Ruiz-Ramirez S (2009) Length–weight relationships for coastal fish species from the gillnet artisanal fishery.in the central Mexican Pacific *J Appl Ichthyol* 25 497-498

Sala E, Aburto-Oropeza O, Paredes G, -Thompson G (2003) Spawning aggregations and reproductive behavior of reef fishes in the Gulf of California. *Bull Mar Sci* 72, 103-121

Santamaria-Miranda A, Elorduy-Garay JF, Villalejo-Fuerte M, Rojas-Herrera AA (2003) Desarrollo gonadal y ciclo reproductivo de *Lutjanus peru* (Pisces: Lutjanidae) en Guerrero, Mexico. *Rev Biol Trop* 51, 489-502 [in Spanish]

Zapata FA, Herron PA (2008) Pelagic larval duration and geographic distribution of tropical eastern Pacific snappers. *Mar Ecol Progr Ser* 230, 295-300

Macrouridae

Andrews AH, Cailliet GM, Coale KH (1999) Age and growth of the Pacific grenadier (*Coryphaenoides acrolepis*) with age estimate validation using an improved radiometric ageing technique. Can J Fish Aquat Sci 56, 1339-1350

Burton EJ (1999) Radiometric age determination of the giant grenadier (*Ambloplites pectoralis*) using ^{210}Pb : ^{226}Ra disequilibria. MSc thesis in Mar Sci, San Fran State Univ

Burton EJ, Andrews AH, Coale KH, Cailliet GM (1999) Application of radiometric age determination to three long-lived fishes using ^{210}Pb : ^{226}Ra disequilibria in calcified structures: a review P 77-87 in (JM Musick, ed), Life in the Slow Lane: Ecology and Conservation of Long-lived Marine Animals. Am Fish Soc

Datsky AV (2017) Biological Features of the Common Fish Species in Olyutorsky-Navarin Region and the Adjacent Waters of the Bering Sea: 2. Families Macrouridae, Clupeidae, and Osmeridae. J Ichthyol 57, 106-121

Hoff GR, Buckley TW, Drazen JC, Duncan KM (2000) Biology and ecology of *Nezumia liolepis* and *N. stelgidolepis* from the west coast of North America. J Fish Biol 57, 662-680

Kitazawa N (2000) Maturity of roundnose grenadier *Coryphaenoides acrolepis* in Sagami Bay. Bull Kanagawa Prefect Fish Res Inst 5, 1-5

Rodgveller CJ, Clausen DM, Nagler JJ, Hutchinson C (2010) Reproductive Characteristics and Mortality of female giant grenadiers in the Northern Pacific Ocean. Mar Coast Fish Dyn Manag Ecosys Sci 2. 73-82

Stein DL, Percy WG (1982) Aspects of reproduction, early life history, and biology of macrourid fishes off Oregon, USA. Deep-Sea Res 29(11A), 1373-1379

Wilson RR Jr (1988) Analysis of growth zones and microstructure in otoliths of two macrourids from the North Pacific abyss. Environ Biol Fish 21, 251-261

Malacanthidae

Bellquist LF, Lowe CG, Caselle JE (2008) Fine-scale movement patterns, site fidelity, and habitat selection of ocean whitefish (*Caulolatilus princeps*). Fish Res 91, 325-335

Cooksey DJ (1980) Age, growth and maturity of the ocean whitefish, *Caulolatilus princeps*. MSc thesis, Calif State Univ Long Beach

Elorduy-Garay JS, Ramirez-Luna S (1994) Gonadal development and spawning of female ocean whitefish, *Caulolatilus princeps* (Pisces: Branchiostegidae) in the Bay of La Paz, B.C.S., Mexico. J Fish Biol 44, 553-564

Elorduy-Garay JS, Ruiz-Cordova SS, Diaz-Urbe JG (2005) Age, growth and mortality of *Caulolatilus princeps* (Pisces: Malacanthidae) from the southern Gulf of California. *Hidrobiologica* 15, 289-297

Mason T (2008) Home range size, habitat use, and the effects of habitat breaks on the movements of temperate gamefishes in a Southern California marine protected area. MSc thesis, Calif State Univ, Long Beach

Moser HG, Sumida BY, Ambrose, DA, Sandknop EM, Stevens EG (1986) Development and distribution of larvae and pelagic juveniles of ocean whitefish, *Caulolatilus princeps*, in the CalCOFI survey region. *Calif Coop Ocean Fish Rep* 27, 162-169

Melamphaidae

Andrianov DP, Bekker VE (1989) Reproductive biology of mesopelagic fishes of the eastern Pacific. *J Ichthyol* 29, 43-48

Bartow KA (2010) Taxonomy and ecology of the deep pelagic fish family Melamphaidae, with emphasis on interactions with a midocean ridge system. PhD thesis, Florida Atlantic Univ, 200p

Belova GV, Savinykh VF (2007) Reproductive biology of the highsnout bigscale *Melamphaes lugubris* (Melamphaidae) from the North-west Pacific. *Trans Pac Res Fish Centre (TINRO)* 150, 189-196

Childress JJ, Taylor SM, Cailliet GM, Price MH (1980) Patterns of growth, energy utilization and reproduction in some meso- and bathypelagic fishes off Southern California. *Mar Biol* 61, 27-40

Shelekhov VA, Savinykh VF (2006) Age and growth of the highsnout bigscale, *Melamphaes lugubris*. North Pacific Marine Science Organization (PICES) Book of Abstracts 59

Merlucciidae

Bailey KM, Francis RC, Stevens PR (1982) The life history and fishery of Pacific whiting, *Merluccius productus*. *Calif Coop Oceanic Fish Rep* XXIII, 81-98

Dark TA (1975) Age and growth of the Pacific hake, *Merluccius productus*. *Fish Bull* 73, 336-355

Ressler PH, Holmes JA, Fleischer GW, Thomas RE, Cooke KC (1980) Pacific hake, *Merluccius productus*, autecology: a timely review. *Mar Fish Rev* 69(1-4), 1-24

- Molidae

Pope EC, Hays GC, Thys TM, Doyle TK, Sims DW, Queiroz N, Hobson VJ, Kubicek L, Houghton JDR (2010) The biology and ecology of the ocean sunfish *Mola mola*: a review of current knowledge and future research perspectives. *Rev Fish Biol Fish* 20, 471-487

- Moridae

Orlov AM, Abramov AA (2002) New data on flat-nosed hake *Antimora microlepis* (Moridae) from the Northwestern Pacific Ocean. *J Ichthyol* 42, 65-73

Moronidae

Clark GH (1938) Weight and age determination of striped bass. *Calif Fish Game* 24, 176-177

Robinson JB (1960) The age and growth of striped bass (*Roccus saxatilis*) in California. *Calif Fish Game* 46, 279-290

Rypel AL (2012) Concordant estimates of countergradient growth variation in striped bass (*Morone saxatilis*) using comparative life-history data. *Can J Fish Aquat Sci* 69, 1261-1265

Sabal MC, Michel CJ, Smith JM, Hampton A, Hayes SG (2019) Seasonal movement patterns of striped bass (*Morone saxatilis*) in their nonnative range. *Est Coast* 42, 567-579

Scofield EC (1931) The striped bass of California. *Calif Dep Fish Game, Fish Bull* 29, 82p

Stevens DE, Kohlhorst DW (2001) Striped bass. Pp 460-464 in *Calif Marine Resources: Status Rep*, Calif Dep Fish Game

Mugilidae

Ellender BR, Taylor Z, Weyl O (2012) Validation of growth zone deposition rate in otoliths and scales of flathead mullet *Mugil cephalus* and freshwater mullet *Myxus capensis* from fish of known age. *Afr Mar Sci*, 455-458

Gonzalez-Castro M, Macchi GJ, Cousseau MB (2020) Studies on reproduction of the mullet *Mugil platanus* Günther, 1880 (Actinopterygii, Mugilidae) from the Mar Chiquita coastal lagoon, Argentina: Similarities and differences with related species. *Ital J Zool* 78, 343-353

Greeley MS Jr, Calder DR, Wallace RA (1987) Oocyte growth and development in the striped mullet, *Mugil cephalus*, during seasonal ovarian recrudescence: relationship to fecundity and size at maturity. Fish Bull 85, 187-200

Ibanez-Aguirre AL, Gallardo-Cabelo M (2004) Reproduction of *Mugil cephalus* and *M. curema* (Pisces: Mugilidae) from a coastal lagoon in the Gulf of Mexico. Bull Mar Sci 75, 37-49

Jacot(1920) Age, growth and scale characters of the mullets, *Mugil cephalus* and *Mugil curema*. Trans Am Microscop Soc 39, 199-229

Ruiz-Campos G, Gonzalez- Acosta AF, De La Cruz-Aguero J (2006) Length–weight and length–length relationships for some continental fishes of northwestern Baja California, Mexico. J appl Ichthyol 22, 314-316

Thomson JM (1963) Synopsis of biological data on the grey mullet *Mugil cephalus* Linnaeus 1758. CSIRO, Fisheries and Oceanography, Fisheries Synopsis, Vol 1

Warburton K (1979) Growth and production of some important species of fish in a Mexican coastal lagoon system. J Fish Biol 14, 449-464

Mullidae

Lucano-Ramirez G, Ruiz-Ramirez S, Rojo-Vasquez JA (2006) Composición por tallas y ciclo reproductivo de *Pseudupeneus grandisquamis* (Pisces: Mullidae) en el Pacífico central Mexicano. Rev Biol Trop 54,95-207 [in Spanish]

Lucano-Ramirez G, Robles-Ravelero M, Ruiz-Ramírez S, González-Sansón G, Aguilar-Betancourt C, Perez-Toledo A (2019) Biología reproductiva de *Mulloidichthys dentatus* (Perciformes: Mullidae) en el Pacífico tropical mexicano Rev Biol Mar Sci Oceanogr 54, 118-128 [in Spanish]

Morales-Azpeitia R, López Martínez J, Rodríguez Romero J, Ponce- Palafox JT, (2011) Distribution, abundance and reproductive pattern of *Pseudupeneus grandisquamis* Gill, 1863 (Perciformes: Mullidae) and *Urobatis halleri* Cooper, 1863 (Rajiformes: Urolophidae) in the Gulf of California. Investigacion Ciencia Univ Auton Aguascalientes 52, 3-14

Pavlov ZZ, . Emel'yanovaa L, Thuan TB, Hab VT (2011) Reproduction and initial development of manybar goatfish *Parupeneus multifasciatus* (Mullidae). J Ichthyol 51, 604-617

Pavlov N. G. Emel'yanovaa NG, Hab VT, Luong-Thuan TB (2013) Age and growth of manybar goatfish *Parupeneus multifasciatus* (Mullidae) from the Nha Trang Bay of the South China Sea. J Ichthyol 53, 442-449

Muraenidae

Higgins BA, Pearson D, & Mehta RS (2017) El Niño episodes coincide with California moray *Gymnothorax mordax* settlement around Santa Catalina Island, California. J Fish Biol 90, 1570-1583

Mehta RS, Dale KE, Higgins BA (2020) Marine protection induces morphological variation in the California moray, *Gymnothorax mordax*. Integr Comp Biol 60, 522-534

Myctophidae

Andrianov DP, Bekker VE (1989) Reproductive biology of mesopelagic fishes of the eastern Pacific. J Ichthyol 29, 43-48

Belova GV, Savinykh VF (2006) Reproductive biology of the mesopelagic fishes *Tarletonbeania crenularis* and *Ceratoscopelus warmingii* (Osteichthyes: Myctophidae) from the northwestern Pacific. N Pac Mar Sci Org (PISCES) XV, Bk Abs 53

Bystydzienska ZE, Phillips AJ, Linkowski TB (2010) Larval stage duration, age and growth of blue lanternfish *Tarletonbeania crenularis* (Jordan and Gilbert, 1880) derived from otolith microstructure. Environ Biol Fish 89, 493-503

Catul V, Gauns M, Karuppasamy PK (2011) A review on mesopelagic fishes belonging to family Myctophidae. Rev Fish Biol Fish 21, 339-354

Childress JJ, Taylor SM, Cailliet GM, Price MH (1980) Patterns of growth, energy utilization and reproduction in some meso- and bathypelagic fishes off Southern California. Mar Biol 61, 27-40

Conley WJ, Gartner JV (2009) Growth among larvae of lanternfishes (Teleostei: Myctophidae) from the eastern Gulf of Mexico. Bull Mar Sci 84, 123-135

Landaeta MF, Contreras JE, Bustos CA, Munoz G (2015) Larval growth of two species of lanternfish at nearshore waters from an upwelling zone based on otolith microstructure analyses. J appl Ichthyol 31, 106-113

Moku M, Ishimaru K, Kawaguchi K (2001) Growth of larval and juvenile *Diaphus theta* (Pisces: Myctophidae) in the transitional waters of the western North Pacific. Ichthyol Res 48, 385-390

Nishimura A, Nagasawa K, Asanuma T, Aoki H, Kubota T (1999) Age, growth, and feeding habits of lanternfish, *Stenobrachius leucopsarus* (Myctophidae), collected from the near-surface layer in the Bering Sea. Fish Sci 65, 11-15

Sassa C, Oshimo S, Tanaka H, Tsukamoto Y (2014) Reproductive biology of *Benthosema pterotum* (Teleostei: Myctophidae) in the shelf region of the East China Sea. J Mar Biol Assoc UK 94, 423-433 [abstract]

Sassa C, Takahashi Z, Tsukamoto Y (2015) Distribution, hatch-date, growth, and mortality of larval *Benthosema pterotum* (Pisces: Myctophidae) in the shelf region of the East China Sea. J Mar Biol Assoc UK 95, 161-174 [abstract]

Smoker W, Percy WG (1970) Growth and reproduction of the lanternfish *Stenobrachius leucopsarus*. J Fish Res Bd Can 27, 1265-1279

Takagi K, Yatsu A, Moku M, Sassa C (2006) Age and growth of lanternfishes, *Symbolophorus californiensis* and *Ceratoscopelus warmingii* (Myctophidae), in the Kuroshio–Oyashio Transition Zone. IchthyolRes_53, 281-289

Willis JM, Percy WG (1980) Spatial and temporal variations in the population size structure of three lanternfishes (Myctophidae) off Oregon, USA. Mar Biol 57, 181-191 [abstract]

Myliobatidae

Matern SA, Cech JJ Jr, Hopkins TE (2000) Diel movements of bat rays, *Myliobatis californica*, in Tomales Bay, California: evidence for behavioral thermoregulation? Environ Biol Fish 58, 173-182

Klimley AP, Kihlslinger RI, Kelly JT (2005) Directional and non-directional movements of bat rays, *Myliobatis californica*, in Tomales Bay, California. Environ Biol Fish 74, 79-88

Martin LK, Cailliet GM (1988a) Aspects of the reproduction of the bat ray, *Myliobatis californica*, in central California. Copeia 1988(3), 754-762

Martin LK, Cailliet GM (1988b) Age and growth determination of the bat ray, *Myliobatis californica* Gill, in central California. Copeia 1988(3), 762-773

Villavicencio Garayzar CJ (1996) Tallas, proporción de sexos y reproducción de *Myliobatis californica* y *M. longirostris* (Pisces: Myliobatidae) en Baja California Sur, México. Rev Biol Trop, 43, 291-295 (in Spanish with English abstract)

Myxinidae

Barss WH (1993) Pacific hagfish, *Eptatretus stouti*, and black hagfish, *E. deani*: the Oregon fishery and port sampling observations, 1988-92. Mar Fish Rev 35, 19-30

Johnson EW (1994) Aspects of the biology of Pacific (*Eptatretus stoutii*) and black (*Eptatretus deani*) hagfishes from Monterey Bay, California. MSc thesis, Calif State Univ, Fresno

Kato S (1990) Report on the biology of Pacific hagfish, *Eptatretus stoutii*, and the development of its fishery in California. NMFS Rep to Calif Fish Game

Reid Z(1990) Research on the fishery and biology of the hagfish. California Environmental Protection Agency.

vanderMeer ZZ, Kooijman SBALM (2014) Inference on energetics of deep-sea fish that cannot be aged: the case of the hagfish. J Sea Res 94, 138-143

Nematistiidae

Chávez-Arellano R, Jakes-Cota S. Ortega-Garcia S*, Sepulveda C, Aalbers S (2019) Potential use of dorsal fin spines of the roosterfish for age estimation. California Fish and Game 105, 10-20

Ortega-Garcia S, Sepulveda C, Aalbers S; Jakes-Cota U, Rodriguez-Sanchez R. (2017) Age, growth, and length–weight relationship of roosterfish (*Nematistius pectoralis*) in the eastern Pacific Ocean. Fish Bull 115, 117

Nemichthyidae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Nettastomatidae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Osmeridae

Barton MB, Moran JR, Vollenweider JJ, Heintz, Boswell KM (2017) Latitudinal dependence of body condition, growth rate, and stable isotopes of juvenile capelin (*Mallotus villosus*) in the Bering and Chukchi Seas. Polar Biol 40, 1451-1463

Bennett WA (2005) Critical assessment of the delta smelt population in the San Francisco Estuary, California. San Fran Estuar Watershed Sci 3(2), p

Benson IM, Kastle CR, Helser TE, Short JA, Anderl DM (2019) Age interpretation in eulachon (*Thaleichthys pacificus*) as suggested by otolith microchemical signatures. Environ Biol Fish 102, 629-643

Brooks ML, Fleishman E, Brown LR, Lehman PW, Werner I, Scholz N, Mitchelmore C, Lovvorn JR, Johnson ML, Schlenk D, van Drunick S, Drever JI, Stoms DM, Parker AE, Dugdale R (2012) Life histories, Salinity zones, and sublethal contributions of contaminants to pelagic fish declines Illustrated with a case study of San Francisco Estuary, California, USA. Est Coasts 35, 603-621

Burlak OV, Zhukova KA (2020) Reproductive biology of rainbow smelt *Osmerus dentex* (Osmeridae) of the Amur River. J Ichthyol 60, 462-469

Chigbu P (2000) Population biology of longfin smelt and aspects of the ecology of other major planktivorous fishes in Lake Washington. J Freshwat Res 15, 543-5xx

Chigbu P, Sibley TH (1994) Relationship between abundance, growth, egg size and fecundity in a landlocked population of longfin smelt, *Spirinchus thaleichthys*. J Fish Biol 45, 1-15

Clarke AD, Lewis A, Telmer KH, Shrimpton JM (2007) Life history and age at maturity of an anadromous smelt, the eulachon *Thaleichthys pacificus* (Richardson). J Fish Biol 71, 1479-1493

Damon LJ, Slater SB, Baxter RD, Fujimura RW (2016) Fecundity and reproductive potential of wild female delta smelt in the upper San Francisco Estuary, California. Calif Fish Game 102, 188-210

Datsky AV (2017) Biological features of the common fish species in Olyutorsky-Navarin region and the adjacent waters of the Bering Sea: 2. Families Macrouridae, Clupeidae, and Osmeridae. Ichthyol 57, 106-121

Doyle MJ, Busby MS, Duffy-Anderson JT, Picquelle SJ, Matarese AC (2002) Early life history of capelin (*Mallotus villosus*) in the northwest Gulf of Alaska: a historical perspective based on larval collections, October 1977–March 1979. ICES J Mar Sci 59, 997-1005

Gjoesaeter H (1998) The population biology and exploitation of capelin (*Mallotus villosus*) in the Barents Sea. Sarsia 83, 453-496

Gjoesaeter, H; Loeng, H (1987) Growth of the Barents Sea capelin, *Mallotus villosus*, in relation to climate. Environ Biol Fish 20, 293-300

Grossman GD (2016) Predation on fishes in the Sacramento–San Joaquin delta: current knowledge and future directions. San Fran Est Watershed Sci 14, issue 2, article 8, 9rw9b5tj

- Gustafson RG, Ford MJ, Teel D, Drake JS (2010) Status Review of eulachon (*Thaleichthys pacificus*) in Washington, Oregon, and California. NOAA Tech Memo NMFS-NWFSC-105
- Haldorson L, Craig P (1984) Life history and ecology of a Pacific-Arctic population of rainbow smelt in coastal waters of the Beaufort Sea. Trans Am Fish Soc 113, 33-38
- Hubbs CL (1925) A revision of the osmerid fishes of the North Pacific. Proc Biol Soc Wash 38 49-56
- Kurobe T, Park MO, Javidmehr A, Teh F-C, Acuña SC, Corbin CJ, Conley AJ, Bennett WA, Teh SJ (2016) Assessing oocyte development and maturation in the threatened Delta Smelt, *Hypomesus transpacificus*. Environ Biol Fish 99, 423-432
- Latour RJ (2016) Explaining patterns of pelagic fish abundance in the Sacramento-San Joaquin delta. Ests Coasts 39, 233-247
- Merz JE, Bergman PS, Melgo JF, Hamilton S (2013) Longfin smelt: spatial dynamics and ontogeny in the San Francisco Estuary, California. Calif Fish Game 99, 122-148
- Middaugh DP, Hemmer MJ, Penttila DE (1987) Embryo ecology of the Pacific surf smelt, *Hypomesus pretiosus* (Pisces: Osmeridae). Pac Sci 41, 41-53
- Miller BS, Simenstad CA, Moulton LL, Karp WA, Fresh KL, Funk FC (1977) Puget Sound Baseline Program nearshore fish survey: data summary report FRI-UW-7711. Fish Res Inst, College of Fisheries, Univ Wash, Seattle
- Moulton LL (1974) Abundance, growth, and spawning of the longfin smelt in Lake Washington. Trans Am Fish Soc 103, 46-52
- Moyle PB, Brown LR, Durand JR, Hobbs JA (2016) Delta Smelt: Life history and decline of a once-abundant species in the San Francisco Estuary SF Est Watershed Sci 14(2), item 09k9f76s
- Moyle PB, Herbold B, Stevens DE, Miller LW (1992) Life history and status of delta smelt in the Sacramento-San Joaquin Estuary, California. Trans Am Fish Soc 121(1), 67-77
- Nobriga ML, Rosenfield JA (2016) Population dynamics of an estuarine forage fish: disaggregating forces driving long-term decline of longfin smelt in California's San Francisco Estuary. Trans Am Fish Soc 145, 44-58
- Olafsdottir AH, Anderson JT (2010) Growth and survival of Icelandic capelin *Mallotus villosus* larvae. Mar Ecol Progr Ser 403, 231-241
- Quinn T, Krueger K, Pierce K, Penttila D, Perry K, Hicks T, Lowry D (2012) Patterns of surf smelt, *Hypomesus pretiosus*, Intertidal spawning habitat use in Puget Sound, Washington State. Ests Coasts 35, 1214-1228

Rice CA (2006) Effects of Shoreline Modification on a Northern Puget Sound Beach: microclimate and embryo mortality in surf smelt (*Hypomesus pretiosus*). *Ests Coasts* 29, 63-71

Rose KA,, Kimmerer WJ, Edwards KP, Bennett WA (2013) Individual-based modeling of delta smelt population dynamics in the Upper San Francisco Estuary: I. model description and baseline results. *Trans Am Fish Soc* 142, 1238-1259

Rosenfield JA, Baxter RD (2007) Population dynamics and distribution patterns of longfin smelt in the San Francisco Estuary. *Trans Am Fish Soc* 136, 1577-1592

Saruwatari T, Lopez JA, Pietsch TW (1997) A revision of the osmerid genus *Hypomesus* Gill(Teleostei: Salmoniformes), with the description of a new species from the Southern Kuril Islands. *Spp Diversity* 2, 59-82

Skurikhina LA, Oleinik AG, Kukhlevsky AD, Kovpak NE, Frolov SV, Sendek DS (2018) Phylogeography and demographic history of the Pacific smelt *Osmerus dentex* inferred from mitochondrial DNA variation. *Polar Biol* doi.org/10.1007/s00300-018-2250-4

Sommer T, Mejia FH, Nobriga ML (2011) The spawning migration of delta smelt in the Upper San Francisco Estuary. *San Fran Est Watershed Sci* 9(2), 1-16

Willson-MF, Armstrong RH, Hermans MC, Koski K (2006) Eulachon: a review of biology and an annotated bibliography. NMFS, AFSC Processed Rep 2006-7

Paralichthyidae

Barnes CL (2015) Growth, mortality, and reproductive potential of California halibut (*Paralichthys californicus*) off Central California. MSc thesis, San Jose State Univ, 101p

Barnes CL, Starr RM (2018) Reproductive tactics of California halibut (*Paralichthys californicus*): combining spawning season, interspawning interval, and batch fecundity to estimate annual reproductive output for a multiple batch spawning fish. *Calif Coop Oceanic Fish Invest Rep* 59, 102-114

Barnes CCL, Starr RM, Reilly PN (2015) Growth, mortality, and reproductive seasonality of California halibut (*Paralichthys californicus*: a biogeographic approach. *Calif Coop Oceanic Fish Invest Rep* 56, 110-118

Clark GH (1950) The California halibut (*Paralichthys californicus*) and an analysis of the boat Catches. *Calif Dep Fish Game, Fish Bull* 32, 52 p

Craig MT, Fodrie FJ, Hastings PA(2004) The Nearshore Fish Assemblage of the Scripps Coastal Reserve, San Diego, California. *Coast Manag* 32, 341-351

Donohoe ZZ (2000) Metamorphosis, growth, and settlement of Pacific sanddab (*Citharichthys sordidus*) to a continental shelf nursery, inferred from otolith microstructure. PhD thesis, Univ Oregon, 232 p

Groce AK (2002) Influence of life history and lipids on the bioaccumulation of organochlorines in demersal fishes. MSc thesis, San Diego State Univ

Haugen DW (ed) (1990) .The California halibut, *Paralichthys californicus*, resources and fisheries. Calif. Dep. Fish Game, Fish Bull. 174, xxx p.

He X, Field JC, -Pearson DE, Lefebvre LS (2016) Age sample sizes and their effects on growth estimation and stock assessment outputs: three case studies from U.S. West Coast fisheries. Fish Res 180, 92-102

He X, Pearson DE, Field JC, Lefebvre L, Key M (2013) Status of the U.S. Pacific Sanddab resource in 2013. SAFE Rep.

Kramer SH (1990) Growth, mortality, and movements of juvenile California halibut *Paralichthys californicus* in shallow coastal and bay habitats of San Diego County, California. Fish Bull 89, 195-207

Lefebvre LS, Payne AM, Field JC (2016) Reproductive dynamics of Pacific sanddab, *Citharichthys sordidus*, off the central coast of California. J Sea Res 107, 100-111

Lesyna KM, Barnes CL (2016) Assessment of length- and age-at-maturity for California halibut (*Paralichthys californicus*), including a histologically-based description of the reproductive cycle. Calif Fish Game 102, 79-99

Love MS, Brooks A (1990) Size and age at first maturity of the California halibut, *Paralichthys californicus*, in the Southern California Bight. Calif Dep Fish Game, Fish Bull 174, 167-174

Love MS, Stephens JS J, Morris PA, Singer MM, Sandhu M (1986) Inshore soft substrata fishes in the Southern California Bight. Calif Coop Oceanic Fish Invest Rep 27, 84-106

MacNair LS, Domeier ML, Chun CSY (2001) Age, growth, and mortality of California halibut, *Paralichthys californicus*, along southern and central California. Fish Bull 99, 588-600

Martinez-Munoz MA, Ortega-Salas AA (1999) Growth and mortality of the fantail sole, *Xystreureys liolepis* (Jordan and Gilbert 1881) off the western coast of Baja California, Mexico. Bull So Calif Acad Sci 98, 66-74

Martinez-Munoz MA, Ortega-Salas AA (2001) Growth and mortality of the bigmouth sole, *Hippoglossina stomata* Eigenmann & Eigenmann 1890 (Pisces: Paralichthyidae), off the western coast of Baja California, Mexico. Bull Mar Sci 69, 1109-1119

Martinez-Munoz MA, Ortega-Salas AA (2010) Growth and mortality of the fish *Citharichthys xanthostigma* (Pleuronectiformes: Paralichthyidae) off the western coast of Baja California, México. Rev Biol Trop 58, 689-705

Maunder M, Reilly P, Tanaka T, Schmidt G, Penttila K (2011) California halibut stock assessment. Calif Dep Fish Game. [https:// www.dfg.ca.gov/marine/sfmp/halibut-assessment.asp](https://www.dfg.ca.gov/marine/sfmp/halibut-assessment.asp)

Oda D (1991) Oda(1991) Development of eggs and larvae of California halibut *Paralichthys californicus* and Fantail Sole *Xystreurys liolepis* (Pisces: Paralichthyidae). Fish Bull 89, 387-402

Protopapadakis L, Penttila K, Dowd WW (2015) Testing a non-lethal method for determining the sex of California halibut, *Paralichthys californicus*, in non-spawning condition. Fish Manag Ecol 22, 432–435

Rodriguez-Romero J, Palacios-Salgado DS, Lopez-Martinez J, Hernandez-Vazquez JS, Velazquez-Abunader JI (2009) The length–weight relationship parameters of demersal fish species off the western coast of Baja California Sur, Mexico J appl Ichthyol 25, 114-116

Sakuma KM, Larson RJ (1995) Distribution of pelagic metamorphic-stage sanddabs *Citharichthys sordidus* and *C. stigmaeus* within areas of upwelling off central California. Fish Bull 93, 516-529

Sears-Hartley, J. 1994. Age, growth, and early life history of *Paralichthys californicus* derived from otolith microstructure. MSc thesis, California State University, Northridge. 54 p

Sunada, J.S., Velez PV, Pattison. CA 1990. Age, size, and sex composition of California halibut from southern California commercial fishery landings, 1983–1988. Calif. Dep. Fish Game, Fish Bull. 174, 303-319

Pentacerotidae

Boehlert GW, Sasaki (1988) Pelagic biogeography of the armorhead, *Pseudopentaceros wheeleri*, and recruitment to isolated seamounts in the North Pacific Ocean. Fish Bull 86, 453-465

Humphreys RL Jr (2000) Otolith-based assessment of recruitment variation in a North Pacific seamount population of armorhead *Pseudopentaceros wheeleri*. Mar Ecol Progr Ser 204, 213-223

Humphreys RL Jr, Tagami DT (1986) Review and current status of research on the biology and ecology of the genus *Pseudopentaceros*. NOAA Tech Rep NMFS-43, 55-62

Humphreys RL Jr, Winans GA, Tagami DT (1989) Synonymy and life history of the North Pacific pelagic armorhead, *Pseudopentaceros wheeleri* Hardy (Pisces: Pentacerotidae). Copeia 1989(1), 142-155

Kiyota M, Nishida K, Murakami C, Yonezaki S (2016) History, biology, and conservation of Pacific endemics 2. The North Pacific armorhead, *Pentaceros wheeleri* (Hardy, 1983) (Perciformes, Pentacerotidae). PacSci 70, 1-20

Murakami C, Yonezaki S, Suyama S, Nakagami N, Okuda O, Kiyota K (2016) Early epipelagic life-history characteristics of the North Pacific armorhead *Pentaceros wheeleri*. Fish Sci 82, 709-718

Somerton DA, Kikkawa BS (1992) Population dynamics of pelagic armorhead *Pseudopentaceros wheeleri* on Southeast Hancock Seamount. Fish Bull 90, 756-769

Uchida RN, Hayasi S, Boehlert GW (eds) (1986) Environment and Resources of Seamounts in the North Pacific: *Proc Workshop MARCH 21-23, 1984 Shimizu, Japan*, NOAA Tech Rep NMFS 43,

Uchida RN, Tagami DT (1984) Groundfish fisheries and research in the vicinity of seamounts in the North Pacific Ocean. Mar Fish Rev 46(2), 1-17

Uchiyama JH, Kazama TK (2003) Updated weight-on-length relationships for pelagic fishes caught in the central North Pacific Ocean and bottomfishes from the Northwestern Hawaiian Islands. NMFS PIFSC Admin Rep H-03-01

Uchiyama JH, Kuba SH, Tagami DT (1983) Length-weight and standard length-fork length relationships of deepsea handline fishes of the Northwestern Hawaiian Islands. Proc Res Inv NWHI UNIH-SEAGRANT-MR-84-01, 209-225

Uchiyama JH, Sampaga JD (1990) Age Estimation and Composition of Pelagic Armorhead *Pseudopentaceros wheeleri*; from the Hancock Seamounts. Fish Bull 88, 217-222

Uchiyama JH, Tagami DT (1983) Life history, distribution, and abundance of bottom fishes in the Northwestern Hawaiian Islands. Proc Res Invest NWHI_UNIH-SEAGRANT-MR-84-01

Yanagimoto T, Humphreys RL Jr (2005) Maturation and reproductive cycle of female armorhead *Pseudopentaceros wheeleri* from the southern Emperor-northern Hawaiian ridge seamounts. Fish Sci 71, 1059-1068

Petromyzontidae

Chase SD (2001) Contributions to the life history of adult Pacific lamprey (*Lampetra tridentata*) in the Santa Clara River of Southern California. Bull So Calif Acad Sci, 74-85

Clemens BJ, Stan van de Wetering S, Stacia A. Sower SA, Schreck- CB 2013) Maturation characteristics and life-history strategies of the Pacific lamprey, *Entosphenus tridentatus*. Can J Zool 91, 775-788

Clems BJ, Weitkamp L, Siwicke K, Wade J, Harris J, Hess J, Porter L, Parker K, Sutton T, Alexei Men. Orlov AM (2019) Marine biology of the Pacific lamprey *Entosphenus tridentatus*. Rev Fish Biol Fish. s11160-019-09578-8

Pholididae

Burgess TD (1978) The comparative ecology of two sympatric polychromatic populations of *Xerperes fucorum* Jordan & Gilbert (Pisces: Pholididae) from the rocky iintertidal zone of central California. J Exp Mar Biol Ecol 35, 43-58

Deforest LG, Busby MS (1986) Development of larval and early juvenile penpoint gunnel (*Apodichthys flavidus*) (family: Pholidae). Fish Bull 104, 476-481

Hughes GW (1985) The comparative ecology and evidence for resource partitioning in two pholidid fishes (Pisces: Pholididae) from southern British Columbia eelgrass beds Can J Zool 63, 76-85

Hughes GW (1986) Observations on the reproductive ecology of the crescent gunnel, *Pholis laeta*, from marine inshore waters of southern British Columbia. Can Fld-Nat 100, 367-370

Wilkie DW (1967) Colour pigments in the penpoint gunnel *Apodichthys flavidus* and their ecological significance. MSc thesis, Zool Dep, Univ British Columbia,, 158 p

Platyrrhynidae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Pleuronectidae

Abookire AA (2006) Reproductive biology, spawning season, and growth of female rex sole (*Glyptocephalus zachirus*) in the Gulf of Alaska. Fish Bull 104,350-359

Abookire AA, Macewicz BJ (2003) Latitudinal variation in reproductive biology and growth of female Dover sole (*Microstomus pacificus*) in the North Pacific, with emphasis on the Gulf of Alaska stock. J Sea Res 50, 187-197

Albert OT (2016) Growth and formation of annual zones in whole otoliths of Greenland halibut, a slow-growing deep-water fish. *Mar Freshwat Res* 67, 937-942

Alton MS, Bakkala RG, Walters GE, Munro PT (1988) Greenland turbot *Reinhardtius hippoglossoides* of the Eastern Bering Sea and Aleutian Islands region. NOAA Tech Rep NMFS 71, 31p

Alverson DL, Chatwin BM (1957) Results from tagging experiments on a spawning stock of petrale sole, *Eopsetta jordani* (Lockington) *J Fish Res Bd Can* 14, 953-974

Boehlert GZ, Mundy BC (1987) Recruitment dynamics of metamorphosing English sole, *Parophrys vetulus*, to Yaquina Bay, Oregon. *Est Coast Shlf Sci* 25, 261-281

Brodziak J, Mikus R (2000) Variation in life history parameters of Dover sole, *Microstomus pacificus*, off the coasts of Washington, Oregon, and northern California. *Fish Bull* 98, 661-673

Butler JL, Dahlin KA, Moser HG (1996) Growth and duration of the planktonic phase and a stage-based population matrix of Dover sole, *Microstomus pacificus*. *Bull Mar Sci* 58, 29-43

Clark WG, Hare JR, Parma AM, Sullivan PJ, Trumble RJ (1999) Decadal changes in growth and recruitment of Pacific halibut (*Hippoglossus stenolepis*). *Can J Fish Aquat Sci* 56, 242-252

Cooper LD (1996) Age and growth in the hornyhead turbot (*Pleuronichthys verticalis*) off Orange County, California. Southern California Coastal Water Res Project (SCCWRP), Annu Rep 1994-1995 [abstract]

Cooper DW, Maslenikov KP, Gunderson DR (2007) Natural mortality rate, annual fecundity, and maturity at length for Greenland halibut (*Reinhardtius hippoglossoides*) from the northeastern Pacific Ocean. *Fish Bull* 105, 296-304

Doyle MJ, Debenham C, Barbeaux SJ, Buckley TW, Pirtle JL, Spies IB, Stockhausen WT, Shotwell SK, Wilson MT, Cooper DW (2018) A full life history synthesis of arrowtooth flounder ecology in the Gulf of Alaska: exposure and sensitivity to potential ecosystem change. *J Sea Res* 142, 28-51

Doyle MJ, Mier KL (2016) Early life history pelagic exposure profiles of selected commercially important fish species in the Gulf of Alaska. *Deep-Sea Res II*, 132, 162-193

Fargo J, Kronlund AR (2000) Variation in growth for Hecate Strait English sole (*Parophrys vetulus*) with implications for stock assessment. *J Sea Res* 44, 3-15

Fargo J, Wilderbuer T (2000) Population dynamics of rock sole (*Lepidopsetta bilineata*) in the North Pacific. *J Sea Res* 44, 123-144

Gadomski DM, Petersen JH (1988) Effects of food deprivation on the larvae of two flatfishes. *Mar Ecol Prog Ser* 44, 103-111

Gao Y (2012) Otoliths speak out: why the Pacific halibut in Puget sound are different. *Environ Biol Fish* 95, 469-479

Garrett DL, Pietsch TW, Utter FM, Hauser L (2007) The hybrid sole *Inopsetta ischyra* (Teleostei: Pleuronectiformes: Pleuronectidae): hybrid or biological species? *Trans Am Fish Soc* 136:460–468

Glubokov AI, Afanasyev PK, Melnikov SP (2017) New data on rex sole *Glyptocephalus zachirus* (Pleuronectidae) in the northwestern part of the Bering Sea. *J Ichthyol* 57, 926-930

Gundersen AC, Stenberg C, Fossen I, Lyberth B, Boje J, Jorgensen OA (2010) Sexual maturity cycle and spawning of Greenland halibut *Reinhardtius hippoglossoides* in the Davis Strait. *J Fish Biol* 77, 211–226

Gunderson DR, Zimmerman M, Nichol DG, Pearson K (2003) Indirect estimates of natural mortality rate for arrowtooth flounder (*Atheresthes stomias*) and darkblotched rockfish (*Sebastes crameri*). *Fish Bull* 101, 175-182

Hagerman FB (1952) The biology of the dover sole, *Microstomus pacificus* (Lockington). *Calif Dep Fish Game, Fish Bull* 85,48p

Haltuch MA, Hamel OS, Piner KR, McDonald P, Kastle CR, Field JC (2013) California Current bomb radiocarbon reference chronology and petrale sole (*Eopsetta jordani*) age validation. *Can J Fish Aquat Sci* 70, 22-31

Haltuch MA, Hicks A (2009) Status of the U.S. petrale sole resource in 2008. *SAFE Rep Final* 11-1-7-09

Hannah R, Parker SJ, Fruh EL (2002) Length and age at maturity of female petrale sole (*Eopsetta jordani*) determined from samples collected prior to spawning aggregation. *Fish Bull* 100, -711-719

Holsman KK, Aydin K, Sullivan J, Hurst T, Kruse GH (2019) Climate effects and bottom-up controls on growth and size-at-age of Pacific halibut (*Hippoglossus stenolepis*) in Alaska (USA). *Fish Oceanogr* 28, 345-358

Hosie MJ, Horton HF (1977) Biology of the rex sole, *Glyptocephalus zachirus*, in waters off Oregon. *Fish Bull* 75, 51-60

Hunter JR, Butler JL, Kimbrell C, Lynn EA (1990) Bathymetric patterns in size, age, sexual maturity, water content, and caloric density of Dover sole, *Microstomus pacificus*. *Calif Coop Ocean Fish Rep* 31, 132-144

Hunter JR, Macewicz BJ, Lo NCH, Kimbrell CA (1992) Fecundity, spawning, and maturity of female Dover sole *Microstomus pacificus*, with an evaluation of assumptions and precision. *Fish Bull* 90, 101-128

Hurst TP, Abookire AA (2006) Temporal and spatial variation in potential and realized growth rates of age-0 year northern rock sole. *J Fish Biol* 68, 905-919

Hurst TP, Abookire AA, Knott B (2010) Quantifying thermal effects on contemporary growth variability to predict responses to climate change in northern rock sole (*Lepidopsetta polyxystra*). *Can J Fish Aquat Sci* 67, 97-107

Inagawa R, Hattori T, Watanabe K, Narimatsu Y, Ito M (2012) Growth pattern and age composition of roughscale sole *Clidoderma asperrimum* off the Pacific coast of northern Honshu, Japan. *Nipp Suis Gakkaishi* 78, 1118-1126 [in Jpn with Engl abstract]

Jacobson LD, Hunter JR (1993) Bathymetric demography and management of Dover sole. *No Am J Fish Manag* 13, 405-420

Kastelle CR, Helser TZ, Wischniowski SG, Loher T, Goetz BJ, Kautzi LA (2016) Incorporation of bomb-produced ^{14}C into fish otoliths: A novel approach for evaluating age validation and bias with an application to yellowfin sole and northern rockfish. *Ecol Model* 320, p79-91

Kobelev EA (1989) Some biological characteristics of the Arctic flounder, *Liopsetta glacialis* in the southeastern area of the Barents Sea. *J Ichthyol* 29, 16-21

Kramer SH (1991) The shallow-water flatfishes of San Diego county. *Calif Coop Oceanic Fish Invest Rep* 32, p128-142

Lane ED (1975) Quantitative aspects of the life history of the diamond turbot, *Hypsopsetta guttulata* (Girard) in Anaheim Bay. *Calif Dep Fish Game, Fish Bull* 165, 153-174

Laurel BJ, Danley C, Haines S (2014) The effects of temperature on growth, development and settlement of northern rock sole larvae (*Lepidopsetta polyxystra*). *Fish Oceanogr* 23, 495-505

Loher T (2008) Homing and summer feeding site fidelity of Pacific halibut (*Hippoglossus stenolepis*) in the Gulf of Alaska, established using satellite-transmitting archival tags. *Fish Res* 92, 63-68

Markle DF, Harris PM, Toole CL (1992) Metamorphosis and an overview of early-life-history stages in Dover sole *Microstomus pacificus*. *Fish Bull* 90, 285-301

Matta ME, Helser TE, Black BA (2016) Otolith biochronologies reveal latitudinal differences in growth of Bering Sea yellowfin sole *Limanda aspera*. *Polar Biol* 39, 2427-2439

Mukhametov IN (2001) On reproduction of the Kamchatka flounder *Atheresthes evermanni* in Pacific waters off the Northern Kuril Islands. *J Ichthyol* 41, 382-396

Nichol DG (1997) Effects of geography and bathymetry on growth and maturity of yellowfin sole, *Pleuronectes asper* in the eastern Bering Sea. *Fish Bull* 95, 494-503

Nichol DG, Acuna EI (2001) Annual and batch fecundities of yellowfin sole, *Limanda aspera*, in the eastern Bering Sea. Fish Bull 99, 108-122

Nunez LA, Hallfredsson EH, Falk-Petersen I-B (2015) Different maturity scales affect estimations of fecundity, TEP and spawning stock size of Greenland halibut, *Reinhardtius hippoglossoides* (Walbaum, 1792). Mar Biol Res 11, 824-833

Orcutt HG (1950) The life history of the starry flounder *Platichthys stellatus* (Pallas). Calif Dep Fish Game, Fish Bull 78, 64p

Orlov AM, Binohlan C (2009) Length–weight relationships of deep-sea fishes from the western Bering Sea J appl Ichthyol 25, 223-227

Orlov AM, Kuznetsova EN, Mukhametov IN (2011) Age and growth of the Pacific halibut *Hippoglossus stenolepis* and the size-age composition of Its catches in the northwestern part of the Pacific Ocean. J Ichthyol 51, 306-323

Pearson DE, McNally SVG (2005) Age, growth, life history, and fisheries of the sand sole, *Psettichthys melanostictus*. Mar Fish Rev 67, 9-18

Peterman RM, Bradford MJ (1987) Density-dependent growth of age 1 English sole (*Parophrys vetulus*) in Oregon and Washington coastal waters. Can J Fish Aquat Sci 44, 48-53

Ralston S (2005) An assessment of starry flounder off California, Oregon, and Washington. Pac Fish Manag Council, Portland OR

Richardson SL (1981) Pelagic eggs and larvae of the deepsea sole, *Embassichthys bathybius* (Pisces: Pleuronectidae), with comments on generic affinities. Fish Bull 79, 163-170

Richardson SL, Dunn JR, Naplin NA (1980) Eggs and larvae of butter sole, *Isopsetta isolepis* (Pleuronectidae), off Oregon and Washington. FishBull 78, 401-417

Roff DA (1981) Reproductive uncertainty and the evolution of iteroparity: why don't flatfish put all their eggs in one basket? Can J Fish Aquat Sci 38, 968-977

Roff DA (1982) Reproductive strategies in flatfish: a first synthesis. Can J Fish Aquat Sci 39, 1696-1698

Roff DA (1991) The evolution of life-history variation in fishes, with particular reference to flatfishes. Neth J Sea Res 27, 197-207

Sampson DB, Al-Jufaily SM (1999) Geographic variation in the maturity and growth schedules of English sole along the U.S. west coast. J Fish Biol 54, 1-17

Schinske JN, Bernardi G, Jacobs DK, Routman EJ (2010) Phylogeography of the diamond turbot (*Hypsopsetta guttulata*) across the Baja California Peninsula. MarBiol 157, 123-134

Seitz AC, Loher T, Norcross BL, Nielsen JL (2011) Dispersal and behavior of Pacific halibut *Hippoglossus stenolepis* in the Bering Sea and Aleutian Islands region. *Aquat Biol* 12, 225-239

Smith RL, Vallarino M, Barbour E, Fitzpatrick Z, Barber WE (eds) (1997) Population biology of the Bering flounder in the northeastern Chukchi Sea. *Am Fish Soc Sci Symp* 19, 127-13

Southern California Coastal Water Research Project (1994) Aspects of the life history of hornyhead turbot, *Pleuronichthys verticalis*, off Southern California. *SCCWRP Annu Rep* 1992-1993

Stark JW (2004) A comparison of the maturation and growth of female flathead sole in the central Gulf of Alaska and southeastern Bering Sea. *J Fish Biol* 64, 876-889

Stark JW (2008a) Age- and length-at-maturity of female arrowtooth flounder (*Atheresthes stomias*) in the Gulf of Alaska. *Fish Bull* 106, 328-333

Stark JW (2011) Contrasting the maturation, growth, spatial distribution and vulnerability to environmental warming of *Hippoglossoides robustus* (Bering flounder) with *H. elassodon* (flathead sole) in the eastern Bering Sea. *Mar Biol Res* 71, issue8, 778-785

Stark JW (2012) Female maturity, reproductive potential, relative distribution, and growth compared between arrowtooth flounder (*Atheresthes stomias*) and Kamchatka flounder (*A. evermanni*) indicating concerns for management. *J appl Ichthyol* 28, 226-230

Stark JW (2012) Contrasting maturation and growth of northern rock sole in the Eastern Bering Sea and Gulf of Alaska for the purpose of stock management. *N Am J Fish Manag* 32, 93-99

Stark JW, Somerton DA (2002) Maturation, spawning and growth of rock soles off Kodiak Island in the Gulf of Alaska. *J Fish Biol* 61, 417-431

Tokranov AM, Zavarina SV (1992) Size-age structure and sex ratios of Alaska plaice, *Pleuronectes quadrituberculatus*, on the western Kamchatka shelf. *J Ichthyol* 32, 27-35

Walters GE, Wilderbuer TK (2000) Decreasing length at age in a rapidly expanding population of northern rock sole in the eastern Bering Sea and its effect on management advice *J Sea Res* 44, 17-26

Webster RA, Clark WG, Leaman BM, Forsberg JE (2013) Pacific halibut on the move: a renewed understanding of adult migration from a coastwide tagging study. *Can J Fish Aquat Sci* 70, 642-653

Wilderbuer TK, Nicol DG, Ianelli J (2016) Assessment of the yellowfin sole stock in the Bering Sea and Aleutian Islands. *No Pac Fish Manag Council, SAFE Rep*

Wilderbuer TK, Zhang CI (1999) Evaluation of the population dynamics and yield characteristics of Alaska plaice, *Pleuronectes quadrituberculatus*, in the eastern Bering Sea. Fish Res 41, 183-200

Youcef WA, Lambert Y, Audet C (2015) Variations in length and growth of Greenland halibut juveniles in relation to environmental conditions. Fish Res 167, 30-47

Zhang CI, Gunderson DR (1990) Yield per recruit and management recommendations for Alaska plaice in the Eastern Bering Sea. No Am J Fish Manag 10,p1-10

Zhang CI, Wilderbuer TK, Walters GE (1998) Biological characteristics and fishery assessment of Alaska plaice, *Pleuronectes quadrituberculatus*, in the Eastern Bering Sea. Mar Fish Rev 60, 17-28

Zimmermann M (1997) Maturity and fecundity of arrowtooth flounder, *Atheresthes stomias*, from the Gulf of Alaska. Fish Bull 95, 598-611

Zimmermann M, Goddard P (1996) Biology and distribution of arrowtooth, *Atheresthes stomias*, and Kamchatka, *A. evermanni*, flounders in Alaskan waters. Fish Bull 94, 358-370

Polyprionidae

Allen LG, Andrews AH (2012) Bomb radiocarbon dating and estimated longevity of giant sea bass (*Stereolepis gigas*). Bull So Calif Acad Sci 111, 1-14

Allen LG (2017) Giants! Or. . .the return of the kelp forest king. Copeia 105(1), 10-13

Benseman SA, Allen LG (2018) Distribution and recruitment of young-of-the-year giant sea bass, *Stereolepis gigas*, off Southern California. Copeia 106, 312-320

Benseman SA, Couffer MC, Allen LG (2019) Behavior of young-of-the-year giant sea bass, *Stereolepis gigas*, off the sandy beaches of Southern California. Bull So Calif Acad Sci 118, 79-86

Chabot CL, Hawk HA, Allen LG (2015) Low contemporary effective population size detected in the Critically Endangered giant sea bass, *Stereolepis gigas*, due to fisheries overexploitation. Fish Res 172, 71-78

Clevenstine AJ, Lowe CG (2021) Aggregation site fidelity and movement patterns of the protected marine predator giant sea bass (*Stereolepis gigas*). Environ Biol Fish
[doi.org_10.1007_s10641-021-01077-9](https://doi.org/10.1007_s10641-021-01077-9)

Hawk HA, Allen LG (2014) Age and growth of the giant sea bass, *Stereolepis gigas*. Calif Coop Ocean Fish Invest Rep 55, 128-134

Love MS, Seeto K, Jainesse C, Nishimoto MM (2018) Spots on sides of giant sea bass (*Stereolepis gigas* Ayres, 1859) are likely unique to each individual. Bull So Cal Acad Sci 117, 77-81

Pomacanthidae

Arellano-Martinez MA, Ceballos-Vazquez BP, Galvan-Magana F (2007) Reproductive cycle of the Cortez angelfish *Pomacanthus zonipectus* (Gill, 1863) (Pomacanthidae) from the Gulf of California, Mexico. Hidrobiologica 17, 241-248

Fernandez-Rivera Melo FJ, Reyes-Bonilla LAH, Lopez-Rocha JA, Salomon-Aguilar CA (2016) Growth and mortality of the king angelfish, *Holacanthus passer* (Actinopterygii: Perciformes: Pomacanthidae), in the central Gulf of California, Mexico. Acta Ichthyol Pisc 46, 293-301

Pomacentridae

Clarke TA (1968). Territorial behavior and population dynamics of the garibaldi, *Hypsypops rubicunda* (Girard), Pisces: Pomacentridae. PhD thesis, UC San Diego. 165pp.

Clarke TA (1970) Territorial behavior and population dynamics of a pomacentrid fish, the garibaldi, *Hypsypops rubicunda*. Ecol Monogr 40, 189-212

DeMartini EE, Barnett AM, Johnson TD, Ambrose RF (1994) Growth and production estimates for biomass-dominant fishes on a Southern California artificial reef. Bull Mar Sci 54, 484-500

Johnson TD, Barnett AM, DeMartini EE, Craft LL, Ambrose RF, Purcell LJ (1994) Fish production and habitat utilization on a Southern California artificial reef. Bull Mar Sci 54, 709-723

Limbaugh C (1964) Notes on the life history of two Californian pomacentrids: garibaldis, *Hypsypops rubicunda* (Girard), and blacksmiths, *Chromis punctipinnis* (Cooper). Pac Sci 18, 41-50

Walters K (2002) A geographic comparison of life histories in two species of fish. MSc thesis, San Francisco State Univ

Wellington GM, Victor BC (1989) Planktonic larval duration of one hundred species of Pacific and Atlantic damselfishes (Pomacentridae). Mar Biol 101, 557-567

Priacanthidae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Psychrolutidae

Drazen JC, Goffredi SK, Schlining B, Stakes DS (2003 Aggregations of egg-brooding deep-sea fish and cephalopods on the Gorda Escarpment: a reproductive hot spot). Biol. Bull. 205: 1–7

Geldof DL, Gidmark NJ (2016) Heterochrony in two sculpins of the genus *Psychrolutes*: growth outpaces skeletal development in an apex predator among Ichthyoplankton. Copeia 104, 683-690
Glubokov AI, Glubokovskii MK, Kobacheva NP (2019) New data on soft sculpin *Malacocottus zonurus* (Psychrolutidae) from the northwestern Bering Sea. J Ichthyol 59, 360-363

Stevenson DE (2015) The validity of nominal new data on species of *Malacocottus* (Teleostei: Cottiformes: Psychrolutidae) known from the Eastern North Pacific with a key to the species. Copeia 103, 22-33

Tokranov AM, Orlov AM (2001) Some biological features of Psychrolutidae in the Pacific waters off Southeastern Kamchatka and the Northern Kuril Islands: Communication. 2. Size-age and sex composition and feeding. J Ichthyology 41, 575-583

Rajidae

Ainsley SM, Ebert DA, Cailliet GM (2011a) Age, growth, and maturity of the whitebrow skate, *Bathyraja minispinosa*, from the eastern Bering Sea ICES J Mar Sci 68, 1426-1434

Ainsley SM, Ebert DA, Cailliet GM (2011b) A comparison of reproductive parameters of the Bering skate, *Bathyraja interrupta*, from two Alaskan large marine ecosystems. Mar Freshwat Res 62, 557-566

Ainsley SM, Ebert DA, Natanson LJ, Cailliet GM (2014) A comparison of age and growth of the Bering skate, *Bathyraja interrupta* (Gill and Townsend, 1897), from two Alaskan large marine ecosystems. Fish Res 154, 17-25

Barnett, LAK, Winton MV, Ainsley SM, Cailliet GM, Ebert DA (2013) Comparative demography of skates: life-history correlates of productivity and implications management. PLoS ONE 8, e65000

Bizarro JT, Vaughn MT (2009) Aspects of the biology and species composition of skates (Rajiformes) from insular waters of southeastern Alaska. Northwest Nat 90, 247-256

Davis C (2006) Age, growth, and reproduction of the rougtail skate *Bathyraja trachura* (Gilbert, 1892) from the eastern North Pacific. MSc thesis, Calif State Univ, Monterey Bay

Davis C, Cailliet GM, Ebert DA (2007a) Age and growth of the rougtail skate *Bathyraja trachura* (Gilbert 1892) from the eastern North Pacific. *Environ Biol Fish* 80, 325-336

Davis C, Cailliet GM, Ebert DA (2007b) Age and growth of the rougtail skate *Bathyraja trachura* (Gilbert 1892) from the eastern North Pacific. *Biology of Skates*. Springer.

Dolganov VN, Ginanova TT (2014) On little known deep-sea skate *Bathyraja abyssicola* (Gilbert, 1896) from the waters of Russia. *Trans Pac Res Fish Cen (TINRO)* 177, 152-155

Downs DE, Cheng YW (2013) Length-length and width-length conversion of longnose skate and big skate off the Pacific coast: implications for the choice of alternative measurement units in fisheries stock assessment. *North Am J Fish Manag* 33, 887-893

Ebert DA (2005) Reproductive biology of skates, *Bathyraja* (Ishiyama), along the eastern Bering Sea continental slope. *J Fish Biol* 66, 618-649

Ebert DA, Smith WD, Cailliet GM (2008) Reproductive biology of two commercially exploited skates, *Raja binoculata* and *R. rhina*, in the western Gulf of Alaska. *Fish Res* 94, 48-57

Farrugia TJ, Goldman KJ, Tribuzio C, Seitz AC (2016) First use of satellite tags to examine movement and habitat use of big skates *Beringraja binoculata* in the Gulf of Alaska. *Mar Ecol Progr Ser* 556, 209-221

Gburski CM, Gaichas SK, Kimura DK (2007) Age and growth of big skate (*Raja binoculata*) and longnose skate (*R. rhina*) in the Gulf of Alaska. *Environ Biol Fish* 80, 337-349

Gertseva V, Matson SE, Taylor I, Bizarro J, Wallace J (2019) Stock assessment of the longnose Skate (*Beringraja rhina*) in state and federal waters off California, Oregon and Washington. *Pac Fish Manag Council*, Portland OR

Haas DL (2011) Age, growth, and reproduction of the Aleutian skate, *Bathyraja aleutica*, from Alaskan waters. MSc thesis, Calif State Univ, Monterey Bay

Haas DL, Ebert DA, Cailliet GM (2016) Comparative age and growth of the Aleutian skate, *Bathyraja aleutica*, from the eastern Bering Sea and Gulf of Alaska. *Environ Biol Fish* 99, 813-828

James KC (2011) Life history characteristics of the starry skate, *Raja stellulata*, from California Waters. MSc thesis, Calif State Univ, San Jose

James KC, Ebert DA, Natanson LJ, Cailliet GM (2014) Age and growth characteristics of the starry skate, *Raja stellulata*, with a description of life history and habitat trends of the central California, U.S.A., skate assemblage *Environ Biol Fish* 97, 435-448

King JR, Helser T, Gburski C, Ebert DA, Cailliet G, Kastle CR (2017) Bomb radiocarbon analyses validate and inform age determination of longnose skate (*Raja rhina*) and big skate (*Beringraja binoculata*) in the north Pacific Ocean. Fish Res 193, 195-206

King JR, McFarlane GA (2010) Movement patterns and growth estimates of big skate (*Raja binoculata*) based on tag-recapture data. Fish Res 101, 50-59

King JR, Surry AM, Garcia S, Starr PJ (2015) Big skate (*Raja binoculata*) and longnose skate (*Raja rhina*) stock assessments for British Columbia. Can Sci Advis Sec, Res Doc 2015070, 329 p

Matta ME (2015) Reproductive biology of the Alaska skate *Bathyraja parmifera*, with comments on an intersexual individual. J Fish Biol 87, 664-678

Matta ME, Gunderson DR (2007) Age, growth, maturity, and mortality of the Alaska skate, *Bathyraja parmifera*, in the eastern Bering Sea. Environ Biol Fish 80, 309-323

Maurer JRF (2009) Life history of two Bering Sea slope skates: *Bathyraja lindbergi* and *B. maculata*. MSc thesis, Calif State Univ, Monterey Bay

McFarlane GA, King JR (2006) Age and growth of big skate (*Raja binoculata*) and longnose skate (*Raja rhina*) in British Columbia waters. Fish Res 78, 169-178

G.A. McFarlane GA, McPhie RP, King JR (2010) Distribution and life history parameters of elasmobranch species in British Columbia waters. CanTech Rep Fish Aquat Sci 2908

McPhie RP, Campagna SE (2009) Bomb dating and age determination of skates (family Rajidae) off the eastern coast of Canada. ICES J Mar Sci 66, 546-560

Orlov AM, Smirnov AA (2011) New Data on Sexual Dimorphism and Reproductive Biology of Alaska Skate *Bathyraja parmifera* from the Northwestern Pacific Ocean. J Ichthyol 51, 590-603

Ormseth OA (2019) Assessment of the skate stock complex in the Gulf of Alaska. No Pac Fish Manag Council, SAFE Rep

Perez CR, Cailliet GM, Ebert DA (2011) J Age and growth of the sandpaper skate, *Bathyraja kincaidii*, using vertebral centra, with an investigation of caudal thorns. J Mar Biol Assoc UK 91, 1145-1156

Perez-Brazen CR, Cailliet GM, Ebert DA (2014) Reproduction of the sandpaper skate, *Bathyraja kincaidii* (Garman 1908) in the eastern North Pacific. Environ Biol Fish 97, 1317-1322

Stevenson DE, ZZ-etal (2008) Emerging patterns of species richness, diversity, population density, and distribution in the skates (Rajidae) of Alaska. Fish Bull 106, 24-39

Taylor IG, -Gertseva V, Stephens A, -Bizzarro J (2019) Status of big skate (*Beringraja binoculata*) off the U.S. Pacific coast in 2019. Pac Fish Manag Council, SAFE Rep

Winton MV, Natanson LJ, Kneebone J, Cailliet GM, Ebert DA (2014) Life history of *Bathyraja trachura* from the eastern Bering Sea, with evidence of latitudinal variation in a deep-sea skate species. J Mar Biol Assoc UK 94, 411-422

Zeiner SJ (1991) Growth characteristics and estimates of age at maturity of two species of skates (*Raja binoculata* and *Raja rhina*) from Monterey Bay, California. MSc thesis, Calif State Univ, Stanislaus

Zeiner SJ, Wolf P (1993) Growth characteristics and estimates of age at maturity of two species of skates (*Raja binoculata* and *Raja rhina*) from Monterey Bay, California). Pp 87-99 in (S Branstetter, ed), Conservation Biology of Elasmobranchs. NOAA Tech Rep 115

Rhincodontidae

Hsu HH, Joung S-J, Hueter RE, Liu KM (2014) Age and growth of the whale shark (*Rhincodon typus*) in the north-western Pacific. Mar Freshwat Res 65, 1145–1154

Norman BM, Stevens JD (2007) Size and maturity status of the whale shark (*Rhincodon typus*) at Ningaloo Reef in Western Australia. Fish Res 84, 81–86

Pauly D (2002) Growth and mortality of the basking shark *Cetorhinus maximus* and their implications for management of whale sharks *Rhincodon typus*. *Occasional Paper of the IUCN Species Survival Commission, Shark Specialist Group*, 199-208.

Pauly, D; International Union for Conservation of Nature and Natural Resources, Gland (Switzerland)

Rhinobatidae

Blanco-Parra MD-P, Marquez-Farias F, Galvan-Magana F (2009) Fishery and morphometric relationships of the banded guitarfish, *Zapteryx exasperata* (Elasmobranchii, Rhinobatidae), from the Gulf of California, Mexico. Pan-Am J Aquat Sci 4, 456-465

Blanco-Parra MD-P, Marquez-Farias JF, Galvan-Magana F (2009) Reproductive biology of the banded guitarfish, *Zapteryx exasperata*, from the Gulf of California, México. J Mar Biol Assoc UK 89, 1655-1662

Downton-Hoffmann CA. (2001) Edad y crecimiento del pez guitarra *Rhinobatos productus* (Ayres 1856) en la costa Occidental de Baja California Sur, México. Masters thesis, CICIMAR-IPN. La Paz, B.C.S., México. 73 pp

Downton-Hoffmann CA (2007) Biology of guitarfish *Rhinobatos productus* (Ayres, 1856) in Baja California Sur, Mexico. PhD thesis, Mar Sci, Center of Interdiscip Mar Sci, 194 p

Marquez-Farias JF (2007) Reproductive biology of shovelnose guitarfish *Rhinobatos productus* from the eastern Gulf of California Mexico. Mar Biol 151, 1445-1454

Romo-Curiel-OE, Sosa-Nishizaki O, . Pérez-Jiménez JC, Rodríguez-Medrano MC (2017) Reproductive cycle and maternal–embryonic nutritional relationship of shovelnose guitarfish *Pseudobatos productus* in the Gulf of California. J Fish Biol 90, 889-905

Timmons M, Bray RN (1997) Age, growth, and sexual maturity of shovelnose guitarfish, *Rhinobatos productus* (Ayres). Fish Bull 95, 349-359

Villavicencio-Garayzar CJ (1993) Reproductive biology of *Rhinobatos productus* (Pisces: Rhinobatidae) in Almejas Bay, Baja California Sur, Mexico. Rev Biol Trop 41 (3B), 777-782

Villavicencio-Garayzar CJ (1995) Reproductive biology of the banded guitarfish, *Zapterix exasperata* (Pisces: Rhinobatidae), in Bahia Almejas, Baja California Sur, Mexico. Ciencias Marinas 21, 141-153

Salmonidae

Armstrong RH, Morrow JE (1980). The dolly varden (*Salvelinus malma*). Pp 99-140 in Charrs: Salmonid Fishes of the Genus *Salvelinus*. Chapter 8.8, Southeast Alaska Conservation Assessment

Balon EK (ed) (1980) Charrs: Salmonid Fishes of the Genus *Salvelinus*. Junk, The Hague, Netherlands

Beamish, R.J. [ed.]. 1995. Climate change and northern fish populations. Can. Spec. Publ. Fish. Aquat. Sci. 121: 739 p.

Berg, L.S., 1962. Freshwater fishes of the U.S.S.R. and adjacent countries. volume 1, 4th edition. Israel Program for Scientific Translations Ltd, Jerusalem

Bigler BS, Welch DW, Helle JH (1996) A review of size trends among North Pacific salmon. Can J Fish Aquat Sci 53, 455–465

Beverton RGH, Holt SJ (1959) A review of the lifespans and mortality rates of fish in nature a the relation to growth and other physiological characteristics. In p. 142-177 in (GEW

Wolstenholme and M O'Connor , eds.), Ciba Foundation, Colloquia in ageing, V. The lifespan of animals. Churchill, London.

Carlander, K.D (1969) Handbook of freshwater fishery biology, volume 1. The Iowa State University Press, Ames. Iowa. 752 p.

Chugunova, N.I (1959) Age and growth studies in fish. A systematic guide for ichthyologists. Israel Program for Scientific Translations. Izdatel'stvo Akademii Nauk SSSR, Moskva. 132 p.

Coad BW, Reist JD (2004) Annotated list of the arctic marine fishes of Canada. Can MS Rep. Fish Aquat. Sci. 2674,112 p

Cox SP, Hinch SG (1997) Changes in size at maturity of Fraser River sockeye salmon (*Oncorhynchus nerka*) and associations with temperature. Can J Fish Aquat Sci 54, 1159-1165

Davidson MA, Vaughan E (1941) Relation of population size to marine growth and time of spawning migration in the pink salmon of southeastern Alaska. J Mar Res 4, 231-246

Drummond Sedgwick, S (1982) The salmon handbook: the life and cultivation of fishes of the salmon family. Andre Deutsch Ltd., London, UK, 247 p.

Eschmeyer, WN, Herald ES, Hammann H (1983) A field guide to Pacific coast fishes of North America. Boston, Houghton Mifflin, 336 p

Fechhelm RG, Craig PC, Baker JS, Gallaway BJ (1985) Fish distribution and use of nearshore waters in the northeastern Chukchi Sea. US Department of Commerce, NOAA, and US Department of the Interior, Minerals Management Service, Outer Continental Shelf Environ Assess Program Final Rep 32, 121-298.

Hankin DG, Nicholas JW, Downey TW (1993) Evidence for inheritance of age of maturity in chinook salmon (*Oncorhynchus tshawytscha*). Can J Fish Aquat Sci 50, 347-358

Helle JH, Hoffman MS (1998). Changes in size and age at maturity of two North American stocks of chum salmon (*Oncorhynchus keta*) before and after a major regime shift in the North Pacific Ocean. N Pac Anadr Fish Comm Bull 1, 81-89

Janowicz ME, Załachowski W, Rybczyk A, Dalton S, Fernandes E, Fontoura NF (2018) Age, growth and reproductive biology of threatened westslope cutthroat trout *Oncorhynchus clarkii lewisi* inhabiting small mountain streams. J. Fish. Biol. 2018:1-13

Jonsson B, Hindar K, Northcote TG (1984) Optimal age at sexual maturity of sympatric and experimentally allopatric cutthroat trout. and dolly varden charr. Oecologia 61, 319-325

Kaeding LR, Koel TM (2011) Age, growth, maturity, and fecundity of Yellowstone Lake cutthroat trout. NW Sci 85, 431-441

Kelson SJ, Miller MR, Thompson TQ, ORourke SM, Carlson SM (2019) Do genomics and sex predict migration in a partially migratory salmonid fish, *Oncorhynchus mykiss*? Can J Fish Aquat Sci 76, 2080-2088

Kendall NW, McMillan JR, Sloat MR, Buehrens TW, Quinn TP, Pess GR (2015) Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): a review of the processes and patterns. Can J Fish Aquat Sci 72, 319-342

Losee JP, Claiborne AM, Dionne PE, Faulkner HS, Seamons TR (2018) Size, age, growth and site fidelity of anadromous cutthroat trout *Oncorhynchus clarkii clarkii* in the Salish Sea. J Fish Biol 93, 978–987

Love MS (2011) Certainly more than you want to know about the fishes of the Pacific coast. . Really Big Press, Santa Barbara, CA, 650 p

Machacek, H. (ed.), 2006. World Records Freshwater Fishing. www.fishing-worldrecords.com, 12

McMillan JR, Dunham JB, Reeves JH, Mills JS, Jordan CE (2012) Individual condition and stream temperature influence early maturation of rainbow and steelhead trout, *Oncorhynchus mykiss*. Environ Biol Fish 93, 343–355

Meyer, KA, Schill DJ, Elle FS, Lamansky JA Jr (2003) Reproductive demographics and factors that influence length at sex maturity of yellowstone cutthroat trout in Idaho. Trans Am Fish Soc 132, 183-195.

Morita K, Fukuwaka M-a (2007) Why age and size at maturity have changed in Pacific salmon. Mar Ecol Progr Ser 335, 289-294

Morita K, Morita SH, Fukuwaka M, Matsuda H (2005) Rule of age and size at maturity of chum salmon (*Oncorhynchus keta*): implications of recent trends among *Oncorhynchus* spp. Can J Fish Aquat Sci 62, 2752-2759

Morrison CM, Gallagher CP, Tierney KB, Howland KL (2021) Freshwater early life growth influences partial migration in populations of Dolly Varden (*Salvelinus malma malma*). Polar Biol 44, 1353–1364

Morrow, JE (1980) The freshwater fishes of Alaska. Univ Brit Columbia. Anim Res Ecology Library. 248 p

Ogura M, Ishida Y, Ito S (1991) Growth variation of coho salmon *Oncorhynchus kisutch* in the western North Pacific. .Nippon Suisan Gakkaishi 57, 1089-1093

Ohms HA, Sloat MT, Reeves GH, Jordan CE, Dunham JB (2014) Influence of sex, migration distance, and latitude on life history expression in steelhead and rainbow trout (*Oncorhynchus mykiss*). Can J Fish Aquat Sci 71, 70-80

Pearcy WG, Brodeur RD, McKinnell SM, Losee J (2018) Ocean ecology of anadromous coastal cutthroat trout. Chapter 8, pp 905-930 in Ocean Ecology of Pacific Salmon and Trout. Am Fish Soc, Bethesda, MD

Peterman, RM, Bradford MJ, Anderson JL (1986) Environmental and parental influences on age at maturity in sockeye salmon (*Oncorhynchus nerka*) from the Fraser River, British Columbia. Can J Fish Aquat Sci 43, 269-27

Pyper BJ, Peterman RM, Lapointe MF, Walters CJ (1999) Patterns of covariation in length and age at maturity of British Columbia and Alaska sockeye salmon (*Oncorhynchus nerka*) stocks. Can J Fish Aquat Sci 56, 1046-1057

Ricker WE (1995) Trends in the average size of Pacific salmon in Canadian catches. Pp 593-602 in (RJ Beamish, ed), Climate Change and Northern Fish Populations. Can. Spec. Publ. Fish. Aquat. Sci. 121, 739 p

Ruggerone GT, Connors BM (2015) Productivity and life history of sockeye salmon in relation to competition with pink and sockeye salmon in the North Pacific Ocean. Can J Fish Aquat Sci 72, 818–833

Ruiz-Campos G, Pister EP, Compean-Jimenez GA 1997. Age and growth of Nelson's trout, *Oncorhynchus mykiss nelsoni* from Arroyo San Rafael, Sierra San Pedro Martir, Baja California, Mexico. SW Nat 42(1), 74-85.

Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titua RG, Sogard SM, Mangel M (2009) Steelhead life history on California's central coast: insights from a state-dependent model. Trans Am Fish Soc 138, 532-548

Siegel JE, Adkison MD, McPhee MV (2018) Changing maturation reaction norms and the effects of growth history in Alaskan Chinook salmon. Mar Ecol Progr Ser 595, 187-202

Siegel JE, McPhee MV, Adkison MD (2015) Evidence that marine temperatures influence growth and maturation of western Alaskan chinook salmon. Mar Coast Fish: Dyn Manag Ecosys Sci 9, 441-456

Sloat MR, Reeves GH (2014) Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. Can J Fish Aquat Sci 71, 491-501

Thrower FP, Hard JJ, Joyce JE (2004) Genetic architecture of growth and early life-history transitions in anadromous and derived freshwater populations of steelhead. J Fish Biol 65(Suppl A), 286-307

Townsend, C.R. (1989) Population cycles in freshwater fish. J. Fish Biol. 35(Suppl.A):125-131

Trotter PC (1989) Coastal cutthroat trout: a life history compendium. Trans Am Fish Soc 118, 463-473

Underwood TJ, Millard MJ, Thorpe LA (1996) Relative abundance, length frequency, age, and maturity of Dolly Varden in nearshore waters of the Arctic National Wildlife Refuge, Alaska. Trans Am Fish Soc 125, 719-728

Vollestad LA, Peterson J, Quinn TP (2004) Effects of freshwater and marine growth rates on early maturity in male coho and chinook salmon. Trans Am Fish Soc 133, 495-503

Wells BK, Grimes CB, Field JC, Reiss CS (2006) Covariation between the average lengths of mature coho (*Oncorhynchus kisutch*) and Chinook salmon (*O. tshawytscha*) and the ocean environment. Fish Oceanogr 15, 67-79

Zavarina, L.O (2003) Biological structure of chum salmon from the northeastern coast of Kamchatka.. P. 531-540 in (SA Syniakov, NI Naumenko, YP Diakov, OG Zolotov, BB Vronsky, eds.), Condition of Biological Resources of the Northwest Pacific. Petropavlovsk-Kamchatsky, KamchatNIRO.

Zhivotovsky LA, Tochilina TG, Shaikhaev EG, Pogodin VP, Malinina TV, Gharrett AJ (2016) Hybrids between chum *Oncorhynchus keta* and pink *Oncorhynchus gorbuscha* salmon: age, growth and morphology and effects on salmon production. J Fish Biol 89, 2098–2106

Scaridae

Lea RN, Rosenblatt RH (2000) Observations on fishes associated with the 1997-98 El Nino off California. Calif Coop Ocean Fish Invest Rep 41, 117-129

Sciaenidae

Allen LG, Franklin MP (1992) Abundance, distribution, and settlement of young-of-the-year white seabass *Atractoscion nobilis* in the Southern California Bight, 1988-89. Fish Bull 90, 633-641

Blake C, Blake BF (1981) Age determination in six species of fish from a Mexican Pacific coastal lagoon. J Fish Biol 18, 471-478

Clardy SD, Brown-Peterson NJ, Peterson MS, Leaf RT (2014) Age, growth, and reproduction of southern kingfish (*Menticirrhus americanus*): a multivariate comparison with life history patterns in other sciaenids. Fish Bull 112, 178-197

DeMartini EE, Fountain RK (1981) Ovarian cycling frequency and batch fecundity in the queenfish *Seriphus politus*: attributes representative of serial spawning fishes. Fish Bull 79,: 547-560

Donohoe CJ (1997) Age, growth, distribution, and food habits of recently settled white seabass, *Atractoscion nobilis*, off San Diego County, California. Fish Bull 94, 709-721

Gherard KE, Erisman BE, Aburto-Oropeza O, Rowell K, Allen LG (2013) Fishery-dependent estimates of growth, development, and reproduction in Gulf corvina (*Cynoscion othonopterus*). Bull So Calif Acad Sci 112, 1-18

Haydock I (1971) Gonad maturation and hormone-induced spawning of the Gulf croaker, *Bairdiella icistia*. Fish Bull 69, 157-180

Joseph (1962) Growth characteristics of two southern California surf fishes, the California corbina and spotfin croaker, family Sciaenidae. Calif Dep Fish Game, Fish Bull 119, 54p

Leung E, Allen LG (2016) Year-class strength predicts commercial catch 11 years later for white seabass, *Atractoscion nobilis*, off southern California. Calif Fish Game 102, 175-182

Love MS, McGowen GE, Westphal W, Lavenberg RJ, Martin L (1984) Aspects of the life history and fishery of the white croaker, *Genyonemus lineatus* (Sciaenidae), off California. Fish Bull 82, 179-198

Lowerre-Barbieri SK, Chittenden ME Jr, Barbieri LR (1996) Variable spawning activity and annual fecundity of weakfish in Chesapeake Bay. Trans Fish Soc 125, 532-545

Miller EF (2010) Structural irregularities in sagittal otoliths of black croaker (*Cheilotrema saturnum*) from Southern California. Bull So Calif Acad Sci 109, 18-21

Miller EF, Goldberg S, Nunes J, Burkes N, Kuratomi J (2008) The reproductive biology of two common surfzone associated sciaenids, yellowfin croaker (*Umbrina roncadore*) and spotfin croaker (*Roncadore stearnsii*), from Southern California. Bull So Calif Acad Sci 108, 152-159

Miller EF, Pondella DJ II, Allen LG, Herbinson KT (2008) The life-history and ecology of black croaker, *Cheilotrema saturnum*. Calif Coop Oceanic Fish Invest Rep, 49, 191-201

Miller EF, Pondella DJ II, Beck DS, Herbinson KT (2011) Decadal-scale changes in southern California sciaenids under different levels of harvesting pressure ICES J Mar Sci_68, 2123-2133

Miller EF, Williams JP, Pondella DJ II (2011) Queenfish (*Seriphus politus*) and white croaker (*Genyonemus lineatus*) larval growth parameters. Calif Coop Oceanic Fish Invest Rep 52, 75-79

Miller EF, Williams JP, Pondella DJ II, Herbinson-KT (2009) Life history, ecology, and long-term demographics of queenfish. Mar Coast Fish Dynam Ecosys Sci 1, 187-199

Moore SL (2001) Age and growth of white croaker (*Genyonemus lineatus*) off Palos Verdes and Dana Point, California. So Calif Coast Wat Res Project (SCCWRP) Annu Rep, Iss 1999-2000

Pondella DJ II, Froeschke JTZZ, Wetmore LS, Miller E, Valle CL, Medeiros L (2008) Demographic parameters of yellowfin croaker, *Umbrina roncadore* (Perciformes, Sciaenidae), from the Southern California Bight. Pac Sci 62, 555-568

Riedel R, Costa-Pierce BA (2001) Review of the fisheries of the Salton Sea, California, USA: past, present, and future. Rev Fish Sci 9, 239-270, abstract only

Riedel R, Costa-Pierce BA (2002) Review of the fisheries of the Salton Sea, California, USA: past, present, and future. Rev Fish Sci 10, 77-112

Riedel R, Caskey LM, Hurlburt SH (2007) Length-weight relations and growth rates of dominant fishes of the Salton Sea: implications for predation by fish-eating birds. Lake Reserv Manag 23, 528-533

Watson W (1982) Development of eggs and larvae of the white croaker, *Genyonemus lineatus* Ayres (Pisces: Sciaenidae), off the southern California coast. Fish Bull 80, 403-419

Williams JP, Allen LG, Steele MA, Pondella DJ II (2007) El Niño periods increase growth of juvenile white seabass *Atractoscion nobilis* in the Southern California Bight. Mar Biol 152, 193-200

Williams JP, Claiss JT, Pondella DJ II (2012) Patterns of life history and habitat use of an Important recreational fishery species, spotfin croaker, and their potential fishery implications. Mar Coast Fish Dynam, Manag Ecosys Sci 4, 71-84

Scomberosocidae

Hughes SE (1974) Stock composition, growth, mortality, and availability of Pacific saury, *Cololabis saira*, of the Northeastern Pacific Ocean. Fish Bull 72, 121-131

Suyama S, Nakagami M, Naya M, Ueno Y (2012) Comparison of the growth of age-1 Pacific saury *Cololabis saira* in the western and the central North Pacific. Fish Sci 78, 277-285

Suyama S, Oshima K, Nakagami M, Kawabata A (2011) Seasonal changes in otolith and somatic growth in age-0 Pacific saury *Cololabis saira*. Fish Sci 77, 223-233

Suyama S, Sakurai Y, Shimazaki K (1996) Age and growth of Pacific saury *Cololabis saira* (Brevoort) in the western North Pacific Ocean estimated from daily otolith growth increments. Fish Sci 62, 1-7

Watanabe Y, Butler JL, Mori T (1988) Growth of Pacific saury, *Cololabis saira*, in the northeastern and northwestern Pacific Ocean. Fish Bull_86, 489-498

Yatsu A (2019) Review of population dynamics and management of small pelagic fishes around the Japanese Archipelago. Fish Sci 85, 611-639

Scombridae

Aguirre-Villaseñor H, Morales-Bojórquez E, Morán-Angulo RE, Madrid-Vera J, Valdez-Pineda MC (2006xx) Biological indicators for the Pacific sierra (*Scomberomorus sierra*) fishery in the southern Gulf of California, Mexico. *Ciencias Marinas* 32, 471–484

Allman R, Ailloud L, Austin R, Falterman B, Farley J, Krusic-Golub K, Lang E, Pacicco A, Satoh K (2020) Report of the international workshop on the ageing of yellowfin and bigeye tuna. Coll Vol Sci Pap ICCAT 77, 32-46

Andrews AH Pacicco A, Allman R, Falterman BJ, Lang ET, Golet W (2020) Age validation of yellowfin (*Thunnus albacares*) and bigeye (*T. obesus*) tuna of the northwestern Atlantic Ocean. Can J Fish Aquat Sci <https://doi.org/10.1139/cjfas-2019-0328>

Bayliff WH (ed) (1980) Synopses of biological data on eight species of scombrids. Special Scientific Rep No 2, 570 p. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Bayliff WH (1980) Synopsis of biological data on the northern bluefin tuna, *Thunnus thynnus* (Linnaeus, 1758), in the Pacific Ocean. Pp 261-294 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. . Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Black G (1979) Maturity and spawning of the Pacific bonito, *Sarda chiliensis lineolata*, in the eastern North Pacific. Calif Dep Fish Game, Mar Res Tech Rep No 41, 40 p

Bok T, Oray IK (2001) Age and growth of bullet tuna *Auxis rochei* (Risso,1810) in Turkish waters. Coll Vol Sci Pap, ICCAT 52, 8-18

Calkins TP (1980) Synopsis of biological data on the bigeye tuna, *Thunnus obesus* (Lowe, 1839). in the Pacific Ocean. Pp 213-260 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. . Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Castro Hernández, JJ; Santana Ortega, AT (2000) Synopsis of biological data on the chub mackerel (*Scomber japonicus* Houttuyn, 1782). FAO Fish Synopsis. No. 157, 77 p.. Rome, FAO.

Cole JS (1980) Synopsis of biological data on the yellowfin tuna, *Thunnus albacares* (Bonnaterre, 1788), in the Pacific Ocean. Pp 71-150 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. . Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Collins RA, MacCall AD (1977) California's Pacific bonito resource, its status and management. Calif Dep Fish Game, Mar Res Tech Rep No. _35

Farley JH, Williams AJ, Davies CR, Clear NP, Eveson JP, Hoyle SD, Nicol SJ (2012) Population biology of albacore tuna in the Australian region. Final report for FRDC project 2009/012, Wealth from Oceans National Research Flagship, Commonwealth Sci Invest Res Org (CSIRO) Marine and Atmos, Hobart, Australia [3] Secretariat of the Pacific Community, Noumea, New Caledonia. Available at <http://www.frdc.com.au/research/reports>

Foreman TJ (1980) Synopsis of biological data on the albacore tuna, *Thunnus alalunga* (Bonnaterre, 1788), in the Pacific Ocean. Pp 17-70 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. . Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Forsberg ED (1980) Synopsis of biological data on skipjack tuna, *Katsuwonus pelamis* (Linnaeus, 1758), in the Pacific cean. Pp 295-360 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. . Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Gluyas-Millan MG, Castonguay M, Quinonez-Velazquez C (1998) Growth of juvenile Pacific mackerel, *Scomber japonicus* in the Gulf of California. Scientia Marina 62, 225-231

Hogarth WT (1976) Life history aspects of the wahoo *Acanthocybium solandri* (Cuvier & Valenciennes) from the coast of North Carolina. PhD thesis, Dep Zool, Univ No Carolina

Hunter JR, Kimbrell CA (1980) Early life history of Pacific mackerel, *Scomber japonicus*. Fish Bull. 78, 89-101

Ishihara T, Abe O, Shimose T, Takeuchi Y, Aires-da-Silva A (2017) Use of post-bomb radiocarbon dating to validate estimated ages of Pacific bluefin tuna, *Thunnus orientalis*, of the North Pacific Ocean. Fish Res 189, 35–41

Khan MZ (2004) Age and growth, mortality and stock assessment of *Euthynnus affinis* (Cantor) from Maharashtra waters. Indian J Fish 51(2), 210-213

Knaggs,EH, Parrish RH (1973) Maturation and growth of Pacific mackerel *Scomber japonicus* Houttuyn. Calif Fish Game 59, 114-120

Leong, R (1977) Maturation and induced spawning of captive Pacific mackerel, *Scomber japonicus*. Fish Bull 75, 205-211.

Muthiah C (1985) Maturation and spawning of *Euthynnus affinis*, *Auxis thazard*, and *Auxis rochei* in the Mangalore inshore area during 1979-82. Pp 71-85, article 11, in (SE Silas, ed), Tuna Fish Excl Econ Zone India, Biol Stk Assess. Central Mar Fish Res Instit (CMFRI) 36. Indian Council of Agricultural Research, P.B. No. 2704, Cochin 682 031, India

McBride RS, Richardson AK, Maki KL (2008) Age, growth, and mortality of wahoo, *Acanthocybium solandri*, from the Atlantic coast of Florida and the Bahamas. Mar Freshwat Res 59, 799–807

Muhlia-Melo AF (1980) Synopsis of biological data on the black skipjack tuna, *Euthynnus lineatus* Kishinouye, 1920. Pp 361-394 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Oxenford HA, Murray PA, Luckhurst BE (2003) The biology of wahoo (*Acanthocybium solandri*) in the Western Central Atlantic. Gulf Carib Res 15, 33-49

Schaefer KM (1980) Synopsis of biological data on the chub mackerel, *Scomber japonicus* Houttuyn, 1782, in the Pacific Ocean. Pp 395-445 in (WH Bayliff, ed), Synopses of biological data on eight species of scombrids. Special Scientific Rep No 2. Inter-American Tropical Tuna Commission (IATTC), La Jolla, California

Taghavi-Motlagh SA, Hashemi SA, Kochanian P (2010) Population biology and assessment of kawakawa (*Euthynnus affinis*) in coastal waters of the Persian Gulf and Sea of Oman (Hormozgan Province). Iranian J Fish Sci 9(2), 315-326

Tao Y, Mingru C, Jianguo D, Zhenbin L, Shengyun Y (2012) Age and growth changes and population dynamics of the black pomfret (*Parastromateus niger*) and the frigate tuna (*Auxis thazard thazard*), in the Taiwan Strait. Lat. Am. J. Aquat. Res 40, 649-656

Scorpaenidae

Love MS, Axell B, Morris P, Collins R, Brooks A (1987) Life history and fishery of the California scorpionfish, *Scorpaena guttata*, within the Southern California Bight. Fish Bull 85, 99-116

Munk MH, He X, Budrick J (2017) Status of California Scorpionfish (*Scorpaena guttata*) off Southern California in 2017. Pac Fish Manag Council, Portland, OR.
<http://www.pcouncil.org/groundfish/stock-assessments/>

Scyliorhinidae

Balart EF, GonzalezGarcia J, Villavicencio-Garayzar C (2000) Notes on the biology of *Cephalurus cephalus* and *Parmaturus xaniurus* (Chondrichthyes: Scyliorhinidae) from the west coast of Baja California Sur, México. Fish Bull 98, 219-221

Cross JN (1988) Aspects of the biology of two scyliorhinid sharks, *Apristurus brunneus* and *Parmaturus xaniurus*, from the upper continental slope off Southern California. Fish Bull 86, 691-702

Flammang BE, Ebert DA, Cailliet GM (2008) Reproductive biology of deep-sea catsharks (Chondrichthyes: Scyliorhinidae) in the eastern North Pacific. Environ Biol Fish 81, 35-49_s10641-006-9162-9

Nelson DR, Johnson RH (1970) Diel activity rhythms in the nocturnal, bottom-dwelling sharks, *Heterodontus francisci* and *Cephaloscyllium ventriosum*. Copeia 1970,(4), 732-739

- Scytalinidae

Setran DC, Behrens DW (1990) New ecological information on *Scytalina cerdale* (Pisces: Scytalinidae) from a central California rocky intertidal zone. Environ Biol Fish 29, 107-117

Sebastidae

Andrews AH, Kerr LA, Cailliet GM, Brown TA, Lundstrom CC, Stanley RD (2007) Age validation of canary rockfish (*Sebastes pinniger*) using two independent otolith techniques: lead-radium and bomb radiocarbon dating. Mar Freshwat Res 58, 531-541

Armstrong J, Aydin K, Barbeaux S, Bryan M, Cunningham C, Davis O, et al (2018) Appendix B. Stock Assessment and Fishery Evaluation Report for the groundfish resources of the Gulf of Alaska. SAFE Rep, No Pac Fish Manag Council, Portland, OR

Aydin K., Barbeaux S, Clausen D, Conners ME, Courtney D, DiCosimo J, et al (2015) Appendix B. Stock Assessment and Fishery Evaluation Report for the groundfish resources of the Bering Sea/Aleutian Islands regions. SAFE Rep, No Pac Fish Manag Council, Portland, OR

Beckmann AT, Gunderson DR, Miller BS, Buckley RM, Goetz B (1998) Reproductive biology, growth, and natural mortality of Puget Sound rockfish, *Sebastes emphaeus* (Starks, 1911). Fish Bull 96, 352-356

Benet DL, Dick EJ, Pearson DE (1990) Life history aspects of greenspotted rockfish (*Sebastes chlorostictus*) from Central California. NOAA-TM-NMFS-SWFSC-446

Bennett JT, Boehlert GW, Turekian KK (1982) Confirmation of longevity in *Sebastes diploproa* (Pisces: Scorpaenidae) from $^{210}\text{Pb}/^{226}\text{Ra}$ measurements in otoliths. Mar Biol 71, 209-215

Berkel C, Cacan E (2021) Analysis of longevity in Chordata identifies species with exceptional longevity among taxa and points to the evolution of longer lifespans Biogerontology
<https://doi.org/10.1007/s10522-021-09919-w>

Beyer ZZ, Sogard SZ, Harvey ZZ, Field JC (2015) Variability in rockfish (*Sebastes* spp.) fecundity: species contrasts, maternal size effects, and spatial differences. Environ Biol Fish 98, 81-100

Bizzarro JJ, Gilbert-Horvath EA, Dick J, Berger AM, Schmidt KT, Pearson D, Petersen C, Kautzi LA, Miller R, Field JC, Garza JC (2020) Genetic identification of blue rockfish (*Sebastes mystinus*) and deacon rockfish (*S. diaconus*) to enable life history analyses for stock assessment. Fish Bull 118, 37-50

Boehlert GW (1980) Size composition, age composition, and growth of canary rockfish, *Sebastes pinniger*, and splitnose rockfish, *S. diploproa*, from the 1977 rockfish survey. Mar Fish Rev 42, 57-63

Boehlert GW, Kappenman RF (1980) Variation of growth with latitude in two species of rockfish (*Sebastes pinniger* and *S. diploproa*) from the Northeast Pacific Ocean. Mar Ecol Progr Ser 3, p1-10

Boehlert GW, Yoklavich MM, Chelton DB (1989) Time series of growth in the genus *Sebastes* from the Northeast Pacific Ocean. Fish Bull 87, 791-806

Butler JL, Jacobson LD, -Barnes JT, Moser HG (2003) Biology and population dynamics of cowcod (*Sebastes levis*) in the southern California Bight. Fish Bull 101, 260-280

Chen L-C (1971) Systematics, variation, distribution, and biology of rockfishes of the subgenus *Sebastomus* (Pisces, Scorpaenidae, *Sebastes*). Bull Scripps Inst Oceanogr 18, 115 p

Claisse JT, Love MS, Meyer-Gutbrod EL, Williams CM, Pondella DA II (2019) Fishes with high reproductive output potential on California offshore oil and gas platforms. Bull Mar Sci 95, 515-534

Clausen, DM, Heifetz J (2002) The northern rockfish, *Sebastes polyspinis*, in Alaska: commercial fishery, distribution, and biology. Mar Fish Rev 64, 1-28

Coates J, Gunderson DR, LaFrance L, Miller BSZ, Goetz B, Palsson WA (2007) Changes in growth and recruitment of the Puget Sound Rockfish (*Sebastes emphaeus*) in northern Puget Sound. Pp 223-236 in Management of North Pacific Rockfishes Heifetz J, DiCosimo J, Gharrett AJ, Love MS, O'Connell VM, Stanley RD, eds). Procs Symp Biol, Assess Manag No Pac Rockfishes, September 13-15, 2005, Univ AK, Fairbanks, Alaska Sea Grant College Program AK-SG-07-01

Colton MA, Larson RJ (2007) Aspects of the life history of treefish, *Sebastes serriceps* (Sebastidae). Calif Coop Oceanic Fish Invest Rep 48, 177-190

Conrath CL (2017) Maturity, spawning omission, and reproductive complexity of deepwater rockfish. Trans Am Fish Soc 146, 495-507

Conrath CL (2019) Reproductive potential of light dusky rockfish (*Sebastes variabilis*) and northern rockfish (*S. polyspinis*) in the Gulf of Alaska. Fish Bull 117, 140-150

Conrath CL, Hulson P-JF (2021) Temporal variability in the reproductive parameters of deepwater rockfishes in the Gulf of Alaska. Fish Res_nnn,pyyy-zzz

Conrath CL, Knoth B (2013) Reproductive biology of Pacific Ocean Perch in the Gulf of Alaska. Mar Coast Fish: Dyn Manag Ecosys Sci 5, 21-27

Conrath CL, Rooper CN, Wilborn RE, Knoth BA, Jones DT (2019) Seasonal habitat use and community structure of rockfishes in the Gulf of Alaska. Fish Res
<https://doi.org/10.1016/j.fishres.2019.105331>

Dick EJ, Berger A, Bizzarro J, Bosley K, Cope J, Field J, et al (2017) The combined status of blue and deacon rockfishes in U.S. waters off California and Oregon in 2017. Pacific Fishery Management Council, Portland, OR. <http://www.pcouncil.org/groundfish/stock-assessments/>

Dick EJ, Beyer S, Mangel M, Ralston S (2017) A meta-analysis of fecundity in rockfishes (genus *Sebastes*). Fish Res 187, 73–85

Dick EJ, and other Plan Team members (2018) Status of the Pacific coast groundfish fishery. Stock Assessment and Fishery Evaluation (SAFE): Description of the Fishery. Pac Fish Manag Council, Portland OR

Drake JS, Berntson EA, Cope JM, Gustafson RG, Holmes EE, Levin PS, Tolimieri N, Waples,R, Sogard SM, Williams GD. (2010) Status review of five rockfish species in Puget Sound, Washington: bocaccio (*Sebastes paucispinis*), canary rockfish (*S. pinniger*), yelloweye rockfish (*S. ruberrimus*), greenstriped rockfish (*S. elongatus*),and redstripe rockfish (*S. proriger*). NOAA Tech Memo NMFS-NWFSC-108, 234 p

Echeverria TW (1986) Sexual dimorphism in four species of rockfish genus *Sebastes* (Scorpaenidae). Environ Biol Fish 15, 181-190

Echeverria TW (1987) Thirty-four species of California rockfishes: maturity and seasonality of reproduction. Fish Bull 85, 229-250

Eldridge MB, Whipple JA, Bowers MJ, Jarvis BM, Gold J (1991) Reproductive performance of yellowtail rockfish, *Sebastes flavidus*. Environ Biol Fish 30, 91-102

Field JC, Dick EJ, MacCall AD (2007) Stock assessment model for the shortbelly rockfish, *Sebastes jordani*, in the California Current. NOAA-Tech Memo TM-NMFS-SWFSC-405

Field JC, Dick EJ, Pearson D, MacCall AD (2009) Status of bocaccio, *Sebastes paucispinis*, in the Conception, Monterey and Eureka INPFC areas for 2009. Groundfish Analysis Team, Fish Ecol Div, SW Fish Sci Cent, 110 Shaffer Rd, Santa Cruz CA 9506

Field-MacCall A, Ralston S, Love MS, Miller BS (2010) Bocaccionomics: the effectiveness of pre-recruit indices for assessment and management of bocaccio. Calif Coop Oceanic Fish Invest Rep 51, 77-90

Fields RT (2016) Spatial and temporal variation in rosy rockfish (*Sebastes rosaceus*) life history traits. MSc thesis, Calif State Univ, Monterey Bay

Frale BW, Wagman DW, Frierson TN, Aguilar A., Sidlauskas BL (2015). A new species of *Sebastes* (Scorpaeniformes: Sebastidae) from the northeastern Pacific, with a redescription of the blue rockfish, *S. mystinus* (Jordan and Gilbert, 1881). Fish Bull, 113, 355-357

Frey PH, Head MA, Keller AA (2015) Maturity and growth of darkblotched rockfish, *Sebastes crameri*, along the U.S. west coast. Environ Biol Fish 98, 2353-2365

Gallagher MB, Heppell SS (2010) Essential habitat Identification for age-0 rockfish along the Central Oregon Coast. Mar Coast Fish: Dyn Manag Ecosys Sci 2, 60-72

Gertseva VV, Cope JM (2011) Population dynamics of splitnose rockfish (*Sebastes diploproa*) in the Northeast Pacific Ocean. Ecol Model 222, 973-981

Gertseva VV, -Cope JM, Matson SE (2010) Growth variability in the splitnose rockfish *Sebastes diploproa* of the northeast Pacific Ocean: pattern revisited. Mar Ecol Progr Ser 413, 125-136

Gharrett AJ, Matala AP, Peterson EP, Gray AK, Li Z (2005) Two genetically distinct forms of rougheye rockfish are different species. Trans Am Fish Soc 134, 242-260

Gharrett AJ, Mecklenburg CW, Seeb LW, Li Z, Matala AP, Gray, AK, Heifetz J (2006) Do genetically distinct rougheye Rockfish sibling species differ phenotypically? Trans Am Fish Soc 135, 792-800

Gunderson DR, -Zimmerman M, Nichol DG, Pearson K (2003) Indirect estimates of natural mortality rate for arrowtooth flounder (*Atheresthes stomias*) and darkblotched rockfish (*Sebastes crameri*). Fish Bull 101, 175-182

Haldorson L, Love M (1991) Maturity and fecundity in the rockfishes, *Sebastes* spp., a review. Mar Fish Rev 53, 25-31

Hannah RW (2014) Length and age at maturity of female copper rockfish (*Sebastes caurinus*) from Oregon waters based on histological evaluation of ovaries OR Dep Fish Wildlf, Info Rep 2014-04

Hannah RW, Blume MTO (2011) Maturity of female quillback (*Sebastes maliger*) and china rockfish (*S. nebulosus*) from Oregon waters based on histological evaluation of ovaries. OR Dep Fish Wildlf, Info Rep 2011-01

Hannah RW, Blume MTO, Thompson JE (2009) Length and age at maturity of female yelloweye rockfish (*Sebastes rubberimus*) and cabezon (*Scorpaenichthys marmoratus*) from Oregon waters based on histological evaluation of maturity. OR Dep Fish Wildlf, Info Rep 2009-04

Hannah RW, Kautzi LA (2012) Age, growth and female maturity of vermilion rockfish (*Sebastes miniatus*) from Oregon waters. OR Dep Fish Wildlf, Info Rep 2012-05

Hannah RW, Kautzi LA (2015) Age, growth and female length and age at maturity of redbanded rockfish (*Sebastes babcocki*) from Oregon waters. OR Dep Fish Wildlf, Info Rep 2015-03

Hannah RW, Wagman DW, Kautzi LA (2015) Cryptic speciation in the blue rockfish (*Sebastes mystinus*): age, growth and female maturity of the blue-sided rockfish, a newly identified species, from Oregon waters. OR Dep Fish Wildlf, Info Rep 2015-01

Harris JP, Hutchinson C, Wildes S (2019) Using otolith morphometric analysis to improve species discrimination of blackspotted rockfish (*Sebastes melanostictus*) and rougheye rockfish (*S. aleutianus*). Fish Bull, 117, 234-244

Hattori T, Narimatsu Y, Ito M, Ueda Y, Fujiwara K, Kitagawa D (2007) Growth changes in bighead thornyhead *Sebastolobus macrochir* off the Pacific coast of northern Honshu, Japan. Fish Sci 73, 341-343

He X, Pearson DE, Dick EJ, Field JC, Ralston S, MacCall AD (2007) Status of the widow rockfish resource in 2007, Final Rep. NMFS, SW Fish Sci Cent, Fish Ecol Div, Santa Cruz, CA 95060

Head MA, Cope JL, Wulfinch MH (2020) Applying a flexible spline model to estimate functional maturity and spatio-temporal variability in aurora rockfish (*Sebastes aurora*). Environ Biol Fish 103, 1199-1216

Heifetz J, Dicosimo J, Gharrett AJ, Love MS, OConnell VM, Stanley RD (eds) (2005) Proc Symp Biol Assess Manag No Pac Rockfishes, September 13-15, 2005, Univ AK, Fairbanks, Alaska Sea Grant College Program AK-SG-07-01

Helser TE, -Lai H-L, Stewart IJ (2004) A Bayesian hierarchical meta-analysis of growth for the genus *Sebastes* in the Eastern Pacific Ocean. ICES Annual Science Conference on Dynamics and Exploitation of Living Resources, Vigo, Spain, Sept. 22-25, 2004 Marine Resources, ICES C.M. 2004/K:21

Holder AM, Field JC (2019) An exploration of factors that relate to the occurrence of multiple brooding in rockfishes (*Sebastes* spp.). Fish Bull 117, 180-188

Hutchinson CE, Kestelle CR, Kimura DK (2007) Using radiometric ages to develop conventional ageing methods for shorttraker rockfish (*Sebastes borealis*). Pp 237-249 in Heifetz J, Dicosimo J, Gharrett AJ, Love MS, OConnell VM, Stanley RD, eds), Proc Symp Biol Assess Manag No Pac Rockfishes, September 13-15, 2005, Univ AK, Fairbanks, Alaska Sea Grant College Program AK-SG-07-01

Hyde JR, -Kimbrell CA, Budrick JE, Lynn EA, Vetter RD (2008) Cryptic speciation in the vermilion rockfish (*Sebastes miniatus*) and the role of bathymetry in the speciation process. Molec Ecol 17, xxx-xxx

Hyde JR, Vetter RD (2009) Population genetic structure in the redefined vermilion rockfish (*Sebastes miniatus*) indicates limited larval dispersal and reveals natural management units. Can J Fish Aquat Sci 66, 1569-1581

Jacobson LD, Vetter RD (1996) Bathymetric demography and niche separation of thornyhead rockfish: *Sebastolobus alascanus* and *Sebastolobus altivelis*. Can J Fish Aquat Sci 53, 600-609

Johansson ML, Banks MA, Glunt KD, Hassel-Finnegan HM, Buonaccorsi VP (2008) Influence of habitat discontinuity, geographical distance, and oceanography on fine-scale population genetic structure of copper rockfish (*Sebastes caurinus*). Molec Ecol 17, 3051-3061

Jones MK, Mulligan T (2014) Juvenile rockfish recruitment in Trinidad Bay, California. Trans Am Fish Soc 143, 543-551

Jorgensen SJ, Kaplan DM, Klimley AP, Morgan SG, OFarrell MR, Botsford LW (2006) Limited movement in blue rockfish *Sebastes mystinus*: internal structure of home range. Mar Ecol Progr Ser 327, 157-170

Kestelle CR, Helser TE, Wischniowski SG, Loher T, Goetz BJ, Kautzi LA (2016) Incorporation of bomb-produced ¹⁴C into fish otoliths: a novel approach for evaluating age validation and bias with an application to yellowfin sole and northern rockfish. Ecol Model 320, 79-91

Keller AA, Frey PH, Wallace JR, Head MA, Wetzel CR, Cope JM, Harms JH (2018) Canary rockfishes *Sebastes pinniger* return from the brink: catch, distribution and life history along the US west coast (Washington to California). Mar Ecol Prog Ser 599, 181-200

Keller AA, Harms JH, Wallace JR, Jones C, -Benante JA, Chappell A (2019) Changes in long-lived rockfishes after more than a decade of protection within California's largest marine reserve. Mar Ecol Prog Ser 623, 175-193

Keller AA, Molton KJ, Hicks AC, Haltuch M, Wetzel C (2012) Variation in age and growth of greenstriped rockfish (*Sebastes elongatus*) along the U.S. west coast (Washington to California). Fish Res 119, 80-88

Key M, MacCall AD, Bishop T, Leos B (2005) Stock assessment of the gopher rockfish (*Sebastes carnatus*). First Assessment for [stock No of Pt Conception of] the species, prep by the Calif Dep Fish Game and NMFS, SW Fish Sci Center, Santa Cruz Laboratory

Key M, MacCall AD, Field J, Aseltine-Neilson D, Lynn K (2008) The 2007 Assessment of Blue Rockfish (*Sebastes mystinus*) in California. SAFE Rep. for West Pac Council, prep by California Department of Fish & Game, c/o NMFS/SWFSC, Santa Cruz Lab and by NMFS, SW Fish Sci Center, Fish Ecol Div, Santa Cruz Lab

Kline DE (1996) Radiochemical age verification for two deep-sea rockfishes *Sebastolobus altivelis* and *S. alascanus*. MSc thesis, San Jose State Univ

Kolora RR, Owens GL, Vazquez JM, Stubbs A, Chatla K, Jainese C, Seeto K, , McCrea M, Sandel MW, Vianna JA, , Maslenikov K, Bachtrog D, Orr JW, Love M, Sudmant PH (2021) Origins and evolution of extreme life span in Pacific Ocean rockfishes. Science 374, 842-847

Koya Y, Hamatsu T, Matsubara T (1995) Annual reproductive cycle and spawning characteristics of the female kichiji rockfish *Sebastolobus macrochir*. Fish Sci 61, 203-208

Laidig TE, Pearson DE, Sinclair LL (2003) Age and growth of blue rockfish (*Sebastes mystinus*) from central and northern California Fish Bull 101, 800-808

Laidig TE, Ralston S (1995) The potential use of otolith characters in identifying larval rockfish (*Sebastes* spp.). Fish Bull 93, 166-171

Laidig TE, Ralston S, Bence JR (1991) Dynamics of growth in the early life history of shortbelly rockfish *Sebastes jordani*. Fish Bull 89, 611-621

Laidig TE, Sakuma KM (1998) Description of pelagic larval and juvenile grass rockfish, *Sebastes rastrelliger* (family Scorpaenidae), with an examination of age and growth. Fish Bull 96, 788-796

Laidig TE, Sakuma KM, Hyde JR, Watson W, Lawley CT (2008) Identification, description, and daily growth of pelagic larval and juvenile squarespot rockfish, *Sebastes hopkinsi* (family Sebastidae) Calif Coop Oceanic Fish Invest Rep 49, 212-221

Laidig TE, Sakuma KM, Nishimoto MM (1996) Description of pelagic larval and juvenile *Sebastes saxicola* (family Scorpaenidae), with an examination of larval growth. Fish Bull 94, 289-299

Laidig TE, Sakuma KM, Stannard JA (2004) Description and growth of larval and pelagic juvenile pygmy rockfish (*Sebastes wilsoni*) (family Sebastidae). Fish Bull 102, 452-463

Lea RN, McAllister RD, venTresca DA (1999) Biological aspects of nearshore rockfishes of the genus *Sebastes* from Central California with notes on ecologically related sport fishes. Calif Dep Fish Game, Fish Bull 177, 212p

Lefebvre LS, Field JC (2015) Reproductive complexity in a long-lived deepwater fish, the blackgill rockfish. *Trans Am Fish Soc* 144, 383-399

Lefebvre LS, Sabrina G. Beyer SG, David M. Stafford DM, Neosha S. Kashef NS, Dick EJ, Sogard SM, Field JC (2018) Double or nothing: plasticity in reproductive output in the chilipepper rockfish (*Sebastes goodei*). *Fish Res* 204, 258-268

Lenarz WH, Echeverria TW (1991) Sexual dimorphism in *Sebastes*. *Environ Biol Fish* 30, 71-80

Love MS, Caselle JE, Herbinson KA (1998) Declines in nearshore rockfish recruitment and populations in the southern California Bight as measured by impingement rates in coastal electrical power generating stations. *Fish Bull* 96, 492-501

Love MS, Johnson K (1998) Aspects of the life histories of grass rockfish, *Sebastes rastrelliger*, and brown rockfish, *S. auriculatus*, from southern California. *Fish Bull* 87, 100-109

Love MS, McCrea M, Kui L (2018) Aspects of the life histories of pinkrose rockfish (*Sebastes simulator*) and swordspine rockfish (*Sebastes ensifer*) with notes on the subgenus *Sebastomus*. *Bull So Calif Acad Sci* 117, 64-76

Love MS, Morris P, McCrae M, Collins R (1990) Life history aspects of 19 rockfish species (Scorpaenidae: *Sebastes*) from the Southern California Bight. NOAA Tech Rep NMFS 87, 38p

Love MS, Westphal WV (1981) Growth, reproduction, and food habits of olive rockfish, *Sebastes serranoides*, off central California. *Fish Bull* 79, 533-545

Love MS, Yoklavich M, Thorsteinson L (2002) The Rockfishes of the Northeast Pacific. Univ Calif Press, Berkeley & Los Angeles, CA. 404 pp

MacCall AD [retired] (2005, unpubl ms) Assessment of vermilion rockfish in southern and northern California (draft VR_00781). NOAA, NMFS, SW Fish Sci Center, Fish Ecol Div, 110 Shaffer Rd, Santa Cruz, CA 95060

MacCall A [retired], Field J, Pearson D (2009, unpubl ms) Life history and assessment of the warthog (bronzespotted rockfish, *Sebastes gilli*), draftv8b. NMFS, SW Fish Sci Cent, Fish Ecol Div, 110 Shaffer Rd, Santa Cruz, CA 95060

Malecha PW, Hanselman DH, Heifetz J (2007) Growth and mortality of rockfishes (Scorpaenidae) From Alaska waters. NOAA Tech Memo NMFS-AFSC-172, 61 p

Markel RW, Lotterhos KE, Robinson CLK (2017) Temporal variability in the environmental and geographic predictors of spatial-recruitment in nearshore rockfishes. *Mar Ecol Progr Ser* 574, 97-111

Matta ME, Helser TE, Black BA (2018) Intrinsic and environmental drivers of growth in an Alaskan rockfish: an otolith biochronology approach. *Environ Biol Fish* 101, 1571-1587.

Matthews KR (1989) A comparative study of habitat use by young-of-the-year, subadult, and adult rockfishes on four habitat types in central Puget Sound. *Fish Bull* 88, 223-239

Meyers-Cherry NL, Nakamura R, Ruttenberg BI, Wendt DE (2016) Gopher rockfish (*Sebastes carnatus*) life history in south-central California. *Calif Coop Oceanic Fish Invest Rep* 57, 63-70

Miller JA, Shanks AL (2004) Evidence for limited larval dispersal in black rockfish (*Sebastes melanops*): implications for population structure and marine-reserve design. *Can J Fish Aquat Sci* 61, 1723-1735

Munk KM (2001) Maximum ages of groundfishes in waters off Alaska and British Columbia and considerations of age determination. *Alaska Fish Res Bull* 8, 12-21

National Marine Fisheries Service (NMFS). 2013. Groundfish essential fish habitat synthesis: a report to the Pacific Fishery Management Council. NOAA NMFS NW Fish Sci Center, Seattle, WA, April 2013. 107 p

O'Farrell MR, Yoklavich MM, Love MS (1998) Assessment of habitat and predator effects on dwarf rockfishes (*Sebastes* spp.) using multi model inference. *Environ Biol Fish* 85, 239-250

Orr JW, Blackburn JE (2004) The dusky rockfishes (Teleostei: Scorpaeniformes) of the North Pacific Ocean: resurrection of *Sebastes variabilis* (Pallas, 1814) and a redescription of *Sebastes ciliatus* (Tilesius, 1813). *Fish Bull* 102, 328-348

Orr JW, Hawkins S (2008) Species of the rougheye rockfish complex: resurrection of *Sebastes melanostictus* (Matsubara, 1934) and a redescription of *Sebastes aleutianus* (Jordan and Evermann, 1898)(Teleostei: Scorpaeniformes). *Fish Bull* 106, 111-134

Pacific Fish Management Council (2018) Status of the Pacific coast groundfish fishery. Stock Assessment and Fishery Evaluation, Description of the fishery. Pac Fish Manag Council, Portland, OR 97220

Pacific States Marine Fisheries Commission (1999) 52nd Annual Report of the Pacific States Marine Fisheries Commission for the year 1999.

Palsson WA (2001) The development of criteria for establishing and monitoring no-take refuges for rockfishes and other rocky habitat fishes in Puget Sound. *Wash Dep Fish Wildlf, Puget Sound Research* 2001, 10 p

Parker SJ, Berkeley SA, Golden JT, Gunderson DR, Heifetz J, Hixon MA, Larson R, Leaman BM, Love MS, Musick J, O'Connell VM, Ralston S, Weeks HJ, Yoklavich MM (2000). Management of Pacific rockfish *Fisheries* 25, 22-30

Pavoine S, Love MS, Bonsall MB (2009) Hierarchical partitioning of evolutionary and ecological patterns in the organization of phylogenetically-structured species assemblages: application to rockfish (genus: *Sebastes*) in the Southern California Bight. *Ecol Lett* 12, 898-908

Pearson KE, Gunderson DR (2003) Reproductive biology and ecology of shortspine thornyhead rockfish, *Sebastolobus alascanus*, and longspine thornyhead rockfish, *S. altivelis*, from the northeastern Pacific Ocean. *Environ Biol Fish* 67, 117-136

Pearson DE, Hightower JE (1991) Spatial and temporal variability in growth of widow rockfish (*Sebastes entomelas*). NOAA Tech. Memo. NMFS-SWFSC-167

Pearson DE, Hightower JE, Chan JTH (1991) Age, growth, and potential yield for shortbelly rockfish. *FishBull* 89, 403-409

Pena BJ, Allen LG (2017) Size structure and growth rates of vermilion rockfish, *Sebastes miniatus*, along California coast. Abstract Contrib Pap, 98th Annu Mtng, Western Society Naturalists (WSN), Pasadena, Calif, Nov16-19,2017

Phillips(1964) Life history studies on ten species of rockfish (genus *Sebastes*). *Calif Fish Game*, *Fish Bull* 126, 1-70

Piner KR, Wallace JR, Hamel OS, Mikus R (2006) Evaluation of ageing accuracy of bocaccio (*Sebastes paucispinis*) rockfish using bomb radiocarbon. *Fish Res* 77, 200-206

Poltev YN (2011) On reproduction of longfin thornyhead *Sebastolobus macrochir* (Scorpaeniformes: Sebastidae) in Waters off the eastern coast of the Kurils. *J Ichthyol* 51, 324-330

Rasmuson LK, Rankin PS, Kautzi LA, Berger A, Blume MTO, Lawrence KA, Bosley K (2021) Cross-shelf variability of deacon rockfish *Sebastes diaconus*: age, growth, and maturity in Oregon waters and their effect on stock status. *Mar Coast Fish: Dyn Manag Ecosys Sci* 13, 379-395

Rocha-Olivares A, Vetter RD (1999) Effects of oceanographic circulation on the gene flow, genetic structure, and phylogeography of the rosethorn rockfish (*Sebastes helvomaculatus*). *Can J Fish Aquat Sci* 56, 803-813

Rodgveller CR, Lunsford CJ, Fujioka JT (2012) Effects of maternal age and size on embryonic energy reserves, developmental timing, and fecundity in quillback rockfish (*Sebastes maliger*). *Fish Bull* 110, 36-45

Rodriguez-Santiago MA, Rosales-Casian JA (2008) Abundance and size composition of vermilion rockfish, *Sebastes miniatus* (Jordan and Gilbert 1880), from sport fishing catches of San Quintín, Ensenada, Baja California, Mexico. *Bull So Calif Acad Sci* 107, 25-32

Rogers JB, Pikitch EK (1989) Life History Characteristics of Commercially Important groundfish species off California, Oregon, and Washington. NOAA Tech Memo FNWC-166, 39 p

Sakuma KM, Laidig TE (1995) Description of larval and pelagic juvenile chilipepper, *Sebastes goodei* (family Scorpaenidae), with an examination of larval growth. Fish Bull 93, 920-930

Sampson C, Francis C, Field J, Jiao Y, (2013) Stock Assessment Review (STAR) panel report for aurora rockfish. Pacific Fishery Management Council.

Schmidt KT (2014) Life history changes in female blue rockfish, *Sebastes mystinus*, before and after overfishing, in Central California. MSc thesis, Calif State Univ, Monterey Bay

Shaw FR, Gunderson DR (2008) Notes on the life history traits of the rosethorn rockfish, *Sebastes helvomaculatus*. Calif Fish Game 94, 123-136

Sivasundar A, Palumbi SR (2010) Life history, ecology and the biogeography of strong genetic breaks among 15 species of Pacific west coast rockfish, *Sebastes*. Mar Biol 157, 1433-1452

Six LD, Horton HF (1977) Analysis of age determination methods for yellowtail rockfish, canary rockfish, and black rockfish off Oregon. Fish Bull 75, 405-414

Sogard SM, Berkeley SA, Fisher R (2008) Maternal effects in rockfishes *Sebastes* spp.: a comparison among species. Mar Ecol Progr Ser 360, 227-236

Stanley RD, Kronlund AR (2005) Life history characteristics for silvergray rockfish (*Sebastes brevispinis*) in British Columbia waters and the implications for stock assessment and management. Fish Bull 103, 670-684

Stephens A, MacCall A (2004) A multispecies approach to subsetting logbook data for purposes of estimating CPUE. Fish Res 70, 295-306

Stevens MM, Andrews AH, Cailliet GM, Coale KH (2004) Radiometric validation of age, growth, and longevity for the blackgill rockfish (*Sebastes melanostomus*). Fish Bull 102, 711-722

Studebaker RS (2006) Use of rocky intertidal areas by juvenile *Sebastes* in Northern California. MSc thesis, Humboldt State Univ

Sullivan JY, Shotwell SK, Hanselman DH, Hulson PJF, Williams BC, Yasumiichi EM, Ferris-Sullivan BE (2021) 13. Assessment of the rougheye and blackspotted rockfish stock complex in the Gulf of Alaska. No Pac Fish Manag Council GOA SAFE Rep

Tagart JV, Wallace FR, Ianelli JN (2000) Status of the yellowtail rockfish resource in 2000. Joint WA Dep Fish Wildlf and NMFS Rep

TenBrink T T, Helser TE (2021) Reproductive biology, size, and age structure of harlequin rockfish: spatial analysis of life history traits. *Mar Coast Fish: Dynam Manag Ecosys Sci* 13, 463-477

TenBrink TT, Spencer PD (2013) Reproductive biology of Pacific Ocean Perch and northern rockfish in the Aleutian Islands. *No Am J Fish Manag* 33, 373-383

Thompson JE, Hannah RW (2010) Using cross-dating techniques to validate ages of aurora rockfish (*Sebastes aurora*): estimates of age, growth and female maturity. *Environ Biol Fish* 88, 377-388

Thorson JT, Dorn MW, Hamel OS (2019) Steepness for West Coast rockfishes: results from a twelve-year experiment in iterative regional meta-analysis. *Fish Res*
<https://doi.org/10.1016/j.fishres.2018.03.014>

Tribuzio CA, Coutre K, Echave KB (2017) 16. Assessment of the Other Rockfish stock complex in the Gulf of Alaska. NOAA_17529_DS1

Vaux F, -Rasmuson LK, Kautzi LA, Rankin PS, Blume MTO, Lawrence KA, Bohn S, O'Malley KG (2019) Sex matters: Otolith shape and genomic variation in deacon rockfish (*Sebastes diaconus*). *Ecol Evol* 9, 13153-13173

Watson W, Robertson LL (2004) Development of kelp rockfish, *Sebastes atrovirens* (Jordan and Gilbert 1880), and brown rockfish, *S. auriculatus* (Girard 1854), from birth to pelagic juvenile stage, with notes on early larval development of black-and-yellow rockfish, *S. chrysomelas* (Jordan and Gilbert 1880), reared in the laboratory (Pisces: Sebastidae). *NOAA Prof Pap NMFS* 3, 30 p

Watters DL (1993) Age determination and confirmation from otoliths of the bank rockfish, *Sebastes rufus* (Scorpaenidae). MSc thesis, San Jose State Univ

West JE, Helser TE, ONeill SM (2014) Variation in quillback rockfish (*Sebastes maliger*) growth patterns from oceanic to inland waters of the Salish Sea. *Bull Mar Sci* 90,747-761

Wheeler SG, Anderson TW, Bell TW, Morgan SG, Hobbs JA (2017) Regional productivity predicts individual growth and recruitment of rockfishes in a northern California upwelling system. *Limnol. Oceanogr* 62, 754-767

Wilson, JR (2011) Marine reserves and the management of small-scale fisheries. PhD in Environ Sci Manag, Univ Calif Santa Barbara

Wilson JR, Kay MC, Colgate J, Qi R, Lenihan HS (2012) Small-scale spatial variation in population dynamics and fishermen response in a coastal marine fishery. *PLoS ONE* 7(12), e52837

Wilson JR, Prince JD, Lenihan HS (2010) A management strategy for sedentary nearshore species that uses Marine Protected Areas as a reference. *Mar Coast Fish: Dyn Manag Ecosys Sci* 2, 14-27

Yoklavich MM, Love MS, Forney KA (2007) A fishery-independent assessment of an overfished rockfish stock, cowcod (*Sebastes levis*) using direct observations from an occupied submersible. *Can J Fish Aquat Sci* 64, 1795-1804

Serranidae

Allen LG, Block HE (2012) Planktonic larval duration, settlement, and growth rates of the young-of-the-year of two sand basses (*Paralabrax nebulifer* and *P. maculatofasciatus*: fam. Serranidae) from Southern California. *Bull So Calif Acad Sci* 111, 15–21

Allen LG, Hovey TE, Love MS, Smith JTW (1995) The life history of the spotted sand bass (*Paralabrax maculatofasciatus*) within the Southern California Bight. *Calif Coop Oceanic Fish Invest Rep* 36, 193-203

Avendano-Ibarra RA, De Silva-DaVila R, Hernandez-Rivas ME (2009) Reproductive Strategies of Sea Basses based on Larval Abundance in Magdalena Bay, Mexico, 1982–1986. *No Am J Fish Manag* 29, 205-215

Butler-Moser-et al (1982) Developmental stages of three California sea basses (*Paralabrax*, Pisces, Serranidae). *Calif Coop Oceanic Fish Invest Rep* 23, 252-268

Cordes JF, Allen LG (1997) Estimates of age, growth, and settlement from otoliths of young-of-the-year kelp bass (*Paralabrax clathratus*). *Bull So Calif Acad Sci* 96, 43-60

Erisman BE, Allen LG (2006) Reproductive behaviour of a temperate serranid fish, *Paralabrax clathratus* (Girard), from Santa Catalina Island, California, U.S.A. *J Fish Biol* 68, 157-184

Erisman BE, Allen LG, Claisse JT, Pondella DJ II, Miller EF, Murray JH (2011) The illusion of plenty: hyperstability masks collapses in two recreational fisheries that target fish spawning aggregations. *Can J Fish Aquat Sci* 68, 1705-1716

Erisman BE, Cota-Nieto JJ, Moreno-Baez M, Aburto-Oropeza O (2017) Vulnerability of spawning aggregations of a coastal marine fish to a small-scale fishery. *Mar Biol* 164, 100-1XX_s00227-017-3135-8

Findlay AM, Allen LG (2002) Temporal patterns of settlement in the temperate reef fish *Paralabrax clathratus*. *Mar Ecol Progr Ser* 238, 237-248

Hovey TE, Allen LG (2000) Reproductive patterns of six populations of the spotted sand bass, *Paralabrax maculatofasciatus*, from Southern and Baja California. *Copeia* 2000(2), 459-468

Hovey CB, Allen LG, Hovey TE (2002) The reproductive pattern of barred sand bass (*Paralabrax nebulifer*) from Southern California. Calif Coop Oceanic Fish Invest Rep 43, 174-181

Jarvis ET, Loke-Sith KA, Evans K, Kloppe RE, Young KA, Valle CF (2014) Reproductive potential and spawning periodicity in barred sand bass (*Paralabrax nebulifer*) from the San Pedro Shelf, Southern California. Calif Fish Game 100, p289-309

Love MS, Brooks A, Bussatto, Stephens J, Gregory PA (1996) Aspects of the life histories of the kelp bass, *Paralabrax clathratus*, and barred sand bass, *R nebulifer*; from the southern California Bight. Fish Bull 94, 472-481

Miller EF, Erisman B (2014) Long-term trends of Southern California's kelp and barred sand bass populations: a fishery-independent assessment. Calif Coop Fish Invest Rep 55, 119-127

Oda DL, Lavenberg RJ, Rounds JM (1993) Reproductive biology of three California species of *Paralabrax* (Pisces: Serranidae). Calif Coop Fish Invest Rep 34, 122-132

Paterson CN, Chabot CL, Robertson JM, CoteNieto JJ, Erisman B, Allen LG (2015) The genetic diversity and population structure of barred sand bass, *Paralabrax nebulifer*: a historically important fisheries species off southern and Baja California. Calif Coop Oceanic Fish Invest Rep 56, 97-109

Pondella DJ II, Allen LG, Craig MT, Ginter B (2006) Evaluation of eelgrass mitigation and fishery enhancement structures in San Diego Bay, California. Bull Mar Sci 78, 115-131

Pondella DJ II, Allen LG, Rsaes Casian JA, Hovey TE (2001) Demographic parameters of golden spotted rock bass *Paralabrax auroguttatus* from the northern Gulf of California. Trans Am Fish Soc 130, 686-691

Tanner RL, Obaza AK, Ginsburg DW (2019) Secondary production of kelp bass *Paralabrax clathratus* in relation to coastal eelgrass *Zostera marina* habitat in a Southern California Marine Protected Area. Bull So Calif Acad Sci 118, 158-172

Teesdale GN, Wolfe BW, -Lowe CG (2015) Patterns of home ranging, site fidelity, and seasonal Spawning migration of barred sand bass caught within the Palos Verdes Shelf Superfund Site. Mar Ecol Prog Ser 539, 255-269

Young PH (1962) The kelp bass (*Paralabrax clathratus*) and its fishery, 1947–1958. Calif Dep Fish Game, Fish Bull 122, 67p

Somniosidae

Courtney DI, Adkison-MD, Sigler MF (2016) Risk analysis of plausible incidental exploitation rates for the Pacific sleeper shark, a data-poor species in the Gulf of Alaska. *No Am J Fish Manag* 36, 523-548

Nielsen J, Hedeholm RB, Heinemeier J, Bushnell PG, Christensen JS, Olsen J, et al (2016) Eye lens radiocarbon reveals centuries of longevity in the Greenland shark (*Somniosus microcephalus*). *Science* 353, 702-704

Yano K, Stevens JD, Compagno LJV (2007) Distribution, reproduction and feeding of the Greenland shark *Somniosus (Somniosus) microcephalus*, with notes on two other sleeper sharks, *Somniosus (Somniosus) pacificus* and *Somniosus (Somniosus) antarcticus*. *J Fish Biol* 70, 374-390

Sparidae

Matus-Nivon E, Ramirez-Sevilla R, Martinez-Pecero R, et al (1989) Description of larval and juvenile Pacific porgy *Calamus brachysomus* Lockington (Pisces, Sparidae). *Invest Mar CICIMAR* 4, 141-150

Sphyraenidae

Bottinelli DJ, Allen LG (2007) A re-evaluation of age, growth, and batch fecundity in the California barracuda, *Sphyraena argentea*, from Southern California based on specimens taken from 2000 to 2002. *Calif Fish Game* 93, 167-199

Pinkas L (1966) A management study of the California barracuda *Sphyraena argentea*. *Calif Dep Fish Game, Fish Bull* 134, 58p

Schultze DL (1981) An evaluation of the California barracuda resource and its management. *Calif Dep Fish Game, Mar Res Admin Rep* 80-6

Walford LA (1932) The California barracuda (*Sphyraena argentea*). *Calif Dep Fish Game, Fish Bull* 37, 72p

Sphyrnidae

Anislado-Tolentino V, Cabello MG, Linares FC, Mendoza CR (2008) Age and growth of the scalloped hammerhead shark, *Sphyrna lewini* (Griffith & Smith, 1834) from the southern coast of Sinaloa, México. *Hidrobiologica* 18, 31-40

Coelho R, Fernandez-Carvalho J, Amorim S, Santos MN (2011) Age and growth of the smooth hammerhead shark, *Sphyrna zygaena*, in the Eastern Equatorial Atlantic Ocean, using vertebral sections. *Aquat Liv Res* 24, 351-357

Drew M., White WT, Dharmadi, Harry AV, Huveneers C (2015) Age, growth and maturity of the pelagic thresher *Alopias pelagicus* and the scalloped hammerhead *Sphyrna lewini*. *J Fish Biol* 86, 333-354

Frazier BS, Driggers WB III, Adams DH, Jones CM, Loefer JK (2014) Validated age, growth and maturity of the bonnethead *Sphyrna tiburo* in the western North Atlantic Ocean. *J Fish Biol* 85, 688-712

Liu K-M, Chen C-T (1999) Demographic analysis of the scalloped hammerhead, *Sphyrna lewini*, in the Northwestern Pacific. *Fish Sci* 65, 218-223

Liu K-M, Tsai WP (2011) Catch and life history parameters of pelagic sharks in the Northwestern Pacific. Keelung, Chinese Taipei, ISC Shark Working Group Workshop. 12pp

Nava P, Marquez-Farias JF (2014) Size at maturity of the smooth hammerhead shark, *Sphyrna zygaena*, captured in the Gulf of California. *Hidrobiologica* 24, 129-135.

Passerotti MS, Carlson JK, Piercy AN, Campana SE (2010) Age validation of great hammerhead shark (*Sphyrna mokarran*), determined by bomb radiocarbon analysis. *Fish Bull* 108, 346–351

Rosa D, Coelho R, Fernandez-Carvalho J, Santos MN (2017) Age and growth of the smooth hammerhead, *Sphyrna zygaena*, in the Atlantic Ocean: comparison with other hammerhead species. *Mar Biol Res* 23, 300-313

Squalidae

DellApa A, Bangley CW, Rulifson RA (2015) Who let the dogfish out? A review of management and socio-economic aspects of spiny dogfish fisheries. *Rev Fish Biol Fish* 25, 273-295

Ebert DA, White WT, 4, Goldman KJ, Compagno LJV, Daly–Engel TS, Ward RD (2010) Resurrection and redescription of *Squalus suckleyi* (Girard, 1854) from the North Pacific, with comments on the *Squalus acanthias* subgroup (Squaliformes: Squalidae). *Zootaxa* 2612, 22–40

Gertseva V, Taylor IG (2011) Status of the spiny dogfish shark resource off the continental U.S. Pacific Coast in 2011. Pac Fish Manag Council, SAFE Rep

Gunderson DR, Dygert PH (1988) Reproductive effort as a predictor of natural mortality rate. *J Cons int Explor Mer* 44, 200-209

Hauser L, Franks JH, Vega N, Gallucci V (2007) Potential for sustainable expansion of the dogfish (*Squalus acanthias*) fishery in the northeast Pacific. Final Rep to NOAA-SK, 36 p

Matta ME, Tribuzio CA, Ebert DA, Goldman KJ, Gburski CM (2017) Age and growth of elasmobranchs and applications to fisheries management and conservation in the Northeast Pacific Ocean. Adv Mar Biol 77, 179-220

McFarlane GA, King JR (2003) Migration patterns of spiny dogfish (*Squalus acanthias*) in the North Pacific Ocean. Fish Bull 101, 358-367

McFarlane GA, McPhie RP, King JR (2010) Distribution and life history parameters of elasmobranch species in British Columbia waters. Can Tech Rep Fish Aquat Sci 2908

Saunders MW, McFarlane GA (1993) Age and length at maturity of the female spiny dogfish, *Squalus acanthias*, in the Strait of Georgia, British Columbia, Canada. Environ Biol Fish 38, 49-57

Taylor IG, Gallucci VF (2009) Unconfounding the effects of climate and density dependence using 60 years of data on spiny dogfish (*Squalus acanthias*). Can J Fish Aquat Sci 66, 351-366

Tribuzio CA (2010) Life history, demography, and ecology of the spiny dogfish "Squalus acanthias" in the Gulf of Alaska. PhD thesis, Univ Alaska, Fairbanks

Tribuzio CA, Echave K, Hulson P, Rodgveller C, Shotwell SK (2012) Evaluation of stock structure for the shark complex in the Gulf of Alaska and Bering Sea/Aleutian Island. BSAI Shark Complex. _Draft SAFE Rep, 29 p. No Pac Fish Manag Council, Portland, OR

Tribuzio CA, Echave K, Rodgveller C, Hulson PJ (2016) Assessment of the shark stock complex in the Bering Sea and Aleutian Islands. BSAI_Shark_Complex_SAFE Rep, 1915-1962. No Pac Fish Manag Council, Portland, OR

Tribuzio, CA, Gallucci VF, Bargmann, GG (2009) A survey of demographics and reproductive biology of spiny dogfish (*Squalus acanthias*) in Puget Sound, WA. Pp 181-194 in (V F Gallucci, GA McFarlane, G Bargmann, eds), Biology and Management of Dogfish Sharks. American Fisheries Society: Bethesda, MD

Tribuzio CA, Kruse GH (2011) Demographic and risk analyses of spiny dogfish (*Squalus suckleyi*) in the Gulf of Alaska using age and stage-based population models. Mar Freshwat Res 62, 1395-1406

Tribuzio CA, Kruse GH (2012) Life history characteristics of a lightly exploited stock of *Squalus suckleyi*. J Fish Biol 80, 1159-1180

Tribuzio CA, Kruse GH, Fujioka JT (2010) Age and growth of spiny dogfish (*Squalus acanthias*) in the Gulf of Alaska: analysis of alternative growth models. Fish Bull 108, 119-135

Tribuzio CA, Matta ME, Gburski C, Atkins N, Bubley W (2017) Methods for the preparation of Pacific spiny dogfish, *Squalus suckleyi*, fin spines and vertebrae and an overview of age determination. Mar Fish Rev 78, 1-13

Tribuzio CA, Matta ME, Gburski C, Blood C, Bubley W, Kruse GH (2018) Are Pacific spiny dogfish lying about their age?: a comparison of ageing structures for *Squalus suckleyi*. Mar Freshwat Res 62, 1395-1406

Squatinae

Cailliet GM, Mollet HF, Pittenger GG, Bedford D., Natanson LJ (1992) Growth and demography of the Pacific angel shark (*Squatina californica*), based upon tag returns off California. Aus J Mar Freshwat Res 43, 1313-1330

Natanson LJ, Cailliet GM (1986) Reproduction and development of the Pacific angel shark, *Squatina californica*, off Santa Barbara, California. Copeia 1986(4), 987-994

Natanson LJ, Cailliet GM (1990) Vertebral growth zone deposition in Pacific angel sharks. Copeia 1990(4), 1133-1145

Romero-Caicedo AF, Galván-Magaña F, Hernández-Herrera A, Carrera-Fernández M (2016) Reproductive parameters of the Pacific angel shark *Squatina californica* (Selachii: Squatinae). J Fish Biol 88, 1430-1440

Sternorhynchidae

Eduardo LN, Mincarone MM, Villarins BT, Frédou T, Lira AS, Bertrand A, Lucena-Frédou F (2019) Length- weight relationships of eleven mesopelagic fishes from oceanic islands of the southwestern Tropical Atlantic. J Appl Ichthyol 35, 605-607

Stichaeidae

Antonenko DV, Pushchina OI, Lachugin PV (2004) Distribution and biological features of the long-snouted blenny *Lumpenus longirostris* (Stichaeidae) in waters of Primorye (the Sea of Japan). J Ichthyol 44, 747-751

Coleman RM (1992) Reproductive biology and female parental care in the cockscomb prickleback, *Anoplopoma fimbria* (Pisces: Stichaeidae). Environ Biol Fish 35, 177-186

Dunbrack R, Green JM (2017) Life history differences and latitudinal variation in recruitment in two species of Arctic-Boreal perciform fishes, the fish doctor *Gymnelus viridis* and the Arctic shanny *Stichaeus punctatus*. J Ichthyol 57, 380-392

Keats-Steele-Green-Martel(1993) Diet and population size structure of the Arctic shanny, *Stichaeus punctatus* (Pisces: Stichaeidae), at sites in eastern Newfoundland and the eastern Arctic. Environ Biol Fish 37, 173-180

Marliave JB, DeMartini EE (1977) Parental behavior of intertidal fishes of the stichaeid genus *Xiphister*. Can J Zool 55, 60-63

Marshall W.H, Wylie-Echeverria T (1992) Age, length, weight, reproductive cycle, and fecundity of the monkeyface prickleback (*Cebidichthys violaceus*). Calif Fish Game 78, 57–64

Meyer- Ottesen CA, Hop H, Falk-Petersen S, Christiansen JS (2014) Reproduction and sexual dimorphism of daubed shanny (Teleostei: *Leptoclinus maculatus*) in Svalbard waters. Polar Biol 37, 809-815

Peppar JL (1965) Some features of the life history of the cockscomb prickleback, *Anoplarchus purpureus* Gill. MSc thesis, Univ British Columbia. 159 pp.

Shiogaki M (1983) On the life history of the stichaeid fish *Chirolophis japonicus*. Jpn J Ichthyol 29, 446-455

Wingert RC (1975) Comparative reproductive cycles and growth histories of two species of *Xiphister* (Pisces: Stichaeidae), from San Simeon, California. MA thesis, Calif State Univ, Fullerton

Stomiidae

Fisher JP, Pearcy WG (1983) Reproduction, growth and feeding of the mesopelagic fish *Tactostoma macropus* (Melanostomiidae) . Mar Biol 74, 257-267

Kawaguchi K, Moser HG (1993) Development and distribution of the early life history stages of the mesopelagic fish *Tactostoma macropus* (Stomiidae) in the transitional waters of the North Pacific. Jpn J Ichthyol 2, 161-172

Orlov A, Binohlan C (2009) Length–weight relationships of deep-sea fishes from the western Bering Sea. J appl Ichthyol 25, 223-227

Stromateidae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Syngnathidae

Bayer RD (1980) Size, seasonality, and sex ratios of the bay pipefish (*Syngnathus leptorhynchus*) in Oregon. Northwest Sci 54, 161-167

Fritzsche RA (1980) Revision of the Eastern Pacific Syngnathidae (Pisces: Syngnathiformes), including both recent and fossil forms. Proc Calif Acad Sci 42, 181-227

Ortega-Salas AA, Reyes-Bustamante H (2006) Fecundity, survival, and growth of the seahorse *Hippocampus ingens* (Pisces: Syngnathidae) under semi-controlled conditions. Rev Biol Trop 54, 1099-1102

Synodontidae

Morales-Nin B (1996) Abundance, distribution and growth of *Synodus evermanni* Jordan and Bollman, 1890 and *Synodus scituliceps* Jordan and Gilbert, 1882 (Pisces: Synodontidae) from the Mexican Pacific shelf. Publ Espec Inst Oceanogr (English Abs)

Tetragonuridae

Love MS (2011) Certainly More than You Want to Know about the Fishes of the Pacific Coast: a Postmodern Experience. Really Big Press, Santa Barbara, CA.650 pp

Triakidae

Au DW, Smith SE (1997) A demographic method with population density compensation for estimating productivity and yield per recruit of the leopard shark (*Triakis semifasciata*). Can J Fish Aquat Sci 54, 415-420

Barker AM, Nosal AP, Lewallen EA, Burton RS (2015) Genetic structure of leopard shark (*Triakis semifasciata*) populations along the Pacific coast of North America. J Exp Mar Biol Ecol 472, 151–157

Cailliet GM (1992) Demography of the Central California population of the leopard shark (*Triakis semifasciata*). Aus J Mar Freshwat Res 43, 183-193

Ebert DA, Ebert TB (2005) Reproduction, diet and habitat use of leopard sharks, *Triakis semifasciata* (Girard), in Humboldt Bay, California, USA). Mar Freshwat Res 56, 1089-1098

Francis MP, Mulligan KP (1998) Age and growth of New Zealand school shark, *Galeorhinus galeus*. NZ J Mar Freshwat Res 32, 427-440

Kusher DA, Smith SE, Cailliet GM (1992) Validated age and growth of the leopard shark, *Triakis semifasciatus*, with comments on reproduction. Environ Biol Fish 35, 187-203

Lewallen EA, Anderson TW, Bohonak AJ (2007) Genetic structure of leopard shark (*Triakis semifasciata*) populations in California waters. Mar Biol 152, 599-609

Moulton PL, Walker TI, Saddler SR (1992) Age and growth studies of gummy shark, *Mustelus antarcticus* Günther, and school shark, *Galeorhinus galeus* (Linnaeus), from southern Australian waters Aus J Mar Freshwat Res 43, 1241-1267

Navia AF, GIRALDO A, Mejia-Falla PA (2006) Notes on the biology and diet of *Mustelus lunulatus* in the central fishing zone in the Pacific Ocean off Colombia. Investigaciones Marinas, Valparaiso 34, 217-222 (in Spanish with Engl Abs)

Nosal AP, Cartamil DP, Ammann AJ, Bellquist LF, Ben-Aderet NJ, Blincow KM, Burns ES, Chapman ED, Freedman RM, Klimley AP, Logan RK, Lowe CG, Semmens BX, White CF, Hastings PA. (2021) Triennial migration and philopatry in the critically endangered soupfin shark *Galeorhinus galeus*. J Appl Ecol J Appl Ecol. 58, 1570-1582

Perez-Jimenez JC, Sosa-Nishizaki O (2010) Determining reproductive parameters for population assessments of two smoothhounds (*Mustelus californicus* and *Mustelus lunulatus*) from the northern Gulf of California, Mexico. Bull Mar Sci 86, 3-13

Pondella DJ II, Allen LG (2008) The decline and recovery of four predatory fishes from the Southern California Bight. Mar Biol 154, 307-313

Ripley WE (1946) The biology of the soupfin shark *Galeorhinus zyopterus* and its fishery. Calif Fish Game Fish Bull 64, 7-37

Russo RA (2019a) Primary and secondary nursery areas for leopard and brown smoothhound sharks in San Francisco Bay, California. Calif Fish Game 105, 21-30

Russo RA (2019b) Determining sexual maturity in male leopard sharks in San Francisco Bay, California. Calif Fish Game 105, 197-204

Smith SE, Abramson NJ (1990) Leopard shark *Triakis semifasciata* distribution, mortality rate, yield, and stock replenishment estimates based on a tagging study in San Francisco Bay. Fish Bull 88, 371-381

Smith SE, Au DW, Show C (1998) Intrinsic rebound potentials of 26 species of Pacific sharks. Mar Freshwat Res 49, 663-678

Smith SE, Mitchell RA, Fuller D (2003) Age-validation of a leopard shark (*Triakis semifasciata*) recaptured after 20 years. FishBull 101, 194-198

Yudin KG, Cailliet GM (1990) Age and Growth of the gray smoothhound, *Mustelus californicus*, and the brown smoothhound, *M. henlei*, sharks from Central California. Copeia 1990(1), 191-204

Trichiuridae

Chen Y-S, Jeng Z-Y, Wang H-Y (2020) Changes in reproduction for subtropical cutlassfish *Trichiurus japonicus* reflect variation in temperature and population demography. Mar Ecol maec.12578.

Mikhaylin SV (1982) Data on the distribution and biology of *Lepidopus xantusi* and *Trichiurus nitens* (Trichiuridae) in the southeastern Pacific. J Ichthyol 22, 24-32

Shih N-T, Hsu K-C, Ni I-H (2011) Age, growth and reproduction of cutlassfishes *Trichiurus* spp. in the southern East China Sea. J appl Ichthyol 27, 1307-1315

Wang H-Y, Heino M (2018) Adaptive and plastic variation in growth and maturation of the cutlassfish *Trichiurus japonicus* in the subtropical Pacific Ocean. Fish Bull 116, 171-182

Trichodontidae

McDonald-Paul-Vogeler, Doyle ZZ (1997) Biological investigations on Pacific sandfish (*Trichodon trichodon*) in the Northern Gulf of Alaska. Pp 87-4 in Forage Fishes in Marine Ecosystems. Am Fish Soc pp 87-94

Triglidae

Schmitter-Soto JJ, Castro-Aguirre JL (1991) Age and growth of *Prionotus stephanophrys* (Osteichthyes: Triglidae) in the western coast of Baja California Sur, Mexico. Rev Biol Trop 39, 23-29

Schmitter-Soto JJ, Castro-Aguirre JL (1994) Age and growth of three searobins (Pisces: Triglidae) off the western coast of Baja California Sur, México Rev Biol Trop 42, 271-279

Schmitter-Soto JJ, Castro-Aguirre JL (1996) Fluctuaciones espacio-temporales del crecimiento en *Prionotus stephanophrys* (Scorpaeniformes: Triglidae) del Pacífico sudcaliforniano, México Rev Biol Trop 44, 271-279

Uranoscopidae

Sutton CP (2004) Estimation of age, growth, and mortality of giant stargazer (*Kathetostoma giganteum*) from Southland trawl surveys between 1993 and 1996. N Z Fish Assess Rep, issue 38

Urolophidae

Babel JS (1967) Reproduction, life history, and ecology of the round stingray, *Urolophus halleri* Cooper. Calif Dep Fish Game, Fish Bull 137, 1-104

Hale LF, Lowe CG (2008) Age and growth of the round stingray *Urolophus halleri* at Seal Beach, California J Fish Biol 73, 510-523

Vaudo JJ, Lowe CG (2006) Movement patterns of the round stingray *Urolophus halleri* (Cooper) near a thermal outfall J Fish Biol 68, 1756

Zaniolepididae

DeMartini EE & Anderson ME (1980) Comparative survivorship and life history of painted greenling (*Oxylebius pictus*) in Puget Sound, Washington and Monterey Bay, California. Environ Biol Fish 5, 44-47

Johnson CR, Adams JP (1970) Biology of *Zaniolepis latipinnis* (Zaniolepididae) in relation to vertical distribution. Copeia 1970(4), 769-771

Xiphiidae

DeMartini EE, Uchiyama JH, Williams HA (2000) Sexual maturity, sex ratio, and size composition of swordfish, *Xiphias gladius*, caught by the Hawaii-based pelagic longline fishery. Fish Bull 98, 489-498

DeMartini EE, Uchiyama JH, Humphreys RL Jr, Sampaga JD, Williams HA (2007) Age and growth of swordfish (*Xiphias gladius*) caught by the Hawaii-based pelagic longline fishery. Fish Bull. 105, 356-367

Sun, C.-L., S.-P. Wang S-P, Yeh S-Z (2002) Age and growth of the swordfish (*Xiphias gladius* L.), in the waters around Taiwan determined from anal fin rays. Fish Bull 100, 822–835

Zaproridae

Smith KR, Somerton DA, Yang M-S, Nichol DG (2004) Distribution and biology of prowfish (*Zaprora silenus*) in the northeast Pacific. Fish Bull 102, 168-178

Zoarcidae

Anderson ME (1980) Aspects of the natural history of the midwater fish *Lycodapus mandibularis* (Zoarcidae) in Monterey Bay, California. Pac Sci 34, 181-194

Dunbrack R, Green JM (2017) Life history differences and latitudinal variation in recruitment in two species of Arctic-Boreal perciform fishes, the fish doctor *Gymnelus viridis* and the Arctic shanny *Stichaeus punctatus*. J Ichthyol 57, 380-392

Ferry LA (1994) Natural history of two-line eelpouts, *Bothrocara brunneum* (Family Zoarcidae). MSc thesis in Mar Biol, San Francisco State Univ

Ferry-Graham LA, Drazen JC, Franklin V (2007) Laboratory observations of reproduction in the deep-water zoarcids *Lycodes cortezianus* and *Lycodapus mandibularis* (Teleostei: Zoarcidae). Pac Sci 61, 129-139

Green JM, -Mitchell LR, Reynolds JB (eds) (1997) Biology of the fish doctor, an eelpout, from Cornwallis Island, Northwest Territories, Canada. Am Fish Soc Symp 19, 140-147

Hildebrandt N, Bergmann M, Knust R (2011) Longevity and growth efficiency of two deep-dwelling Arctic zoarcids and comparison with eight other zoarcid species from different climatic regions. Polar Biol 34, 1523-1533

Lancraft, T.M. 1982. Aspects of the natural history of *Melanostigma pammelas*. (Pisces: Zoarcidae). MA thesis, Univ Calif, Santa Barbara, 137 p.

Matallanas J, Rucabado J, Lloris D, PilarOlivar MP (1990) Early stages of development and reproductive biology of the South-American eelpout *Austrolycus depressiceps* Regan, 1913 (Teleostei: Zoarcidae). Scientia Marina 54, 257-261

Zuercher R, Kliever RG, Cailliet GM (2019) Life history of the deep-water persimmon eelpout (*Eucryphycus californicus*, family: Zoarcidae), and its use of drift vegetation as an ecological subsidy. Environ Biol Fish 102, 1161-1178