

INDEX

- Able, K. W., 139
Abudefduf troschelli (Panamic sergeant major), 92f, 235t, 237f, 241f
 abundance/richness, 5, 9, 81, 547–48; habitat and, 109–12; in kelp bed/rocky reef, 228–32; patterns of, 6–8, 626f, 630; in rocky intertidal zone, 215, 217, 221
 abyssal grenadier (*Coryphaenoides armatus variabilis*), 353t, 369, 372f, 374–76
Acanthochromis polyacanthus, 503
Acanthocybium solandri (wahoo), 333, 337t
Acanthogobius flavimanus (yellowfin goby), 95f, 533t, 612t, 613–14, 618; in bays and estuaries, 126t, 130t, 131, 132f, 142f
 acanthopterygians, 55, 57, 63, 73–74
Acanthurus spp. (surgeonfishes), 329, 523t
 Achiridae (American soles), 175, 179t, 180t
Acipenser spp. (sturgeon), 125, 587, 602; *A. medirostris* (green sturgeon), 18t, 125, 126t, 128f, 130t; *A. transmontanus* (white sturgeon), 18t, 125, 126t, 128f, 518t
 acoustico-lateralis system, 343
 acoustic signals, in courtship, 493
 acoustic telemetry, 527–30, 539
 actinopterygian fishes, 56–57, 63, 172
 activity patterns, 543–49; biological clocks and, 548–49; dawn and dusk changeover and, 548; day-night differences in, 547–48; diel, 188, 507, 544, 545f; egestion and excretion, 546; nocturnal, 545–46. *See also* under feeding; movement
 Adams, P. B., 421
 Adey, W. H., 5–6
 Agonidae (poachers), 9f, 10f, 70, 189, 263t, 505; *A. acipenserinus*, 17t, 21t; on continental shelf, 175, 178t, 180t; reproduction and, 499
Agonomalus mozinoi (kelp poacher), 70
Agonopsis sterletus (southern spearnose poacher), 19t, 24t, 178t, 183f, 196f
Agonopsis vulsa, 17t, 21t
 Ahlstrom, Elbert A., 269, 305
 albacore (*Thunnus alalunga*), 18t, 23t, 89f, 389, 401f; fisheries and, 573–74, 575, 584; in pelagic zone, 326f, 327f, 330, 336t, 337t; seasonal movement of, 536t, 542
Albatrossia pectoralis. *See* *Coryphaenoides pectoralis*
Albula vulpes (bonefish), 20, 135t, 145t, 518t, 532t
Alcichthys alcicornis (elkhorn sculpin), 499, 501–2
Alepisaurus ferox (longnose lancetfish), 327f, 330, 519t
 Alepocephalidae (slickheads), 106t, 349t, 353t, 354; *A. bairdii*, 519t; *A. tenebrosus* (California slickhead), 20t, 21t, 101f, 353t
 Aleutian Province, 3, 4f
 Alevizon, W. S., 457
 algae, 5, 74, 218–19, 220, 329, 422; *Gelidium robustum*, 457, 459–60; *Gracillaria* spp., 111, 159; herbivores and, 394, 403, 404; *Pterygophora californica*, 228, 246, 437–38, 474; *Sargassum* spp., 228, 324, 419, 430, 436f
 alien species, 611–19; in bays and estuaries, 130–31, 142f; Japanese gobies, 130t, 131, 134, 142f, 614, 618–19; overview of, 611–12; rainwater killifish, 130, 131, 134f, 612t, 614; yellowfin goby, 95f, 126t, 130t, 131, 132f, 142f, 612t, 613–14, 618. *See also* American shad; *Morone saxatilis* (striped bass); *specific aliens*
 Allen, Larry G., 5, 9, 75, 81, 109, 506, 540, 541; bay and estuary study, 6, 10, 122, 135, 137–38, 140, 143; on harbors, 158, 159, 163; on pollutants, 601; on surf zone, 151, 153, 154; on trophic relations, 398, 403
 Allen, M. J., 174, 186, 187, 188, 189, 193, 197
Alloclinus holderi (island kelpfish), 19t, 24t, 92f, 162t, 236t, 477, 534t; cleaner behavior of, 555, 556f, 557t
Alloctytus folletti, 17t, 21t
Alloctytus verrucosus, 520t
Allosmerus elongatus, 17t
Allothunnus fallai, 20t, 24t
 almaco jack (*Seriola rivoliana*), 20t, 25t, 62
 Almany, G. R., 435, 437, 443, 444, 445
Alopias superciliosus, 17t, 21t, 518t
Alopias vulpinus (thresher shark), 19t, 233t, 542; bigeye thresher, 89f, 326f, 337t; and commercial fishing, 579, 591
Alosa sapidissima. *See* American shad
Altrichthys spp., 503
 Ambrose, R. F., 135
 ambushers, 59–62, 68, 190, 395
 American eel (*Anguilla rostrata*), 612t, 618
 American Indians, 624, 627
 American shad (*Alosa sapidissima*), 88, 157t, 587; as alien species, 130–31, 612t, 613f, 615; in bays and estuaries, 93f, 126t, 128f, 130t
Ammodytes hexapterus (Pacific sand lance), 17t, 75, 153t, 296f, 494–95, 522t
 amphibious fishes, 208, 209, 213
 Amphiprioninae, 503, 522t
Amphistichus spp. (surfperch): *A. argenteus* (barred surfperch), 18t, 150t, 151f, 154f, 155t, 160t; *A. koelzi* (calico surfperch), 19t, 97f, 150t, 151, 152f, 153t, 585t; *A. rhodoterus*, 17t. *See also* Embiotocidae
 anadromous fishes, 27, 88; alien, 615–16; in bays and estuaries, 124, 125, 128f; phyogeography and, 29–32. *See also* *Oncorhynchus* spp.
 Anaheim Bay, 135
Anarhichthys ocellatus (wolf-eel), 17t, 75, 75f, 91f, 255; in kelp bed/rocky reef, 236t, 240t; reproduction and, 485, 497, 498, 522t
Anchoa spp. (anchovies), 519t, 570, 573–74, 574, 579; commercial fishing of, 569, 576, 578–79; larval stage, 304f. *See also* *Engraulis mordax*
Anchoa compressa (deepbody anchovy), 18t, 95f, 160t, 173; in bays and estuaries, 126t, 134f, 136t, 138t, 145t; in surf zone, 150t, 151, 154, 159f; trophic relations and, 396f, 401
Anchoa delicatissima (slough anchovy), 19t, 396f, 401–2; in bays and estuaries, 93f, 126t, 134f, 138t, 145t
 anchovy and sardine regimes, 15, 281–82, 283–84f, 289; climate change and, 624, 625f, 633; competition and, 450; PDO and, 629–30, 629f, 630f, 631; trophic relations and, 403, 405
 Andersen's lanternfish (*Diaphus anderseni*), 351t
 Anderson, Todd W., 229, 421, 542; predation studies of, 436, 438–41, 442, 444
 Andrews, A. H., 373
 anemones (Actrinidae), 220, 560–61, 606
 angel shark (*Squatina californica*), 19t, 95f, 184, 233t, 390, 518t; commercial fishing and, 579, 591; movement of, 538t, 540; in surf zone, 153, 156f, 157t, 158t; and telemetry, 527

- anglerfish (ceratoids), 324, 358, 406, 487, 520t, 558f. *See also* dreamers (*Oneirodes*)
- Anguillidae (freshwater eels), 329, 348, 349t; *A. anguilla* (European eel), 307; *A. rostrata* (American eel), 612t, 618
- Anisotremus davidsonii* (sargo), 19t, 66, 67f, 74, 89, 94f, 136t, 534t; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 235t, 239f, 241f; phylogenetic relations and, 43, 44f, 45t, 46t
- Anisotremus interruptus* (burrito grunt), 235t
- Anomalopidae (flashlight fishes), 66, 370f, 520t, 558f. *See also* California flashlight fish
- Anomalops katoptron* (anomalopid), 558
- Anoplarchus insignis* (slender cockscomb), 17t, 21t, 235t
- Anoplarchus purpureus* (high cockscomb), 17t, 35t, 90f, 498, 522t; in rocky intertidal zone, 209t, 210, 213, 215, 216f, 220; trophic relations and, 399f, 403
- Anoplogaster cornuta* (fangtooth), 311f, 402f, 520t; as bathypelagic fish, 348, 352t, 358, 366t
- Anoplopoma fimbria* (sablefish), 18t, 101f, 106t, 306, 521t, 532t; in bathypelagic zone, 353t, 369, 372f; commercial fisheries and, 574, 575, 578, 580; on continental shelf, 172, 178f, 185f, 187; foraging guild, 192f, 197f; pollution and, 598
- Anoplopomatidae (sablefishes), 106t, 175, 180t
- Anopterus pharao* (daggertooth), 327f, 330, 519t
- Antemarius* spp., 20t, 24t, 520t
- Antimora microlepis* (Pacific flatnose), 18t, 21t, 353t, 372f
- Antimora rostrata*, 519t
- Apadocreedia vanderholsti*, 522t
- Apodichthys flavidus* (penpoint gunnel), 17t, 90f, 111f, 129f, 151, 153t, 485; in rocky intertidal zone, 209t, 215t, 216f, 221
- Apogonidae (cardinalfishes), 66, 234t, 506, 547, 548; *A. guadalupensis* (Guadalupe cardinalfish), 19t, 24t, 234t, 485; *A. lineatus* (mouth-brooding cardinalfish), 502, 521t; *A. pacificus* (pink cardinalfish), 7t, 20t, 25t, 234t
- Apristurus brunneus* (brown cat shark), 18t, 21t, 101f, 177t, 184, 185f, 197f
- Apristurus kampae* (longnose cat shark), 353t, 372f
- Argentina sialis* (Pacific argentine), 19t, 22t, 100f, 177t, 182f, 195f, 519t
- Argentinidae (argentines), 106t, 180t, 354
- Argyropelecus* spp. (hatchetfish), 306, 350t, 354, 356f, 367; *A. affinis* (slender hatchetfish), 350t, 354, 356f, 359, 366t; *A. hemigymmus* (spurred hatchetfish), 350t, 361; *A. lychnus* (tropical hatchetfish), 350t, 356f, 359, 366t; *A. sladeni* (lowcrest hatchetfish), 350t, 354, 356f, 366t
- Aristostomias scintillans* (shining loosejaw), 350t, 356, 357f
- Arius graeffei*, 519t
- Arius planiceps* (ariid catfish), 485
- Armor, C., 123
- Arnoglossus japonicus*, 308f
- arrow goby (*Clevelandia ios*), 11, 19t, 34t, 159, 161t, 523t, 533t, 600; in bays and estuaries, 93f, 126t, 128f, 129f, 132f, 134f, 136t, 138t, 143f, 145t; larval stages of, 294, 298, 303, 304f; symbiosis and, 559, 559f; trophic relations of, 396f, 401
- arrowtooth flounder (*Atheresthes stomias*), 17t, 179t, 185f, 197f
- Artedius* spp. (sculpins), 295, 296f; *A. corallinus* (coralline sculpin), 18t, 22t, 91f, 234t, 238f; *A. creaseri* (roughcheek sculpin), 75, 92f, 238f; *A. fenestralis*, 17t, 500t, 507; *A. meanyi*, 17t; *A. notospilotus* (bonehead sculpin), 18t, 153t
- Artedius harringtoni* (scalyhead sculpin), 17t, 91f, 213, 234t, 238f; reproduction and, 498, 500t, 502
- Artedius lateralis* (smoothhead sculpin), 18t, 90f, 500t; feeding of, 392, 394; in rocky intertidal zone, 209t, 215t, 217f, 220f
- Ascelichthys rhodorus* (rosytip sculpin), 17t, 91f, 209t, 499, 500t, 507
- Asemichthys taylori* (small-bodied sculpin), 486, 500t
- Assurger anzac*, 20t, 24t
- Asterotheca infraspinata*, 17t
- Asterotheca pentacantha*, 17t, 22t
- Atheresthes stomias* (arrowtooth flounder), 17t, 179t, 185f, 197f
- Atherinidae (smelt and grunions), 73, 304, 306, 321, 323, 329. *See also* *Leuresthes tenuis/sardina* (grunion)
- Atherinops affinis* (topsmelt), 19t, 73, 73f, 88–89, 94f, 475, 532t, 622; in bays and estuaries, 127t, 128f, 129f, 134f, 138t, 143f, 145t, 398; feeding and trophic relations of, 389, 394, 396f, 404; fisheries and, 585t; habitat of, 102, 103, 105t; in kelp bed/rocky reef, 233t, 239f; larval forms of, 303, 304f; in surf zone, 150t, 151–52f, 153t, 161t
- Atherinopsidae (silversides), 105, 106t, 149, 233t; and commercial fisheries, 574; genetic divergence in, 621–22, 622f; in harbors, 159f, 160t; larval stage of, 289, 304; in surf zone, 150, 154, 155, 156
- Atherinopsis californiensis* (jacksmelt), 18t, 73, 89, 94f, 233t, 303, 585t; in bays and estuaries, 127t, 128f, 129f, 134f, 138t, 145t; genetic divergence in, 622; in surf zone, 150t, 151f, 153t, 157t, 161t
- Atlantic menhaden (*Brevoortia tyrannus*), 469, 546, 600
- Atlantic-Pacific connection, 68–69
- Atlantic salmon (*Salmo salar*), 612t, 618, 619
- Atlantic wreckfish (*Polyprion americanus*), 497, 521t
- Atractoscion nobilis* (white sea bass), 19t, 96f, 111f, 336t, 337t, 432, 603; commercial fishing of, 574, 579, 588; in kelp bed/rock reef, 235t, 239f, 241f; larval stages of, 274, 275f; pollutants and, 598, 603; recreational fishing of, 580, 582, 584; reproduction and, 494; in surf zone, 151, 154f, 155t, 156f, 157t, 158t
- Attia, P., 375
- Aulostomus chinensis*, 520t
- aurora rockfish (*Sebastes aurora*), 18t, 22t, 101f, 278
- Autorhynchus flavidus* (tubesnout), 18t, 91f, 111f, 504, 520t; in kelp bed/rocky reef, 233t, 240f, 243f
- Auxis rochei*, 20t, 24t
- Auxis thazard*, 20t, 24t
- Avise, John C., 5
- Avocettina bowersii* (snipe eel), 349t
- Avocettina infans* (blackline snipe eel), 354, 355f
- Axoclinus carminalis*, 522t
- ayu (*Plecoglossus altivelis*), 612t, 618
- Azurina hirundo* (swallowtail damsel), 235t
- bacterial endosymbionts, 558
- Bagre panamensis*, 20t, 24t
- Bailey, T. G., 359, 360t, 364, 376
- Baja California, 27, 41, 43–46t; bays and estuaries in, 120, 121f, 136–37; kelp reef species, 92f, 239–40
- Bajacalifornia burraigei* (sharpchin slickhead), 349t
- bald sculpin (*Clinocottus recalvus*), 19t, 90f, 209t, 215t, 216f, 394, 403
- Balistes polylepis* (finescale triggerfish), 21t, 23t, 236t, 309f, 523t; on continental shelf, 179t, 194f; reproduction and, 485
- Balistidae (triggerfishes), 175, 179t, 180t, 236t
- balloonfish (*Diiodon holocanthus*), 7t, 21t, 25t, 523t
- balloon trawl, 575, 577
- Baltz, D. M., 618
- banded guitarfish (*Zapteryx exasperata*), 20t, 135t, 137f, 233t; on continental shelf, 177f, 183f, 194, 196f
- bank rockfish (*Sebastes rufus*), 18t, 23t, 100f, 102, 105t, 262f
- Barathronus pacificus*, 353t
- barcheek pipefish (*Syngnathus exilis*), 111f, 151f, 154f, 155t, 598
- barge fishing, 580, 582–83, 583f, 588
- Barham, E. G., 368
- barracuda (*Sphyræna argentea*), 19t, 61, 96f, 236t, 293, 330, 332f, 390, 523t; abundance of, 336, 337t; and commercial fishing, 574, 575f, 587; larval stages of, 295f, 297f, 300f, 302, 311; and recreational fishing, 581, 582, 583, 584, 585t; seasonal movement of, 541–42; in surf zone, 150t, 154, 155t, 158t
- in harbors, 157t, 158t, 159f, 160–61t; in kelp bed/rocky reef, 239f, 241f
- barracuda (*Sphyræna ensis*), 7t, 20t, 25t, 236t
- barracudinas (Paralepididae), 279f, 351t; *Lestidiops ringens*, 277, 280, 351t, 355f, 370f
- barred pipefish (*Syngnathus auliscus*), 20t, 88, 97f, 111f, 152; in bays and estuaries, 93f, 126t, 134f, 145t
- barred sand bass. *See* *Paralabrax nebulifer*
- barred surfperch (*Amphistichus argenteus*), 18t, 150t, 151f, 154f, 155t, 160t
- barreleye (*Macropinna microstoma*), 349t
- Barry, J. P., 133
- basketweave cusk-eel (*Ophidion scrippsae*), 96, 98f, 163, 181f, 182f; larval stages of, 277t, 280f, 293, 295f, 297f, 298; trophic relations and, 397f
- basking shark (*Cetorhinus maximus*), 19t, 326f, 336t, 390, 518t
- bass. *See* *Morone saxatilis*; *Paralabrax* spp.; Serranidae (seabasses)
- Bass, A. H., 491
- Bassozetes* spp. (cusk-eel), 360. *See also* cusk-eels (Ophidiidae)
- Bathophilus filifer* (threadfin dragonfish), 350t
- Bathophilus flemingi* (highfin dragonfish), 351t, 357f
- Bathylagopus* spp.: *B. infraspinus*, 21t; *B. nigripinnis*, 17t, 21t; *B. pentacanthus* (big-eye poacher), 178t, 197f
- Bathygobius* spp., 211, 493, 498
- Bathylagidae (deep-sea smelts), 349t, 354, 365, 370t; *B. bericoides*, 310f; *B. nigrigenys* (blackchin blacksmelt), 349t; *B. ochotensis* (popeye blacksmelt), 276t, 277–78f, 365, 519t
- Bathylagus pacificus* (Pacific blacksmelt), 310f, 349t, 354, 355f, 370f; swim bladder of, 365, 366t
- Bathylagus wesethi* (snubnose blacksmelt), 349t, 354, 355f, 360t, 370f; larval stages of, 276t, 277f, 279f, 280, 285–86, 289f, 368; swimbladder of, 365, 366t

- Bathylchmops exilis* (javelin spookfish), 349t
 Bathymasteridae (ronquils), 100f, 186, 238f;
Rathbunella spp., 18t, 72, 72f, 235t, 522t,
 532t
 bathypelagic zone (deep sea), 320, 321f,
 342–77; benthic and bathypelagic fishes,
 369–77; bioluminescence and, 342, 343,
 345t, 354, 356; chemistry of buoyancy in,
 365–68; ecological groupings in, 363–64;
 feeding in, 364–65; geographic variations,
 359; geologic features, 344, 346; grenadier
 spp., 353t, 369, 373–77, 390; growth and
 reproduction, 368–69; mesopelagic fishes
 and, 343, 345t, 346, 348, 406; midwater
 fishes, 348, 354–59; physical features of,
 344–48; trophic relations in, 402f, 406;
 vertical distribution in, 363; water types
 and currents, 346–48. *See also* bristle-
 mouths; lampfish; *specific species*
Bathyraxa spp. (deep-sea skates): *B. abyssicola*,
 19t, 22t, 353t, 372f; *B. interrupta*, 17t,
 22t; *B. spinosissima*, 19t, 22t; *B. trachura*
 (black skate), 18t, 353t
Bathysaurus mollis (highfin lizardfish), 353t,
 369, 372f, 374
 Batoidae (rays), 184, 539, 574. *See also* bat ray;
 Myliobatiformes; Rajidae; Torpedinidae
 Batrachoididae (toadfishes), 106t, 178t, 180t,
 210, 532t; reproduction and, 493, 503,
 504, 506. *See also* *Porichthys* spp.
 bat ray (*Myliobatis californicus*), 19t, 95f, 233t,
 337t, 429, 518t, 535t; in bays and estuar-
 ies, 127t, 128f, 132f, 134f, 137f; feeding
 of, 394; subsistence fishing and, 568; in
 surf zone, 150t, 153, 156f, 157t, 158t, 160t
 Baumgartner, T. R., 624
 Baxter, J. L., 588
 bay blenny (*Hypsoblennius gentilis*), 19t, 95f,
 111f, 236t, 455, 532t; in bays and estuar-
 ies, 126t, 134f, 136t, 137f, 145t
 bay goby (*Lepidogobius lepidus*), 18t, 98f, 155t,
 158, 161t, 453t, 534; in bays and estuar-
 ies, 126t, 128f, 132f; larval stages of, 292,
 294f, 298, 300, 303, 304f
 bay pipefish (*Syngnathus leptorhynchus*), 18t,
 95f, 111f, 429, 475; in bays and estuar-
 ies, 126t, 128f, 129f, 132f, 134f, 136t,
 137f, 145t; in surf zone, 151, 152f, 153t;
 trophic relations and, 396f, 402
 bays and estuaries (BE), 82t, 84f, 85t, 86f, 88,
 119–45; alien species in, 130–31, 618–19;
 analysis of, 104t; background to, 120–22;
 Baja California, 120, 121f, 136–37; central
 California, 121f, 131–33; dendrograms, 8,
 11f, 14f, 124f; ecological classification,
 123–24; future studies, 144–45; high pro-
 ductivity and biomass, 138–39; interan-
 nual variability, 140–42; introduction to,
 119–20; larval assemblages in, 303–5; lati-
 tudinal distribution, 122–23; northern
 California, 124–25, 128–31; nursery func-
 tion of, 142–43, 303–5, 468; salinity in,
 120, 122f, 123f; seasonality in, 139–40;
 southern California, 121f, 133–36;
 species, 93f, 95f; species diversity in,
 137–38; species in jeopardy, 125, 128–30;
 trophic relationships in, 396–98, 400–402
 Beamish, R. J., 306
 bearded eelpout (*Lycónema barbatum*), 18t, 22t,
 179t, 183f, 185f; foraging guild, 196f, 197f
 Beattie, A. J., 554
 Beck, M. W., 119, 143, 223
 Behrents, K. C., 214, 247, 435–36, 438, 441
Bellator gymnotetus (nakedbelly sea robin),
 178t, 183f, 196f
Bellator xenisma, 20t, 24t
 Bellwood, D. R., 57
 Belonidae (needlefishes), 73
 belt transect sampling, 229–30, 231, 232
 Bennett, W. A., 616
Benthalbella spp. (pearleyes): *B. dentata*
 (northern pearleye), 351t, 370f; *B. infans*,
 519t; *B. linguoides* (longfin pearleye), 351t
 benthic eggs, 77
 benthivores, 65, 395
 bathypelagic fishes, 345t
 bathypelagivores, 193, 403
 benttooth bristlemouth. *See* *Cyclothone acclin-*
idens
 Bernardi, Giacomo, 5
 Berry, F. H., 359, 361
 beryciforms, 57, 66
Beryx splendens, 520t
 bigeye poacher (*Bathylagus pentacanthus*),
 178t, 197f
 bigeye thresher shark, 89f, 326f, 337t
 bigeye trevally (*Caranx sexfasciatus*), 7t, 21t
 bigeye tuna, 89f, 326f, 337t
 bigfin eelpout (*Lycodes cortezianus*), 18t, 22t,
 179t, 184, 185f, 197f
 bigmouth sole (*Hippoglossina stomata*), 19t,
 23t, 98f, 236t, 398f, 535t, 606; on conti-
 nental shelf, 179t, 182f, 183f; foraging
 guild, 192f, 195f, 196f; larval stages of,
 293, 295, 302
 bigscale goatfish (*Pseudupeneus grandisquamis*),
 7t
 bigscale pomfret (*Taractes longipinnis*), 327f, 330
 bigscales (*Melamphaes* spp.), 279f, 280, 352t,
 359; *M. acanthomus* (slender bigscale),
 352t; *M. lugubris* (highsnout bigscale),
 277, 308f, 352t, 358f; *M. parvus* (little
 bigscale), 277, 352t; *Scopelogadus mizolep-*
sis bispinosus, 352t, 361, 365, 366t
 big skate (*Raja binoculata*), 18t, 177t, 181f,
 182f, 183f, 518t; foraging guild, 194f,
 195f, 196f
 billfishes, 335t, 429, 530. *See also* *Makaira*;
 swordfish (*Xiphias gladius*)
 biogeography, 3–15, 190, 622; climatic per-
 spectives and, 4–5, 14–15; cluster analy-
 sis of bay and non-bay species, 10–11;
 dendrograms, 7–8, 8, 11f; distribution
 patterns and, 5–6; end points of species
 ranges, 8–9, 12f, 13f; latitude and ordina-
 tion, 11–14, 243; phylogeography and,
 26; regions and provinces, 3–4; species
 richness and, 5, 6–8
 biological clocks, 548–49
 bioluminescence, 342, 343, 345t, 354, 356;
 reproduction and, 491; symbiotic rela-
 tions and, 557–58
 birds, and pollutants, 602
 black-and-yellow rockfish (*Sebastes chrysome-*
las), 19t, 67f, 91f, 162t, 215t, 219t; in
 kelp bed/rocky reef, 234t, 237f, 240f;
 movement of, 531, 536t; recruitment of,
 415, 420; territoriality of, 453t, 455–56f,
 457f; trophic relations and, 399t, 403
 blackbelly dragonfish (*Stomias atriventer*),
 351t, 356, 357f, 366t, 370f; larval stages
 of, 277, 279f, 280; trophic relations of,
 402f, 406
 blackbelly eelpout (*Lycodes pacificus*), 18t, 98f,
 179t, 183f, 192f, 196f, 538t
 black bullhead, 129f
 black croaker (*Cheilotrema saturnum*), 95f,
 145t, 153, 155t, 156f, 157t
 black cusk-eel (*Cherublemma emmelas*), 177t,
 197t
 black eelpout (*Lycodes diapterus*), 17t, 22t,
 354t
 blackeye goby (*Rhinogobiops nicholsii*), 19t,
 60, 61f, 76, 162t, 420, 506; habitat of,
 102, 105t; in kelp bed/rocky reef, 236t,
 238f, 239f, 240f, 247; in mid-depth
 rocky reef, 99f, 255, 260f, 263t; move-
 ment and activity patterns of, 523t, 531,
 534t, 545; predation study of, 431–32f,
 434–37f, 438, 439–42, 460–61; reproduc-
 tion and, 494, 497; territoriality of, 455
 blackfinned snailfish (*Careproctus cypselurus*),
 353t
 blackgill rockfish (*Sebastes* sp.), 101f
 black hagfish (*Eptatretus deani*), 18t, 101f,
 353t, 372f
 blackline snipe eel (*Avocettina infans*), 354, 355f
 blacklip dragonet (*Synchiropus atrilabiatus*), 7t,
 20t, 25t, 179t, 183f, 196f
 black marlin (*Makaira indica*), 20t, 25t, 323,
 333
 blackmouth eelpout (*Lycodapus fierasfer*), 17t,
 22t, 354t
 black perch. *See* *Embiotoca jacksonii*
 black prickleback (*Xiphister atropurpureus*),
 18t, 90f, 236t, 498, 522t; in rocky inter-
 tidal zone, 209t, 210, 211f, 215t, 216f;
 trophic relations and, 394, 399f, 403
 blackrag (*Psenes pellucidus*), 328f, 330
 black rockfish (*Sebastes melanops*), 17t, 70,
 70f, 91f, 111f, 255, 475, 585t; in kelp
 bed/rocky reef, 237f, 240f; rocky inter-
 tidal zone, 215t, 216f, 217t, 219t; in surf
 zone, 151, 152f, 155t
 black skate (*Bathyraxa trachura*), 18t, 353t
 blacksmelt. *See* *Bathylagus* spp.
 blacksmith. *See* *Chromis punctipinnis*
 blackspot wrasse (*Decodon melasma*), 7t, 20t,
 24t
 black swallower (*Chiasmodon niger*), 344,
 522t
 blacktail snailfish (*Careproctus melanurus*),
 18t, 22t, 353t, 372f; on continental shelf,
 178t, 184, 185f, 197f
 blacktip poacher (*Xeneretmus latifons*), 18t,
 23t, 98f, 178t, 183f, 192f, 196f, 532t
 blackwing flyingfish, 326f
 bleeding wrasse (*Polylepium cruentum*), 179t,
 183f, 196f
 Blenniidae (combt tooth blennies), 236t, 289,
 532t; reproduction and, 485, 491, 493,
 502, 504. *See also* *Hypsoblennius* spp.
Blepsias cirrhosus (silverspotted sculpin), 17t,
 151, 152f, 153t; reproduction in, 486,
 499–500t
 blind goby (*Typhlogobius californiensis*), 19t,
 24t, 211, 497, 532t; symbiotic relations
 of, 559–60, 560f
 blob sculpin (*Psychrolutes phrictus*), 353t,
 372f, 521t
 bluebanded goby (*Lythrypnus dalli*), 19t, 23t,
 61f, 92f, 236t; activity patterns of, 534t,
 545; in kelp bed/rocky reef, 236t, 238f,
 247; phylogenetic characteristics of, 43,
 44f, 45t, 46t, 49, 623; predation and,
 431–32f, 434–37f, 438, 439–42, 460–61;
 recruitment of, 419–20, 422; reproduc-
 tion and, 487; symbiotic relations of,
 555, 556f, 557t
 bluebanded ronquil, 100f, 238f
 blue chromis. *See* *Chromis punctipinnis*
 bluefin tuna (*Thunnus thynnus*), 89f, 401f,
 542; commercial fishing of, 573, 578;
 feeding behavior of, 389, 390; in pelagic
 zone, 326f, 330, 336t, 337t
 bluefish (*Pomatomus saltatrix*), 149, 521t
 bluehead wrasse (*Thalassoma bifasciatum*),
 487, 488

- blue lanternfish (*Tarletonbeania crenularis*), 306, 351t, 357f, 359, 368; swimbladder of, 366t, 367; trophic relations of, 402f
- blue marlin (*Makaira nigricans*), 332
- blue rockfish (*Sebastes mystinus*), 18t, 59f, 91f, 215, 416; in deep rock habitat, 255, 260f; fishing and, 584, 585t, 590; in kelp bed/rocky reef, 234t, 237f, 240f, 242, 243f; reproduction and, 494; seasonal movement of, 537t, 542; trophic relations and, 394, 404
- blue shark (*Prionace glauca*), 20t, 89f, 326f, 330, 337t, 389, 518t; movement of, 524, 532t, 540, 542, 543; telemetry and, 527–28; trophic relations and, 401f
- bocaccio (*Sebastes paucispinnis*), 18t, 59, 234t, 334f, 484f, 632; commercial fishing of, 575, 589, 590; in deep rock habitat, 256, 260f, 261f, 263t; epibenthic, 539; habitat for, 100f, 103; larval stages of, 276t, 309f; movement of, 528, 537t, 541; recreational fishing and, 584, 585t; young-of-year (YOY), 111f, 261, 262f
- Bodianus diplotaenia* (Mexican hogfish), 235t
- Bodkin, J. L., 241, 244, 246
- Boehlert, George W., 270, 305
- Bone, Q., 368
- bonefish (*Albula vulpes*), 20, 135t, 145t, 518t, 532t
- bonehead sculpin (*Artedius notospilotus*), 18t, 153t
- bonito (*Sarda chiliensis*), 20t, 87, 89f, 236t, 585t; in coastal pelagic zone, 154, 157t, 158t, 159f, 160t; commercial fisheries and, 574, 575; feeding of, 406; in pelagic zone, 330, 332f, 336t, 337t; pollution and, 600, 601; recreational fishing and, 581, 583, 584; reproduction and, 494; seasonal movement of, 541–42; trophic relations and, 401f
- Booth, F. E., 571
- boreal region, 330
- Borostomias panamensis* (Panama snaggletooth), 351t, 357f, 366t
- Bothidae (lefteye flounders), 176, 180t, 329. *See also* Paralichthyidae
- Bothrionus swanii* (rockhead), 17t, 70
- Bothrocara brunneum* (two-line eelpout), 18t, 22t, 354t, 372f
- Bothrocara molle*, 18t, 22t
- bottletail (*Danaphos oculatus*), 350t, 359, 366t, 370f
- Bowen, B. W., 624
- Boyle, K. S., 403
- Brachyistius frenatus* (kelp perch), 19t, 71f, 75, 111f, 162t, 422, 477, 494; in kelp bed/rocky reef, 92f, 235, 239f, 240f; movement and activity patterns of, 533t, 544, 556f, 557t; predation and, 431f, 438–39, 440f, 442f; trophic relations and, 395, 400f, 404
- Bradbury, J. W., 488
- Brama japonica*, 20t, 22t, 521t
- Brantley, R. K., 491
- Bray, Richard N., 433, 524
- breeding, 531. *See also* reproduction
- Bregmaceros atlanticus*, 519t
- Breitberg, D. L., 455
- Brevoortia tyrannus* (Atlantic menhaden), 469, 546, 600
- Brewer, G. D., 305, 362
- Briggs, J. C., 174
- bristlemouths (Gonostomatidae), 290, 350t, 354, 359, 365, 370f. *See also* Cyclothone spp.
- brittlestar (*Ophiophthalmus normani*), 371, 540
- broadfin lanpfish (*Nannobranchium ritleri*), 351t, 356, 357f, 359, 368, 370f; larval stages of, 276t, 277f, 279, 280, 305; swimbladder of, 366t, 367; trophic relations of, 402f
- bronzedspotted rockfish (*Sebastes gilli*), 18t, 24t, 258t
- bronzestripe grunt (*Orthopristis reddingi*), 178t, 181f, 182f, 194f, 195f
- Brooks, A. J., 249–50
- broomtail grouper (*Mycteroperca xenarcha*), 20t, 23t, 58, 234t
- Brosomphycis marginata* (red brotula), 18t, 186, 233t, 238f, 519t
- brotulas. *See* Bythitidae (brotulas)
- Brown, D. W., 359, 360t
- Brown, James H., 5, 621
- brown algae (*Pterygophora californica*), 228, 246
- brown cat shark (*Apristurus brunneus*), 18t, 21t, 101f, 177t, 184, 185f, 197f
- brown Irish lord (*Hemilepidotus spinosus*), 17t, 91f, 234t, 238f, 296f
- brown rockfish (*Sebastes auriculatus*), 19t, 22t, 111f, 255, 270; in kelp bed/rocky reef, 92f, 234t, 240f; movement of, 536t, 542; in surf zone, 152, 155t
- brown smoothhound (*Mustelus henlei*), 19t, 95f, 127t, 128f, 233t; in surf zone, 153, 156f, 157t, 158t
- brownsnout spookfish (*Dolichopteryx longipes*), 349t
- brown trout (*Salmo trutta*), 612t, 616
- buffalo sculpin (*Enophrys bison*), 17t, 234t, 485, 486, 500
- bulbous dreamer (*Oneirodes eschrichtii*), 352t
- bullet mackerel, 337t
- bull sculpin (*Enophrys taurus*), 17t
- bullseye puffer (*Spherooides annulatus*), 20t, 25t, 95f, 96, 137f, 236t, 237f, 241f
- buoyancy and swim bladders, 365–68
- Bureau of Commercial Fisheries (BCF), 269
- Burge, R. T., 245
- Busby, P. J., 31
- butterflyfishes (Chaetodontidae), 63–64, 235t, 394, 506, 522t; *Chaetodon humeralis*, 20t, 24t, 235t; *Prognathodes falcifer*, 20t, 24t, 220, 235t
- butterfly ray (*Gymnura marmorata*), 20t, 135t, 518t
- butter sole (*Isopsetta isolepis*), 17t, 174, 179t, 181f, 194f, 296f
- Bythitidae (brotulas), 185f, 186, 484, 504; red brotula (*Brosomphycis marginata*), 18t, 186, 233t, 238f, 519t
- cabezon. *See* *Scorpaenichthys marmoratus*
- Cabrillo Beach, 161f, 162, 164f
- Caelorinchus scaphopsis* (shoulderspot), 369
- Caillet, G. M., 154, 364, 371
- Calamus brachysomus* (Pacific porgy), 20, 24t, 137f, 235t, 521t
- CalCOFI. *See* California Cooperative Oceanic Fisheries Investigations (CalCOFI)
- calico rockfish (*Sebastes dallii*), 19t, 98f, 187, 234t, 398f, 416; movement of, 536t, 542
- calico surfperch (*Amphistichus koelzi*), 19t, 97f, 585t; in surf zone, 150t, 151, 152f, 153t
- California barracuda. *See* barracuda (*Sphyrna argentea*)
- California butterfly ray (*Gymnura marmorata*), 20t, 135t, 518t
- California Channel Islands, 33, 36, 46–49, 228, 240, 603, 624
- California clingfish (*Gobiesox rhessodon*), 19t, 90f, 212–13, 215t, 216f, 238f, 403
- California Cooperative Oceanic Fisheries Investigations (CalCOFI), 269–70, 271, 275–302, 313, 361, 362, 368–69, 628. *See also* ichthyoplankton assemblages
- California corbina. *See* corbina
- California Counter Current, 47, 347
- California Current, 47, 320, 329, 347, 368, 629; pelagic fishes and, 333–36; phylogeography and, 38, 40, 41
- California Current region, ichthyoplankton studies in, 269, 270, 280, 283, 287–88, 306. *See also* ichthyoplankton assemblages
- California Department of Fish and Game (CDFG), 168, 169, 269, 587, 588, 617; and recreational fishing, 580, 583, 590, 591; and sardine industry, 571, 572, 573
- California Fish and Game* (journal), 587
- California flashlight fish (*Protomyctophum crockeri*), 351t, 357f, 359, 370f; larval stages of, 276t, 277f, 279f, 280, 306, 368; swimbladder of, 366t, 367
- California grenadier. *See* *Nezumia stelgidolepis*
- California grunion (*Leuresthes tenuis*), 95f, 145t, 161t; in surf zone, 150t, 151f
- California halfbeak (*Hyporhamphus rosea*), 19t, 73, 88, 93f, 135t, 136t, 328
- California halibut. *See* *Paralichthys californicus*
- California headlightfish (*Diaphus theta*), 305, 351t, 356, 357f, 359, 360t, 363, 366t; trophic relations of, 402f
- California killifish (*Fundulus parvipinnis*), 10–11, 19t, 34t, 41, 42f, 88, 533t; abundance of, 142, 143f, 145t; in bays and estuaries, 93f, 120, 126t, 129f, 132f, 134f, 137f, 138; pollution and, 601; trophic relations and, 396f, 401
- California lanternfish (*Symbolophorus californiensis*), 351t, 357f, 358, 359, 402f; larval stages of, 276t, 277f, 279f, 280, 282f; swimbladder of, 366t, 367
- California lizardfish (*Synodus lucioceps*), 18t, 98f, 155t, 538t; on continental shelf, 173, 177, 181f, 182f, 183f; foraging guilds, 192f, 194f, 195f, 196f; larval stages of, 277, 293, 295f, 301, 302f; trophic relations and, 395, 397f, 398f
- California moray (*Gymnothorax mordax*), 19t, 24t, 95f, 545, 555, 558f; activity patterns of, 545, 549; in harbors, 157t, 158t; in kelp bed/rocky reef, 233t, 238f, 239f, 241f, 243f
- California needlefish (*Strongylura exilis*), 73, 135t
- California rattail, 101f, 372f
- California roach, 128f, 129f
- California sardine. *See* *Sardinops sagax*
- California sheephead. *See* *Semicossyphus pulcher*
- California skate (*Raja inornata*), 18t, 98f, 177t, 181f, 182f, 183f; foraging guild, 194f, 195f, 196f
- California slickhead (*Alepocephalus tenebrosus*), 20t, 21t, 101f, 353t
- California smoothtongue. *See* *Leuroglossus stilbius*
- California subregion, 27
- California tonguefish. *See* *Symphurus atriacauda*
- California Transition Zone, 5
- Callionymidae (dragonets), 176, 180t. *See also* blacklip dragonet
- camouflage. *See* cryptic species
- canary rockfish (*Sebastes pinniger*), 18t, 99f, 255, 260f, 263t; fisheries and, 585t, 589, 591
- canneries, 569, 570–74, 579
- cannibalism, 436, 500–502
- Cannon, Ray, 590
- Canthigaster* spp. (pufferfish), 503, 523t

- canyon species, 100f
Caracanthus spp., 520t
 Carangidae (jacks), 9f, 10f, 62, 77, 106, 150, 329, 393; *C. caballus*, 20t, 23t; *C. caninus* (Pacific crevalle jack), 7t, 20t, 25t; *C. sexfasciatus* (bigeye trevally), 7t, 21t; *C. vinctus* (cocinero), 7t, 20t
Carapus spp., 308f, 519t
 Carcharhinidae (sharks), 9f, 10f; *C. brachyurus*, 20t; *C. leucas*, 20t, 24t; *C. longimanus* (whitetip shark), 19t, 25t, 328f, 331; *C. obscurus*, 19t, 24t; *C. remotus*, 24t. *See also* Scyliorhinidae (cat sharks); Squalidae (dogfish sharks); Triakis (hound sharks)
Carcharodon carcharias (white shark), 20t, 22t, 89f, 337t, 390; movement of, 530, 534t, 542; pollution and, 601
 carcinogens. *See* DDT and PCB contamination
 cardinalfishes (Apogonidae), 66, 234t, 506, 547, 548. *See also* Apogon spp.
Careproctus spp. (snailfishes), 353t, 486, 521t. *See also* Liparis spp.
Careproctus melanurus (blacktail snailfish), 18, 22t, 353t, 372f; on continental shelf, 178t, 184, 185f, 197f
 Carey, F. G., 543
Caristius macropus, 521t
 Carlisle, J. G., Jr., 150
 Carlton, J. T., 612
 carnivores, 394. *See also* piscivorous fish
 Carpelan, L. H., 473
 Carr, M. H., 248, 419, 436, 441, 445
 Carroll, R. L., 57
 Cartamil, D. P., 540
 Case, J. F., 358
 Cass-Calay, S. L., 305
 catadromous fishes, 124, 126t, 134f
Cataetx rubrirostris (rubynose brotula), 186
 catch data, 525, 585t
 catch per unit effort (CRUE), 229, 371
 cat sharks. *See* *Apristurus* spp.; Scyliorhinidae (cat sharks)
Caulolatilus princeps (ocean whitefish), 20t, 22t, 35t, 87, 311f, 521t; in kelp bed/rocky reef, 92f, 234t, 237f, 239f; and recreational fishing, 586
Caulophryne spp., 309f, 520t
Cebidichthys violaceus (monkeyface pricklyback), 18t, 75, 90f, 235t, 558; feeding and trophic relations, 389, 394, 399f, 403; movement and activity of, 527, 530, 538t, 543; in rocky intertidal zone, 207, 209t, 211, 212, 215t, 216f, 219, 220f
 Cech, J. J., 123
 Cenozoic influences, 58, 68–70, 621
 Centrarchidae (sunfishes), 393. *See also* *Mola mola* (ocean sunfish)
 Centro Interdisciplinario de Ciencias Marinas (CICIMAR), 270
Centropomus undecimalis, 521t
Centropyge spp., 522t
Centrosyllium nigrum (combtooth dogfish), 353t
Centrostephanus coronatus (sea urchin), 441. *See also* sea urchins
Cephalopholis boenak, 497
Cephaloscyllium ventriosum (swell shark), 21t, 23t, 95f, 153, 156f, 157t, 158t; activity patterns of, 536t, 544, 549; in kelp bed/rocky reef, 233t, 239f
Cepola rubescens, 522t
 ceratoids (angler fish), 324, 358, 406, 487, 520t, 558f
Ceratoscopelus townsendi (dogtooth lampfish), 351t, 358, 366t, 369; larval stages of, 276t, 277f, 279, 280, 305
Ceratoscopelus warmingii (lampfish), 280
Cetengraulis mysticetus, 20t, 24t
Cetomimus sp., 520t
Cetorhinus maximus (basking shark), 19t, 326f, 330, 336t, 518t
 Chaenopsidae (pikeblennies), 236t, 289; *Chaenopsis alepidota* (orangethroat pikeblenny), 19t, 24t, 43, 44f, 45t, 46t, 137f. *See also* *Neoclinus* spp.
Chaetodipterus zonatus, 20t, 25t, 523t
 Chaetodontidae (butterflyfishes), 63–64, 235t, 394, 506, 522t; *C. humeralis* (three-banded butterflyfish), 20t, 24t, 235t; *Prognathodes falcifer*, 20t, 24t, 220, 235t
 chameleon goby (*Tridentiger trigonocephalus*), 612–13, 612t
 chameleon rockfish (*Sebastes phillipsi*), 17t, 23t, 258t
 chameleon wrasse (*Halichoeres dispilus*), 92f, 235t, 237f, 241f
 Channel Islands. *See* California Channel Islands
 Channel Islands clingfish (*Rimicola cabrilla*), 7t, 17t, 24t
Chanos chanos (milk fish), 519t, 612t, 618
Chauliodon macoumi (Pacific viperfish), 351t, 356, 357f, 365, 366t, 370f, 519t; trophic relations and, 395, 402f, 406
 Chavez, F. P., 5, 15, 629, 631
 cheekspot goby (*Ilypnus gilberti*), 95f, 154f, 155t, 159, 161t, 534; in bays and estuaries, 126t, 134f, 136t, 137f, 145t, 303; trophic relations and, 396f, 401
Cheilodactylus spectabilis, 522t
Cheilopogon spp. (flyingfish), 323, 328, 331, 405; *C. heterurus*, 20t, 25t, 520t; *C. pinna-tibarbatus* (California flyingfish), 19t, 22t, 405
Cheilotrema saturnum (black croaker), 19t, 95f, 145t, 153, 535t; in kelp bed/rocky reef, 235t, 239f
Cherublemma emmelas (black cusk-eel), 177t, 197f
 Chess, J. R., 75, 188, 546
 Chess, Tony, 55
Chiasmodon niger (black swallower), 344, 522t
Chilara taylori (spotted cusk-eel), 19t, 98f, 163, 176, 177t, 398f; on continental shelf, 182f, 183f, 185f; foraging guild, 192f, 194f, 195f, 196f, 197f; movement patterns of, 535t, 546
 Childress, J. J., 406
 chilipepper (*Sebastes goodei*), 19t, 100f, 172, 262f; fisheries and, 575, 585t, 590
Chilconger dentatus (thicklip conger), 177t, 183f, 195f, 196f
Chilomycterus reticulatus, 20t, 24t
 Chimaeridae (chimaeras), 106t; on continental shelf, 175, 177t, 180t
 china rockfish (*Sebastes nebulosus*), 17t, 76, 76f, 234t, 240t
 chinook salmon (*Oncorhynchus tshawytscha*), 17t, 153t, 233t, 330, 331f, 630; in bays and estuaries, 93f, 126t, 128f, 130t, 131; fisheries and, 569, 579, 585t, 589, 628; phylogeography and, 30t, 31–32, 32t, 39
Chirolophus decoratus (decorated warbonnet), 7t, 17t, 236t
Chirolophus nugator (mosshead warbonnet), 17t, 236t
Chitonotus pugetensis (roughback sculpin), 19t, 22t, 98f, 398f, 532t; on continental shelf, 176, 178t, 182f, 189; foraging guild, 192f, 195f
Chlamydoselachus anguineus, 17t, 21t, 518t
Chloroscombrus orqueta, 20t, 24t
 Choat, J. H., 57
 Chotkowski, M. A., 221
Chromis alta (silverstripe chromis), 235t
Chromis atrilobata (scissortail chromis), 92, 235t, 241f
Chromis punctipinnis (blacksmith), 19t, 35t, 74, 87, 92f, 157t, 416; cleaner behavior of, 556f, 557f; feeding of, 63f, 390; in harbors, 157t, 159, 162t; in kelp bed/rocky reef, 235t, 237f, 239f, 241f, 242, 243f; larval stages of, 293, 295f, 301, 302; movement and activity patterns of, 535t, 542, 544, 546, 548; trophic relations and, 400f
 chub mackerel. *See* *Scomber japonicus*
 chum salmon (*Oncorhynchus keta*), 17t, 21t, 130t, 233t
 Cichlidae (cichlids), 72, 497. *See also* *Tilapia* spp. (cichlids)
 cigarfish (*Cubiceps* spp.), 333, 523t, 559
 cigar shark (*Isistius brasiliensis*), 561, 561f
 circadian rhythms, 548–49
Cirrhitichthys aureus, 522t
Cirriformia luxuriosa (polychaete worm), 220
Citharichthys spp. (sanddabs), 193, 294, 305, 606–7; *C. fragilis* (gulf sanddab), 7t, 20t, 277f, 278f, 280; *C. gorda* (mimic sanddab), 179t, 183f, 193, 196f; DDT concentrations in, 601, 602f, 603; larval stages of, 276t, 280, 291, 297
Citharichthys sordidus (Pacific sanddab), 98f, 157t, 299, 403, 535t; on continental shelf, 172, 173, 179t, 182f, 258; diel movement of, 543; foraging guild, 192f, 195f; larval stages of, 19t, 98f, 157t, 276t; pollution and, 600, 601, 602f, 605, 606; recreational fishing and, 583, 585t; trophic relations and, 398f, 403
Citharichthys stigmaeus (speckled sanddab), 19t, 60, 61f, 236t, 258, 631; in bays and estuaries, 127t, 128f, 132f; on continental shelf, 96, 97f, 103, 176, 179t, 181f; foraging guilds, 192f, 194f; in harbors, 158, 161t; larval stages of, 276t, 277–78f, 280, 298, 299; movement of, 535t, 543; pollution and, 601, 602f, 606; in surf zone, 152f, 153t, 155t; trophic relations and, 397f, 403
Citharichthys xanthostigma (longfin sanddab), 19t, 23t, 98f, 398f, 535t; on continental shelf, 179t, 181f, 182f, 183f; foraging guild, 193, 194f, 195f, 196f; larval stages of, 276, 277–78f, 280; pollutant concentrations in, 600, 601, 602f
 Clark, Frances, 572, 588
 Clarke, G. L., 365
 Clarke, T. A., 227, 453
 cleaning symbiosis, 555–57, 556f, 557t
Clevelandia ios (arrow goby), 11, 19t, 34t, 159, 161t, 523t, 533t, 600; in bays and estuaries, 93f, 126t, 128f, 129f, 132f, 134f, 136t, 138t, 143f, 145t; larval stages of, 294, 298, 303, 304f; symbiotic relations of, 559, 559f; trophic relations of, 396f, 401
 climate change, 4–5, 14–15, 36–38, 47, 621–33; in Baja California, 41; eutrophication and, 145; global warming and, 632–33; over-fishing and, 625–26; rocky intertidal zone, 205, 221–22, 630; trophic interactions and, 631–32. *See also* ENSO; PDO
 clingfish (*Rimicola* spp.): *R. cabrilla* (Channel Islands clingfish), 7t, 17t, 24t; *R. dimorpha*, 18t, 24t; *R. eigenmanni* (slender clingfish), 19t, 25t, 92f, 111f, 236t, 238f; *R. muscarum* (kelp clingfish), 18t, 111f, 155t, 236t, 238f, 405. *See also* *Gobiesox* spp.

- Clinidae (kelpfish), 9f, 10f, 71–72, 105, 106t, 217f, 236t, 532t; reproduction and, 491, 495
- Clinocottus* spp. (sculpins), 128f, 500;
C. acuticeps (sharpnose sculpin), 17t, 209t, 498f, 500t, 521t; *C. embryum*, 18t, 420; *C. recalvus* (bald sculpin), 19t, 209t, 215t, 394, 500t
- Clinocottus analis* (wooly sculpin), 35t, 90f, 399f, 403, 532t, 628; hypoxia and, 468; parental care in, 500t, 507; in rocky intertidal zone, 209t, 211, 215t, 217f, 220f
- Clinocottus globiceps* (mosshead sculpin), 17t, 21t, 209t, 215t, 217f, 420, 500t; feeding of, 390, 392
- Clupea pallasii* (Pacific herring), 72, 152f, 153t, 214, 304f, 331f, 429; in bays and estuaries, 93f, 127t, 128f, 129f, 132f, 138t; commercial fishing of, 574, 576; on continental shelf, 172, 195f; feeding of, 390, 393, 394; and reproduction, 494
- Clupeidae (herrings), 72, 106t, 149, 304, 570, 574; *C. harengus*, 17t, 303, 325; commercial fishing of, 578; and reproduction, 494, 496
- coastal pelagic (CP) zone, 82t, 85t, 86f, 89f, 91, 96f, 106, 109, 154–56; analysis of, 104t
- coastal species, 8, 17–21t; dendrogram for, 11, 11f; phylogeographic characteristics of, 34–35t
- cocinero (*Caranx vinctus*), 7t, 20t
- cockcombs. *See* *Anoplarchus* spp.
- cod. *See* Gadidae (cods)
- codlings (Moridae), 175, 177t, 180t
- Cohen, A. N., 612
- coho salmon (*Oncorhynchus kisutch*), 18t, 30t, 31, 32, 88, 330, 331f; in bays and estuaries, 126t, 128f, 130t
- Collisella limatula* (limpet), 220
- Cololabis saira* (Pacific saury), 19t, 330, 405, 520t, 624
- coloration: anti-predatory, 428–29; courtship and, 488, 489, 490, 493, 494. *See also* crypsis
- Colpichthys regis* (false grunion), 622
- Columbia River, 31
- Colwell, R. K., 451
- combfishes. *See* *Zaniolepis* spp.
- combtooth blennies (Blenniidae), 236t, 289, 532t; reproduction and, 485, 491, 493, 502, 504
- combtooth dogfish (*Centroscyllium nigrum*), 353t
- commensalism, 554, 558–61
- commercial fisheries, 337t, 525, 567, 568–80, 587; (1850–1899), 568–70; (1900–1950), 570–77; (1951–2001), 577–80; balloon trawls and, 575, 577; canneries and, 570–74, 579; competition with recreational fishing, 589–92; drying/salting, 569–70; fishing techniques, 575–76; freezing and, 569, 575, 577, 579; globalization of, 578–79; landings made (1981–1999), 580t; market fisheries, 574–77; nets, 571, 575, 576, 578f, 579–80, 589, 590–91; pelagic wetfish and, 570, 573–74; purse seines, 571, 572, 573, 575, 577, 588; racism and, 586–87; reduction in, 570–74, 579; regulation of, 572, 573, 580, 586–89; sardines, 569, 570–73, 578–79, 588, 590; technology revolution in, 577–78; traditional, 579–80; trawling and, 570, 575, 577; tuna, 333f, 337t, 570–77, 579
- Commercial Passenger Fishing Vehicles (CPFVs), 580–84, 583f, 584f, 588
- competition, 415, 449–65; in context, 461–62; definition and concept of, 449–51; density-dependent, 449, 453; direct observation of, 450, 454; disturbance and, 472, 477; exploitation, 449; interference, 429, 449; interspecific, 443, 450, 451–61; niche overlap and, 451, 452f, 456, 457, 458f; predation and, 438, 443; resource partitioning and, 450, 454; territoriality and, 450, 451, 453t, 454, 455–56
- Congleton, J. I., 468
- Congridae (conger eels), 175, 177t, 180t, 183f, 195f, 196f
- Connell, J. H., 311, 434, 442, 466
- conservation, 569, 572, 587, 589; recruitment and, 422–23
- continental shelf and slope, 96, 167–99; assemblages, 172–73; deep shelf assemblages, 258t; diel behavior, 188; distribution of families in, 175–76; ecological segregation, 190–95; feeding and foraging, 188–92, 192f, 194–97f; foraging guilds, 190–92; interactions with other organisms, 195–95; life zones, 174–75; mobility and migration and, 187–88; morphological attributes, 184, 186; natural history traits, 185; other regions compared, 196–98; overview of scientific studies, 168–71; physical and biological conditions, 167–68; physical changes with depth, 168; population characteristics, 171–72; sampling methods, 170–71; soft-bottom habitat, 167–68; taxonomic composition, in surveys, 172; topography, 167; trawl survey assessments, 171–73; zoogeographic provinces, 173–74. *See also* inner shelf (IS); outer shelf (OS)
- Cook, A. E., 219–20, 223, 407
- copper rockfish (*Sebastes caurinus*), 17t, 111f, 255, 260f, 270, 585t; on continental shelf, 178t, 182f, 195f; in kelp bed/rocky reef, 92f, 234t, 240t; recruitment of, 415; seasonal movement of, 536t, 539, 542; in surf zone, 152, 155t
- coralline scuplin (*Artedius corallinus*), 18t, 22t, 91f, 234t, 238f
- coral reef communities, 57, 58, 63, 69, 71, 420, 437
- corbina (*Menticirrhus undulatus*), 19t, 95f, 136t, 149, 293, 536t; commercial fishing and, 587; in harbors, 157t, 158t, 160–61t; larval stages of, 293, 295f; pollutants and, 600; in surf zone, 150t, 151
- Cordilleran region, 27, 36
- Cordone, A. J., 130
- Coris julis*, 549
- cornetfish (*Fistularia corneta*), 7t, 14, 20t, 25t
- Cortez angelfish (*Pomacanthus zonipectus*), 235t
- Cortez gregory, 92f, 241f
- Cortez Province, 3, 4f
- Cortez stingray, 137f
- Coryphaena* spp., 519t; *C. hippurus* (dolphinfish), 20t, 22t, 324, 332, 521t
- Coryphaenoides* spp. (grenadiers), 353t, 369, 373–77, 530; *C. acrolepis* (Pacific grenadier), 18t, 22t, 353t, 369, 372f, 373, 374; *C. armatus* (abyssal grenadier), 353t, 369, 372f, 374–76; *C. filifer* (threadfin grenadier), 353t, 369; *C. leptolepis* (ghostly grenadier), 353t, 369, 374, 376; *C. pectoralis* (giant grenadier), 101f, 353t, 369, 372f, 373–74; *C. yaquinae* (rough abyssal grenadier), 353t, 369, 374–77, 540
- Coryphopterus* (goby), 76, 494
- Cosmocampus arctus* (snubnose pipefish), 19t, 92f, 234t, 238f
- C-O sole (*Pleuronichthys coenosus*), 18t, 60, 61f, 70, 236t, 535t
- Costa, M. J., 144
- Costanso, Miguel, 568
- Cottidae (sculpins), 9f, 10f, 90f, 91f, 105, 106t, 329; on continental shelf, 172, 175, 178t, 180t, 186; egg size and, 501t; in kelp bed/rocky reef, 70, 234t; larval stages of, 289, 304f; movement of, 532t–533t, 543; reproduction and, 483, 491, 493, 495, 498–99, 500, 502, 504, 507; in rocky intertidal zone, 209t, 210, 212, 215, 217f. *See also* *Artedius* spp.; *Clinocottus*; *Oligocottus*
- Cottus asper* (prickly sculpin), 126t, 128f, 129f, 132f, 296
- Cottus bairdi* (mottled sculpin), 501
- courtship, 487–95; acoustic signals and, 493; coloration and, 488, 489, 490, 493, 494; functions of, 488–89; spawning and, 488–89, 494
- cow cod (*Sebastes levis*), 18t, 23t, 100f, 103, 537t, 589; in deep rock habitat, 255, 262f, 263t; larval stages of, 285, 288f
- Cowen, R. K., 76, 241, 417, 546
- Coyer, J. A., 544
- crepuscular patterns, 543, 549
- crescent gunnel (*Pholis laeta*), 17t, 21t, 209t, 485, 522t
- crested bigscale (*Poromitra crassiceps*), 352t, 358f, 365, 366t, 370f; trophic relations of, 402f
- crevice kelpfish (*Gibbonsia montereyensis*), 18t, 90f, 111f, 215t, 216f, 238f, 399f
- crisscross prickleback (*Plagiogrammus hopkinsii*), 17t, 24t, 91f, 236t, 238f
- croakers. *See* *Genyonemus* spp.; Sciaenidae (croakers)
- Croker, R. S., 588, 590
- Cross, J. N., 170, 212, 398
- crustaceans, as foodsource, 361, 394, 403, 406, 606
- Cryptacanthodidae, 485, 505, 522t
- cryptic species, 64, 91f, 92f, 106, 109f; predation and, 428–29, 432; in rocky subtidal reefs, 238f; surveys of, 231–32
- Cryptosaras couesii*, 487, 520t
- Cryptotrema corallinum*, 18t, 24t
- Crystallodystes cookei*, 522t
- Cubiceps* spp. (cigar fish), 333, 523t, 559
- cunner (*Tautoglabrus adspersus*), 419
- curlfin turbot (*Pleuronichthys decurrens*), 18t, 98f, 398, 535t, 606; on continental shelf, 179t, 181f; foraging guild, 192f, 194f
- Cushing, D. H., 308–9
- Cushman, J. H., 554
- cuskeels (Ophidiidae), 106, 177t, 180t, 197f, 353t, 369; *Lepophidium* spp., 177t, 182f, 183f, 195f, 196f; movement and activity of, 535t, 544. *See also* *Chilara taylori*
- cutlassfish (*Trichiurus nitens*), 20t, 25t, 327f, 332
- cutthroat trout (*Oncorhynchus clarkii*), 17t, 21t, 29, 130t
- Cyaena* sp. (scyphomedusa), 328f
- Cyclopsetta panamensis*, 308f
- Cyclothone acclinidens* (benttooth bristlemouth), 350t, 354, 356f, 360t, 362–63, 370f, 402f; larval forms of, 276, 280; swimbladder of, 365, 366t, 367
- Cyclothone* spp. (bristlemouths), 354, 365, 368, 370f; *C. atraria*, 519t; *C. pallida* (tan bristlemouth), 350t; *C. pseudopallida* (slender bristlemouth), 276, 280; *C. signata* (showy bristlemouth), 276t, 350t, 354, 356f, 360t, 362–63, 366t, 370f; larval stages of, 276, 277f, 279f, 280, 368

- Cyema atrum*, 519t
- Cymatogaster aggregata* (shiner perch), 18t, 71, 72f, 88–89, 94f, 475, 631; in bays and estuaries, 127t, 128f, 129f, 132f, 134f, 136t, 138t, 145t; on continental shelf, 176, 178t, 181f; distribution patterns of, 546; foraging guilds, 192f, 194f; habitat of, 102, 103, 105t, 111f; in kelp bed/rocky reef, 235t, 248; pollution and, 606; reproduction of, 494; seasonal movement by, 533t, 542; in surf zone, 150, 152f, 153t, 154f, 160–61t; trophic relations and, 396f, 397f, 402, 403
- Cynoglossidae (tonguefishes), 298, 329; on continental shelf, 175, 179t, 180t, 189. *See also* *Symbolophorus* spp.
- Cynoscion* spp. (sea trout), 150; *C. parvipinnis*, 19t, 24t
- Dactylopterus volitans*, 520t
- Dactyloscapus* spp., 522t
- daggertooth (*Anotopterus pharao*), 327f, 330, 519t
- Dagorn, L., 540
- Dalatiidae (dwarf sharks), 329, 353t
- damsel fish (Pomacentridae), 65, 74, 76, 106t, 506, 547; in kelp bed/rocky reef, 235t, 237f; movement of, 535t, 548; reproduction and, 488, 489, 491, 493, 495, 503. *See also* *Chromis* spp.; garibaldi (*Hypsypops rubicundus*)
- Danaphos oculatus* (bottlelight), 306, 350t, 354, 356f
- darkblotched rockfish (*Sebastes crameri*), 17t, 21t, 257, 589
- darkwing flyingfish (*Hirundichthys rondeletii*), 331f
- Darling, J. D. S., 220
- Dascyllus* spp., 503, 522t
- Dasyatis* spp. (stingrays): *D. diptera*, 20t; *D. longa*, 518t; *D. violacea*, 326f, 331, 337t
- data matrix, 83
- Daugherty, 576
- Davidson Current, 347, 361
- Davis, G. E., 229
- Davis, J. C., 584
- Davis, J. L. D., 208, 212
- Davis, M. K., 572
- Dawson, Michael N., 5, 11, 14, 623
- DDT and PCB contamination, 595–603, 605, 607f. *See also* pollution
- de Bary, Anton, 554
- Decapterus muroadsi*, 20t, 23t
- Decodon melasma* (blackspot wrasse), 7t, 20t, 24t
- decorated warbonnet (*Chirolophis decoratus*), 7t, 17t, 236t
- deep bank (DBNK), 82t, 85t, 86f, 99, 104t
- deepbody anchovy (*Anchoa compressa*), 18t, 95f, 159f, 160t, 173; in bays and estuaries, 126t, 134f, 136t, 138t, 145t; in surf zone, 150t, 151, 154; trophic relations and, 396f, 401
- deep reef (DRF), 82t, 84f, 100f, 104t
- deep rocky reef (DDRF), 85t, 86f, 253–64; anthropogenic structures, 260–64; bottom assemblage, 261, 263; midwater assemblages, 260–62; natural outcrops, 255; overview of fish assemblages, 255–59; pipelines and, 264; shell mound and, 264
- deep sea. *See* bathopelagic zone (deep sea)
- deep-sea skate (*Bathyraja abyssicola*), 19t, 22t, 353t, 372f
- deep-sea sole (*Embassichthys bathybius*), 354t, 372f
- deep slope (DSLPL), 85t, 86f
- deepwater cornetfish (*Fistularia corneta*), 7t, 14, 20t, 25t
- deepwater eelpout (*Lycodapus endemoscotus*), 354t
- deepwater serrano (*Serranus aequidens*), 178t, 180t, 196f
- Delolepsis gigantea*, 17t, 21t
- delta smelt (*Hypomesus transpacificus*), 126t, 128, 129, 130t, 131, 473, 615
- DeMartini, E. E., 152, 154, 159, 507; on kelp bed/rocky reef habitat, 231, 241, 244, 245, 246; on movement patterns, 540, 541, 544
- dendrograms, 7–8, 11f, 14f, 86f; of bays and estuaries, 8, 11f, 14f, 124f
- Denman, William, 572
- Denton, E., 73
- Derichthys serpentinus*, 518t
- desiccation, resistance to, 212
- Desmodena lorum*, 20t, 23t
- Desmond, J. S., 138, 139
- detritivores, 394
- Diablo Canyon, 294
- diadromous fishes, 124
- Diamond, J., 451
- diamond stingray, 137f
- diamond turbot (*Hypsopsetta guttulata*), 19t, 181f, 396f, 402, 535t, 600; in bays and estuaries, 127t, 129f, 136t, 137f; larval stages of, 291, 294f, 299
- diaphanous hatchetfish (*Sternopyx diaphana*), 350t, 354, 356f, 402f, 519t
- Diaphus* spp. (headlightfish), 276t; *anderseni* (Andersen's lanternfish), 351t; *D. theta* (California headlightfish), 305, 351t, 356, 357f, 359, 360t, 363, 366t, 402f
- Dicentrarchus labrax* (European sea bass), 497
- diel behavior, 188, 468, 507, 543–44
- diet. *See* feeding
- Dill, W. A., 130
- Diodon* spp. (porcupinefish), 77; *D. holocanthus* (balloon fish), 7t, 21t, 25t, 523t; *D. hystrix*, 21t, 236t
- Diogenichthys atlanticus* (longfin lanternfish), 351t, 370f; larval stages of, 276t, 277f, 279, 282f, 290f, 368
- Diogenichthys laternatus* (Diogenes lanternfish), 351t, 370f; larval stages of, 277, 277f, 279, 280, 287, 293f
- Diplectrum* spp. (sand perch), 496; *D. labarum* (highfin sand perch), 178t, 182f, 183f, 195f, 196f; *D. maximum* (greater sand perch), 7t, 20t, 24t; *D. pacificum* (Pacific sand perch), 178t, 181f, 194f, 521t
- Diplodus sargus*, 521t
- Diplogrammus pauciradiatus*, 523t
- Diplophus taenia* (bristlemouth), 365
- Diplospinus multistratus*, 309f
- Diretmus argenteus*, 311f
- disease, pollution-related, 603–7
- dispersal, 49, 526f
- distribution patterns, 5–6, 8, 76, 546
- disturbance, 466–79; anthropogenic, 467, 473 (*See also* pollution); competition and, 472, 477; defined, 467; direct versus indirect effects, 468; fitness and growth, 472, 473; freshwater inflows and salinity, 473; habitat selection and use, 469–70, 473, 474–75; hypoxia and, 468–73; natural disturbance, 467–68; predation and, 442; rate of, 108; storms, 473–77; theoretical context of, 466–67
- diurnal patterns, 75, 543, 547
- diversity patterns, 83, 85, 107–12, 442–43
- dogfish sharks (Squalidae), 106t, 175, 177t, 180t. *See also* *Squalus acanthias*
- dogtooth lampfish (*Ceratoscopelus townsendi*), 359, 366t, 369, 370f; larval stages of, 276t, 277f, 279, 280, 305
- Dolan, R., 3
- Dolichopteryx longipes* (brown snout spookfish), 349t
- dolphinfish (*Coryphaena hippurus*), 20t, 89f, 324, 326, 331f, 332, 333, 337t
- Dormitator latifrons*, 20t, 24t
- Dorosoma petenense* (threadfin shad), 161t, 612t, 613f, 617
- Dover sole. *See* *Microstomus pacificus*
- dragonfishes (Stomiidae), 329, 350–51t
- Drazen, J. C., 373, 374
- dreamers (Oneirodidae), 17t, 352t, 358, 390, 395, 402t, 520t
- drift algal beds, 111, 151–52, 154f
- driftfish (*Psenes* spp.), 328f, 333, 559
- dusky shark, 337t
- dwarf lanternfish, 370f
- dwarf perch. *See* *Micrometrus minimus*
- dwarf perch (*Micrometrus minimus*), 18t, 111f, 145t, 161t, 213, 235t, 499, 533t; in surf zone, 95f, 150t, 151f, 152, 154f, 155t
- Dyer, B. S., 621
- East Pacific Barrier, 69
- Ebeling, A. W., 359, 363–64, 437–38, 456–57, 466; on kelp bed/rocky reef habitat, 231, 241, 244, 245, 246; on movement and activity patterns, 526, 531, 547; on storm disturbance, 475, 476
- Eber, L. E., 6
- Echeneidae (suckerfishes), 324–25. *See also* *Remora* spp.
- Echeneis naucrates* (sharksucker), 20t, 24t, 324, 329f
- Echinomacrurus occidentalis*, 353t
- Echinorhinus cookei*, 20t, 23t
- ecological classification, 81–112; abundance within species groups, 88t; approach to, 81, 83–85; bay and estuary, 123–24; biological basis for, 85–87; habitat generalists, 102–3; patterns of diversity and, 107–12; physical bases for, 97–102; species richness and, 109–12; taxonomic composition of assemblages, 103, 105–6. *See also* specific habitat
- ecological variables, 5
- economic exclusion zones (EEZs), 525
- ecosystem collapse, 56, 58
- ecosystem conservation, 589–90
- eelgrass (*Zostera marinus*), 111, 134, 397, 475
- eelpout (*Melanostigma pammelas*), 18t, 22t, 348, 352t. *See also* *Lycodapus*; *Lycodes* spp.; Zoarcidae
- eels. *See* *Anguillidae* (freshwater eels)
- egg laying. *See* oviparity (egg laying)
- Eigenmann, Carl and Rosa, 253
- Ekman, S., 69
- elasmobranchs, 94, 484, 504
- electric ray. *See* Pacific electric ray (*Torpedo californica*)
- Elegatis bipinnulatus* (rainbow runners), 324
- elephant seals (*Mirounga angustirostris*), 530, 542, 561
- elkhorn sculpin (*Alcichthys alcicornis*), 499, 501–2
- Elkhorn Slough, 8, 131–32, 133, 400, 619
- Elliot, M., 119
- Elliott, J., 560–61
- El Nino Southern Oscillation. *See* ENSO
- Elops affinis* (machete), 7t, 20t, 518t
- Embassichthys bathybius* (deep-sea sole), 354t, 372f

- Embiotoca jacksonii* (black perch), 5, 19t, 34t, 75, 92f, 111f, 150t, 585t; activity patterns of, 544, 547, 549, 633; in bays and estuaries, 127t, 128f, 132f, 134f, 136t, 145t; in Channel Islands, 48f, 49f; competition and, 450, 457–60, 461–62f, 463f; feeding of, 390; habitat range of, 102, 103, 105t; in harbors, 157t, 159, 161t, 162t; in kelp bed/rocky reef, 235t, 237f, 239f, 240f, 241f, 404; movement of, 533t, 542; phylogeographic structures of, 34t, 38, 39f, 40, 41, 624; predators and, 64, 430, 431; recruitment of, 412f, 415, 416; reproduction of, 496, 522t; in surf zone, 150t, 154f, 155t; symbiotic relations of, 556f, 557t; territoriality of, 453t, 454, 455; trophic relations and, 400f
- Embiotoca lateralis* (striped seaperch), 18t, 75, 92f, 533t, 585t; activity patterns of, 544–45; competition and, 450, 453f, 457–60, 461–62f, 463f; in kelp bed/rocky reef, 235t, 237f, 240f, 404; phylogenetic structures of, 34t, 40; recruitment of, 415, 416; in surf zone, 111f, 151, 152f, 153t
- Embiotocidae (surfperches), 9f, 10f, 106t, 235t, 557t, 591, 630; activity patterns of, 533t, 539, 544; on continental shelf, 175, 178t, 180t, 186, 258t; evolution of, 71, 72, 74, 75, 77; as habitat generalists, 87, 105; pollution and, 602; predation and, 437–38; recreational fishing of, 581; reproduction of, 484, 498, 504, 505; storm disturbance and, 474–77, 478f; subsistence fishing of, 567, 568; in surf zone, 152f, 153t, 155t
- Emblemaria hypacanthus* (signal blennies), 432, 503, 522t
- emergence behavior, 208, 210
- Emery, K. O., 228
- Emmet, R. L., 125
- endangered species, 125, 128–30
- endosymbiotic mutualism, 557
- Engle, J. M., 229
- English sole. *See* *Pleuronectes/Parophrys vetulus*
- Engraulidae (anchovies), 72, 106t, 149, 494. *See also* *Anchoa* spp.; anchovy-sardine regimes
- Engraulis mordax* (northern anchovy), 19t, 72, 186, 233t, 260, 429, 590; abundance of, 89, 91, 160t, 161t, 336t; in bays and estuaries, 125, 127t, 128f, 129, 132f, 138t, 145t; competition and, 450; diel movement of, 543; feeding of, 390, 394; habitat of, 102, 103, 105t; larval stages of, 273–78f, 280, 281–84f, 291, 292, 294f, 299, 303–5; in pelagic zone, 94f, 159f, 330, 332f, 335t, 337t; schooling by, 324; in surf zone, 150t, 151f, 152, 155t; trophic relations and, 397f, 398f, 401f
- Engyophrys sanctilaurentii* (speckletail flounder), 7t, 20t, 25t, 179t, 182f, 195f
- Enhydra lutris* (sea otter), 624, 627, 628
- Enneanectes reticulatus* (flag triplefin), 236t
- Enophrys bison* (buffalo sculpin), 17t, 234t, 485, 486, 500t
- Enophrys taurus* (bull sculpin), 17t
- ENSO (El Niño-Southern Oscillation), 174, 240, 330, 466, 504, 615, 630–31; bay habitat and, 129, 135, 141; distributional patterns and, 4–5, 6, 14–15; La Niña event and, 4–5, 222, 283, 286; larval populations and, 282–84, 285–93f, 298–302, 303, 628; phylogeography and, 37, 41; recruitment and, 414, 417; reef coral damage by, 69; rocky intertidal zone and, 221–22; species density and, 249
- environmental legislation, 588, 591
- environmental potential for polygamy (EPP), 502
- Environmental Protection Agency (USEPA), 168, 170, 602
- Eopsetta exilis*. *See* *Lyopsetta exilis*
- Eopsetta jordani* (petrale sole), 18t, 179t, 182f, 183f, 185f, 193; foraging guild, 193, 195f, 196f, 197f
- epibranchial organs, 392–93
- Epinephelinae* spp., 58, 497
- Epinephelus* spp. (cabrilla): *E. analogus* (spotted cabrilla), 20t, 24t, 234t; *E. guttatus*, 521t; *E. labrifomis* (flag cabrilla), 234t; *E. niphobles*, 20t, 25t; *E. striatus*, 521t
- epipelagic fishes, 320, 330, 342–43, 345t, 368; trophic relations in, 401f, 405–6
- Eptatretus deani* (black hagfish), 18t, 22t, 101f, 353t, 372f
- Eptatretus stoutii*, 18t, 518t
- Erilepsis zonifer*, 17t, 21t
- Ernogrammus walkeri* (masked prickleback), 7t, 17t, 24t
- Erres zachirus* (rex sole), 333. *See also* *Glyptocephalus zachirus*
- Eschmeyer, W. N., 483
- escolar (*Lepidocybium flavobrunneum*), 20t, 22t, 327f, 330
- Estero de Punta Banda, 136t
- estuaries. *See* bay/estuaries (BE)
- Etrumeus teres*, 21t, 23t, 280f
- Eucinostomus* spp. (mojarra): *E. argenteus* (spotfin mojarra), 178t, 181f, 182f, 194f, 195f; *E. curranii*, 20t, 24t; *E. dowii*, 20t, 25t
- Eucyclogobius newberryi* (tidewater goby), 5, 18t, 132f, 614; in bays and estuaries, 126t, 128f, 129f, 130t, 134f; phylogeography and, 34t, 38, 39f
- eulachon (*Thaleichthys pacificus*), 128–29, 130t
- Euleptorhampus longirostris* (half beak), 20t, 24t
- Euleptorhampus viridis* (ribbon halfbeak), 331f, 333
- Eumicrotremus orbis*, 521t
- Euphausia pacifica* (krill), 371
- Euprotomicrus bispinatus* (pygmy shark), 17t, 331
- Euryhaline species, 88
- Eurypegasus draconis*, 520t
- Eurypharynx pelecyanoides* (umbrellamouth gulper), 348, 519t
- Eutaeniophorus festivus*, 310f
- Euthymus* spp.: *E. affinis*, 20t, 25t; *E. lineatus*, 20t, 24t; *E. pelamis*, 22t
- eutrophication, 145
- evolution, 50, 55–77; Cenozoic influences, 68–70; distribution and latitude, 68–70; extinction-resurgence episodes and, 55–56, 62; for feeding effectiveness, 59–62; mainstream feeding relations and, 58–59; modern teleost fishes, 58; new modes of feeding and, 62–68; other evolutionary lines, 72–73; perciformes, 71–72; percoids, 57–58; pleuronectiformes and, 70–71; predation and, 428–33; scorpaeniformes and, 70; smallness and, 62–64; species composition and, 73–77; teleost history and, 55, 68; teleost-scleractinia connection, 56–57
- Exocoetidae. *See* flyingfish (Exocoetidae)
- extinction-resurgence episodes, 55–56, 62, 625
- Fagerstrom, J. A., 56
- fangtooth (*Anoplogaster cornuta*), 311f, 402f, 520t; as bathypelagic fish, 348, 352t, 358, 366t
- fantail sole (*Xystreus liolepis*), 19t, 145t, 155t, 158t; on continental shelf, 179t, 181f, 182f; foraging guilds, 192f, 194f, 195f; larval stages of, 293, 295f, 300, 301f, 302
- Feder, H. M., 227, 555
- Federal Food Administration, 570
- feeding: biting, 390–91; commensalism, 554, 558–61; competition in, 389, 613; by deep sea fishes, 344, 361, 364–65, 375–76; distribution and, 544–46; epibranchial organs, 392–93; factors determining diet, 388–89; larval survival and, 311, 312; mechanisms in, 387–95; oophagy, 484; overview of, 387–88; pharyngeal and oral jaws, 393–94; in rocky intertidal zone, 218–20; by soft-bottom fishes, 188–90; suction and ram feeding, 391–92; suspension feeding, 392–94; types of prey capture, 389–90
- feeding divergence, 59–68; ambushers, 59–62, 68, 190; dissimulace, 64; herbivores, 65, 66f, 74, 207, 390; increased effectiveness in, 59–62; inedibility, 64–66; new modes, 62–68; nocturnality, 66–68; piscivorous fish, 131, 395, 402; planktivores, 63, 65, 131, 189, 420, 547; smallness, 62–64; straightforward rush, 61–62, 68
- Ferrante, Pietro, 571
- Ferren, W. R., Jr., 120
- Fierstine, H. L., 133
- filetail cat shark (*Parmaturus xaniurus*), 19t, 23t, 101f, 353t, 518t
- fin erosion disease, 603, 604f, 605, 607f
- finescale cusk-eel (*Lepophidium microlepis*), 177t, 183f, 196f
- finescale triggerfish (*Balistes polylepis*), 21t, 23t, 236t, 309f, 523t; on continental shelf, 179t, 194f; reproduction and, 485
- Finney, B. P., 624–25
- fishing and fisheries, 540, 575–76; climate change and, 625–26, 627, 633; foreign, 578–79; management of, 461–62, 586–90; overfishing, 567, 572, 573, 576, 580, 588; subsistence, 524, 567–68, 580. *See also* commercial fisheries; recreational fishing
- Fistularia commersonii*, 520t
- Fistularia corneta* (deepwater cornetfish), 7t, 14, 20t, 25t
- Fitch, J. E., 348, 483
- flag cabrilla (*Epinephelus labrifomis*), 234t
- flag flounder (*Perissia taeniopterus*), 179, 183f, 193, 196f
- flag rockfish (*Sebastes rubrivinctus*), 7t, 18t, 99f, 255, 260f, 263t
- flashlight fishes (Anomalopidae), 66, 370f, 520t, 558f. *See also* *Protomyctophum*
- flatfishes. *See* *Pleuronectiformes*
- flathead sole (*Hippoglossoides elassodon*), 21t, 605
- flounders, 179t, 183f, 185f, 195f, 196f. *See also* *Paralichthyidae*; *Pleuronectidae*
- fluffy sculpin (*Oligocottus snyderi*), 18t, 90f, 399f, 500t; in rocky intertidal zone, 209t, 215t, 216f, 217f, 218, 220f
- flyingfish (Exocoetidae), 73, 326f, 331f, 401f; *Cheilopogon heterurus*, 20t, 25t, 520t; *Cheilopogon pinatibarbatus*, 19t, 22t, 405; *Cheilopogon* spp., 323, 328, 331, 405; *Cypselurus californicus*, 73

- Fodiator acutus*, 20t, 24t
 food web. *See* trophic interactions
 foraging behavior, 67–68, 189–90, 190–92, 531; guilds, 190–92, 192f, 194–97f. *See also* feeding
Forcipiger flavissimus (longnose butterflyfish), 63
 Forrester, G. E., 422, 430, 434, 439–40
 freckled rockfish (*Sebastes lentiginosus*), 18t, 25t, 255
 freshwater fishes: in bays and estuaries, 124, 128f, 132f, 134f
 fringed flounder, 137f
 fringeheads (*Neoclinus* spp.), 18t, 19t, 23t, 97f, 236t, 238f, 247; in kelp bed/rocky reef, 236t, 238f, 247
 Fuentes, E. R., 451
Fundulus heteroclitus (mummichog), 139, 470, 520t
Fundulus parvipinnis (California killifish), 10–11, 19t, 34t, 41, 42f, 88, 533t; abundance of, 142, 143f, 145t; in bays and estuaries, 93f, 120, 126t, 129f, 132f, 134f, 137f, 138; pollution and, 601; trophic relations and, 396f, 401
 Gadidae (cods), 175, 180t
Gadus macrocephalus (Pacific cod), 17t, 21t, 605
Galeocerdo cuvier, 20t, 25t
Galeorhinus (soupfin shark), 19t, 89f, 233t, 337t, 576
Gambusia affinis (western mosquitofish), 126t, 132f, 535t, 612t, 616–17
 garibaldi (*Hypsypops rubicundus*), 19t, 65f, 74, 92f, 404, 522t, 555, 589; activity patterns of, 535t, 544; in harbors, 157t, 162t; in kelp bed/rocky reef, 235t, 237f, 239f, 241f, 243f; reproduction and, 489, 490f, 493, 495, 496, 501; territoriality of, 451, 453t; trophic relations and, 400f
 Garrison, L., 197
 Gartner, J. V., Jr., 354, 368, 406
Gasterosteus aculeatus (threespine stickleback), 18t, 153t, 520t; in bays and estuaries, 126t, 128f, 129f, 132f, 142f; reproduction and, 488, 489, 491–92
 Geibel, J. J., 245–46, 539, 542
Gelidium robustum (algae), 457, 459–60
Gempylus serpens (snake mackerel), 21t, 25t, 327f, 329, 332
 gender allocation, 487, 488t, 495–98, 502–4, 505f, 506
 gene flow, 27, 38, 48, 49–50, 623; glaciations and, 33, 623–24
 genetic divergence, 31, 32
 gene trees, 27, 36. *See also* phylogeography
Genyonemus lineatus (white croaker), 19t, 96, 102, 105t, 390, 631; commercial fishing and, 574, 579, 590; on continental shelf, 98f, 172, 178t, 181f, 182f; foraging guild, 192f, 194f, 195f; in harbors, 157t, 159f, 160–61t, 162; larval stages of, 276t, 291, 294, 297f, 299, 305; movement of, 535t, 540; pollutants and, 598, 600, 601, 603, 604, 605–6; recreational fishing and, 582; subsistence fishing and, 568; in surf zone, 150t, 154f, 155t, 156; trophic relations and, 397f
Genyonemus spp. (croakers) larval stages of, 291, 295, 297–98
 Gerkema, M. P., 549
 Gerreidae (mojarra), 175, 180t, 506; spotfin mojarra, 178t, 181f, 182f, 194f, 195f
Gerres cinereus, 521t
 ghostly grenadier (*Coryphaenoides*), 353t, 369, 374, 376
 ghost shrimp (*Neotrypaea* spp.), 559, 560f
 giant grenadier (*Coryphaenoides pectoralis*), 101f, 353t, 369, 372f, 373–74
 giant kelpfish. *See* *Heterostichus rostratus*
 giant kelp (*Macrocystis pyrifera*), 48, 67, 71f, 415, 421, 436, 438–39, 459, 477
 giant lampfish (*Parvilux ingens*), 351t, 366t, 370f
 giant sea bass (*Stereolepis gigas*), 18t, 87, 555, 624; in kelp bed/rocky reef, 92f, 234t, 237f, 239f, 241f; recreational fishing and, 583, 589; reproduction and, 494, 497; trophic relations, 400f, 405
Gibbonsia elegans (spotted kelpfish), 19t, 60, 61f, 92f, 111f, 155t, 532t; hypoxia and, 468; in kelp bed/rocky reef, 236t, 238f; in rocky intertidal zone, 207, 215t, 216f, 217, 220f; in surf zone, 145t, 154f, 155t, 159
Gibbonsia metzi (striped kelpfish), 19t, 90f, 111f, 155t, 236t, 238f; in rocky intertidal zone, 215t, 216f; trophic relations of, 399f, 403
Gibbonsia montereyensis (crevice kelpfish), 18t, 90f, 111f, 215t, 216f, 236t, 399f
Gibbonsia spp. (kelpfishes), 35t, 215t, 477
 Gibson, R. N., 212, 214, 217, 218, 221, 395
Gigantactis spp. (anglerfish), 358, 520t
Gigantura indica, 519t
Gilbertidia sigalutes, 500t, 521t
Gillichthys mirabilis (longjaw mudsucker), 11, 19t, 88, 211, 533t, 617; in bays and estuaries, 93f, 126t, 129f, 132f, 134f, 136t, 138t, 143f; hypoxia and, 472; larval stages of, 303, 304f; phylogenetic relations and, 35t, 43, 44f, 45t, 46t; trophic relations and, 396f, 402
 gill nets, 578f, 590–91
Girella nigricans (opaleye), 19t, 102, 105t, 136t, 150t, 585t; activity patterns of, 534t, 544–45; in harbors, 157t, 158t, 162t; as herbivore, 65, 66f, 74, 394, 403, 404; in kelp bed/rocky reef, 92f, 235t, 237f, 239f, 241f; mutualism of, 556f, 557t, 558; phylogenetic characteristics of, 35t, 41, 43, 44f, 45t, 46t; recruitment of, 414, 419; in rocky intertidal zone, 209t, 215t, 216f
Girella tricuspidata, 522t
 glaciations, 32–33, 36, 37, 41, 69; gene flow and, 33, 623–24
 global warming, 632–33. *See also* climate change
 globalization, in fishing industry, 578–79
Glyptocephalus zachirus (rex sole), 18t, 22t, 101f, 103, 307, 535t; on continental shelf, 179t, 183f; foraging guild, 192f, 194, 195f, 196f, 197f
Gnathophis spp., 19t, 24t, 519t
 goatfishes (Mullidae), 7t, 135t, 329, 506
Gobiesox meandricus (northern clingfish), 18t, 33, 34t, 90f, 236t, 523t; in rocky intertidal zone, 209t, 215t, 216f, 220f, 221; trophic relations and, 399f, 403
Gobiesox spp. (clingfishes), 106t, 217f; *G. eugrammus*, 18t, 25t; *G. papillifer*, 20t, 25t; *G. rhessodon* (California clingfish), 19t, 90f, 212–13, 215t, 216f, 238f, 403; reproduction and, 485
 Gobiidae (gobies), 9f, 10f, 105, 106t, 142f, 506; alien species, 612–14, 619; larval stages of, 289, 294f, 298, 303, 304; movement of, 524, 533–34t; pollution and, 602; reproduction and, 491, 493, 497, 498, 503, 504
Gobiomorphus breviceps, 523t
Gobiosoma bosc (naked goby), 469
 golden lanternfish (*Myctophum aurolaternatum*), 307, 310f, 370f
 golden spotted rock bass (*Paralabrax auroguttatus*), 234t, 497
 golden wrasse (*Halichoeres melanotis*), 235t
 Goldman, K. J., 528, 542
Gonichthys tenuiculus (slendertail lanternfish), 277, 279f, 280, 370f
 gonochorism, 483, 497, 498, 502–3, 504, 507
 Gonostomatidae (bristlemouths), 329, 354, 356f, 359, 365
 gopher rockfish (*Sebastes carnatus*), 19t, 23t, 219t, 258, 270, 585t; in kelp bed/rocky reef, 91f, 234t, 237f, 240; movement of, 531, 536t; recruitment and, 415, 420; territoriality of, 453t, 455–56f, 457f
 Gorbatenko, K. M., 361
 Gosline, W. A., 55, 63, 72, 73
 Gotshall, D. W., 253
Gracillaria spp. (algae), 111, 159. *See also* algae
 Graham, M. H., 624
Gramma loreto, 521t
Grammonus diagrammus, 25t
 Grant, J. W. A., 496
 Grant, W. S., 624
 grass rockfish (*Sebastes rastrelliger*), 18t, 59f, 111f, 234t, 270, 403, 430; in rocky intertidal zone, 90f, 215t, 216f, 217f, 219t; in surf zone, 152, 153t, 155t, 157t
 graveldiver (*Scytalina cerdale*), 17t, 90f, 215t, 216f, 220f, 522t; feeding of, 399f, 403
 gray remora (*Remora brachyptera*), 21t, 25t, 325
 gray smoothhound (*Mustelus californicus*), 19t, 95f, 127t, 132f, 233t; in surf zone, 150t, 151, 153, 156f, 157t, 158t
 greater sand perch (*Diplectrum maximum*), 7t, 20t, 24t
 Greely, T. M., 368
 Green, J. M., 543
 greenblotched rockfish (*Sebastes rosenblatti*), 18t, 23t, 102, 105t, 183f; in deep rock habitat, 100f, 255, 262f, 263t
 greenlings. *See* Hexagrammidae
 green-spotted rockfish (*Sebastes chlorostichus*), 18t, 22t, 178t, 183f, 196f; in deep rock habitat, 100f, 255, 262f, 263t; movement of, 528, 536t, 539
 greenstripe rockfish (*Sebastes longatus*), 18t, 22t, 102, 105t, 537; in deep rock habitat, 100f, 255, 262f, 263t
 green sturgeon (*Acipenser medirostris*), 18t, 125, 126t, 128f, 130t
 grenadiers. *See* *Coryphaenoides* spp.; Macrouridae (grenadiers); *Nezumia* spp.
 Grossman, G. D., 212, 219, 220, 221
 grouper. *See* *Mycteroperca* spp.
 Gruber, D., 291, 297, 298
 grunion. *See* *Leuresthes tenuis/sardina*
 grunts. *See* Haemulidae (grunts)
 grunt sculpin (*Rhamphocottus richardsonii*), 17t, 21t, 213, 500t, 512t
 Grutter, A. S., 555
 Guadalupe cardinalfish (*Apogon guadalupensis*), 19t, 24t, 234t, 485
 guitarfish, 176, 177t, 180, 183f, 189, 233t. *See also* banded guitarfish (*Zapteryx exasperata*). *See* Rhinobatidae; *Zapteryx exasperata*
 gulf grouper (*Mycteroperca jordani*), 19t, 25t, 234t
 Gulf of California, 27, 41–43, 46
 gulf sanddab (*Citharichthys fragilis*), 7t, 20t, 277f, 278f, 280
Gummelichthys spp., 523t

- gunnels (*Pholis* spp.), 9f, 10f, 72, 90f, 209t, 217f; *P. clemensi*, 17t, 21t; *P. laeta* (crescent gunnel), 17t, 21t, 209t, 485, 522t; *P. ornata* (saddleback gunnel), 17t, 153t, 399f; *P. schultzi*, 17t, 21t; *Ulvicola sanctaerosae* (kelp gunnel), 18t, 24t, 72, 72f, 236t. *See also* penpoint gunnel; rockweed gunnel
- Gymnothorax meleagris*, 518t
- Gymnothorax mordax* (California moray), 19t, 24t, 95f, 555, 558f; activity patterns of, 545, 549; in harbors, 157t, 158; in kelp bed/rocky reef, 233t, 238f, 239f, 241f, 243f
- Gymnura marmorata* (butterfly ray), 20t, 135t, 518t
- habitat. *See* ecological classification; *specific habitat*
- Haedrich, R. L., 559
- Haemulidae (grunts), 66, 69, 74, 235t, 393, 534t; bronzestripe (*Orthopristis reddingi*), 178t, 181f, 182f, 194f, 195f; on continental shelf, 175–76, 178t, 180t. *See also* *Anistotremus* spp.; salema (*Xenistius californiensis*)
- Haemulon flavolineatum* (grunt), 546
- Haemulon sciurus*, 521t
- Haemulopsis axillaris* (yellowstripe grunt), 178t, 182f, 194f, 195f
- hagfish (Myxinidae), 103, 106t, 353t, 578–79; black hagfish (*Eptatretus deani*), 18t, 22t, 101f, 353t, 372f
- hakes, 472. *See also* *Merluccius* spp.
- halfbanded rockfish (*Sebastes semicinctus*), 19t, 23t, 99f, 178t, 183f, 196f, 631; in deep rock habitat, 255, 257, 260f, 263t
- halfbeak, 331f, 333. *See also* *Hyporhamphus rosea*
- halfmoon (*Medialuna californiensis*), 19t, 35t, 92f, 157t, 162t, 400f; as herbivore, 65, 66f, 74, 394, 404; in kelp bed/rocky reef, 235t, 237f, 239f, 241f; symbiotic relations and, 556f, 557t, 558
- halfspotted tonguefish (*Symphurus atrimentatus*), 179t, 182f, 195f
- halibut. *See* *Hippoglossus stenolepis*; *Paralichthys californicus*
- Halichoeres semicinctus* (rock wrasse), 19t, 87, 92f, 162t, 400f; cleaner behavior of, 556f, 557t; in kelp bed/rocky reef, 235t, 237f, 239f, 241f, 404; movement and activity patterns of, 534t, 544, 546, 548; phylogenetic relations and, 44f, 45t, 46t, 49; prey dissemblance and, 64; reproduction and, 487, 490, 491f, 496, 497; in surf zone, 150t, 157t
- Halichoeres* spp. (wrasses): *H. chrysus*, 549; *H. dispilus* (chameleon wrasse), 92f, 235t, 237f, 241f; *H. (golden wrasse)*, 235t
- hammerhead shark, 89f, 337t, 538t
- Hanan, D. A., 336
- Harbison, G. R., 307
- harbors, 82t, 157–63
- hardfin marlinsuckers (*Rhombochirus osteochirus*), 25t, 325
- Hare, S. R., 15
- Harengula thrissina*, 20t, 25t
- Hastings, P. A., 432, 623
- hatchetfish (*Sternoptyx* spp.), 365, 366t, 402f, 406, 519t; *S. diaphana* (diaphanous hatchetfish), 350t, 354, 356f, 402f, 519t; *S. obscura*, 350t. *See also* *Argyrolepecus* spp.
- Hayakawa, Y., 499
- Hayden, B. P., 3
- headlight fish. *See* *Diaphus* spp.
- Hecker, B., 175
- Hedgpeth, J. W., 174
- Helfman, G. S., 548
- Hemianthias signifer*, 19t, 311f
- Hemilepidotus* spp., 500; *H. gilberti*, 499; *H. hemilepidotus* (red Irish lord), 17t, 485, 486f, 498, 500t, 521t; *H. spinosus* (brown Irish lord), 17t, 91f, 234t, 238f, 296f
- Hemiramphus* spp., 73; *H. saltator*, 20t, 25t
- Hemitripteris villosus* (Japanese sea raven), 499, 521t
- Heneman, B., 589
- Hensen, V., 271
- Herbinson, K.T., 151, 187
- herbivores, 394, 403, 404
- hermaphroditism, 343, 374, 487, 488t, 506; gonochorism and, 502–3, 504; mating systems and, 496–98
- Hermosilla azurea* (zebraperch), 15, 19t, 87, 544, 630; genetic characteristics of, 44f, 45t, 46t; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 92f, 235t, 237f, 241f, 404; symbiotic relations and, 556f, 558; trophic relations and, 394, 400f
- Herrgesell, P. L., 123
- herring. *See* Clupeidae (herrings)
- Hertig, M., 554
- Hesperonoe advector* (scale worm), 559f
- Hessler, R. R., 373
- Heteroclinus* spp., 522t
- Heterodontus francisci* (horn shark), 19t, 95f, 137f, 394, 429, 518t, 534t; activity patterns of, 544, 546, 549; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 233t, 239f, 241f; in surf zone, 153, 156f
- Heteropriacanthus cruentatus*, 521t
- Heterostichus rostratus* (giant kelpfish), 19t, 103, 105t, 111f, 162t, 418, 522t; in bays and estuaries, 136t, 137f, 145t; cleaner behavior of, 555, 557t; in kelp bed/rocky reef, 71, 71f, 96, 97f, 236t, 238f, 239f; movement of, 532t, 544; predation and, 430, 431f, 436; in surf zone, 150t, 151f, 152, 154f, 155t; trophic relations and, 400f, 405
- Hexagrammidae (greenlings), 70, 91f, 106t, 234t, 263t, 329; on continental shelf, 175, 178t, 180t; larval stages of, 270, 295; reproduction and, 485, 490–91, 495, 501, 507; symbiotic relations of, 560–61. *See also* *Oxylebius pictus* (painted greenling)
- Hexagrammos decagrammus* (kelp greenling), 17t, 215t, 259, 263t, 333, 585t; feeding of, 392; in kelp bed/rocky reef, 70, 71f, 91f, 234t, 237f, 240f; reproduction of, 490, 491f, 495, 501
- Hexagrammos lagocephalus* (rock greenling), 234t
- Hexagrammos superciliosus*, 17t
- Hexanchus griseus*, 18t
- Higgins, Elmer, 572
- highbrow crestfish (*Lophotus cristatus*), 327f, 332
- high cockscomb (*Anoplarchus purpureus*), 17t, 35t, 90f, 498, 522t; in rocky intertidal zone, 209t, 210, 213, 215, 216f, 220; trophic relations and, 399f, 403
- highfin dragonfish (*Bathophilus flemingi*), 351t, 357f
- highfin lizardfish (*Bathysaurus mollis*), 353t, 369, 372f, 374
- highfin sand perch (*Diplectrum labarum*), 178t, 182f, 183f, 195f, 196f
- Hill, C. W., 135
- Hippocampus ingens* (Pacific seahorse), 20t, 25t, 135t, 137f, 484
- Hippoglossina bollmani* (spotted flounder), 179t, 182f, 195f
- Hippoglossina stomata* (bigmouth sole), 19t, 23t, 98f, 236t, 398f, 535t, 606; on continental shelf, 179t, 182f, 183f; foraging guild, 192f, 195f, 196f; larval stages of, 293, 295f, 302
- Hippoglossoides elassodon* (flathead sole), 21t, 605
- Hippoglossus stenolepis* (Pacific halibut), 17t, 525, 580
- Hirundichthys rondeletii* (darkwing flyingfish), 331
- Histrio histrio* (anglerfish), 324
- Hixon, Mark A., 433, 441, 455, 459, 466, 496; on movement patterns, 526, 531, 547
- Hjort, J., 307–8, 309, 434
- Hobson, E. S., 75, 188, 214, 227, 485; on movement patterns, 544, 546, 548; on predation, 440–41; on symbiotic relations, 555, 556, 561
- Hohenshell, A. B., 582
- Holbrook, Sally J., 245, 429, 430–31, 459–60
- Holder, Charles Frederick, 580, 582, 589
- Holocentridae (soldierfishes), 329
- holoepipelagic fish, 324–25, 326–28f, 330t, 331
- Holthymia* spp.: *H. latifrons* (streaklight tubeshoulder), 369; *H. macrops* (bigeye searsid), 350t; *H. melanocephala* (searsid), 350t
- home range, and habitat complexity, 525, 531, 539–40
- honeycomb rockfish (*Sebastes umbrosus*), 19t, 24t, 99f, 255, 260f, 586
- Hoplostethus atlanticus* (orange roughy), 66, 520t
- Horn, Michael H., 5, 9, 13t, 75, 158, 628; bay and estuary study by, 6, 10, 122, 137, 140; on monkeyface pricklyback, 212, 530, 558
- horn shark. *See* *Heterodontus francisci*
- hornyheaded turbot. *See* *Pleuronichthys verticalis*
- hound sharks. *See* *Mustelus* spp.
- Hovden, Knut, 572
- Hovey, T. E., 494, 506
- Howard, D. F., 417, 421
- Howella sp., 521t
- How to Fish the Pacific Coast* (Cannon), 590
- Hubbs, C. L., 11, 123
- human-health risks, pollution and, 601–2
- human impact, 270
- hundred-fathom codling. *See* *Physiculus rastrelliger*
- Hunter, J. R., 305
- Hydrolagus collieri* (spotted ratfish), 18t, 177t, 183f, 185f, 394, 532t; foraging guild, 196f, 197f; habitat of, 101f, 102, 103, 105t
- hydrophones, 528, 530, 540
- Hygophum atratum* (thickhead lanternfish), 277, 279f, 280, 370f
- Hyperoglyphe antarctica*, 523t
- Hyperprosopon anale* (spotfin surfperch), 18t, 97f, 152f, 153t, 155t
- Hyperprosopon argenteum* (wall-eye surfperch), 18t, 67f, 95f, 186, 397f, 585t, 631; activity patterns of, 533t, 544, 546; in coastal pelagic zone, 155t, 157t, 159f, 160–61t; in kelp bed/rocky reef, 235t, 239f; in surf zone, 150–51, 150t, 152f, 153t, 154f

- Hyperprosopon ellipticum* (silver surfperch), 18t, 91f, 235t, 240f, 585t; in surf zone, 152f, 153t
- Hypnys gilberti*, 19t
- Hypomesus pretiosus* (surf smelt), 17t, 72, 494–95, 519; in bays and estuaries, 93f, 127t, 128f, 129f, 132f; in surf zone, 151, 152f, 153t
- Hypomesus* spp. (smelt): *H. nipponensis* (wakasagi), 130t, 131, 612t, 615–16; *H. transpacificus* (delta smelt), 126t, 128, 129, 130t, 131, 473, 615
- Hypoplectrodes maccullochi*, 521t
- Hyporhamphus naos*, 20t, 25t
- Hyporhamphus rosea* (California halfbeak), 19t, 73, 88, 93f, 135t, 136, 328
- Hyporhamphus unifasciatus*, 520t
- hypoxia, 207, 210, 211, 468–73; effects on growth and fitness, 472–73; feeding and trophic interaction, 471; habitat selection and use, 469–70; reproduction and, 471–72
- Hypsoblennius gentilis* (bay blenny), 19t, 95f, 111f, 236t, 455, 532t; in bays and estuaries, 126t, 134f, 136t, 137f, 145t
- Hypsoblennius gilberti* (rockpool blenny), 19t, 90f, 236t, 403; movement of, 531, 532t, 543; in rocky intertidal zone, 209t, 210, 211, 215t, 216f
- Hypsoblennius jenkinsi* (mussel blenny), 19t, 92f, 136t, 236t, 238; movement patterns of, 531, 532t; phylogenetic characteristics of, 43, 44f, 45t, 46t; territoriality of, 451, 453t
- Hypsoblennius* spp. (blennies), 236t, 333, 334f, 522t; larval forms of, 293, 295f, 298, 303, 304f; reproduction of, 492, 493. *See also* Blenniidae (cometooth blennies)
- Hypsopsetta guttulata* (diamond turbot), 19t, 181f, 396f, 402, 535t, 600; in bays and estuaries, 127t, 129f, 136t, 137f; larval stages of, 291, 294f, 299
- Hypsurus caryi* (rainbow seaperch), 18t, 91f, 235t, 239f; cleaner behavior of, 556f, 557t; movement of, 533t, 542, 544–45; niche overlap and, 457, 458f; in surf zone, 152, 155t, 162t
- Hypsypops rubicundus*. *See* garibaldi
- Hyrolagus collicii*, 518t
- Hysteroecarpus traski*, 484f
- Icanberry, J. W., 294, 295
- Icelinus quadriseriatus* (yellowchin sculpin), 19t, 23t, 398f, 403, 532t; on continental shelf, 98f, 173, 178t, 182f; foraging guilds, 192f, 195f
- Icelinus* spp. (sculpins): *I. burchami*, 17t, 22t; *I. cavifrons*, 19t, 23t; *I. filamentosus* (threadfin sculpin), 18t, 22t, 178t, 183f, 196f, 258t; *I. fimbriatus*, 17t, 24t; *I. oculatus*, 18t, 22t; *I. tenuis*, 18t, 22t
- Ichthyococcus irregularis* (bullbog lightfish), 350t
- ichthyoplankton assemblages, 269–313; in bays and estuaries, 142–43, 303–5; CalCOFI samplings, 269–70, 271, 275–302, 313, 361, 362, 368–69, 628; coastal assemblages, 290–303; ENSO events and, 282–84, 285–93f, 298–302, 628; geographic and seasonal distribution, 277–80, 296–98; historic overview, 269–70; interannual and decadal changes, 281–87, 298–302; nearshore, 289–90; ontogenetic stages, 272–75; PDO and, 283, 284, 285–93f, 299–302f, 303; samplings, 270–72; sardine and anchovy regimes, 281–82, 283–84f; specializations, 306–7, 308–11f; trophic relationships, 307–13; vertical distribution, 305–6
- Ichthyos lockingtoni* (medusa fish), 18t, 22t, 328f, 330, 559; larval stages of, 276, 277–78f, 279, 286f
- Icosteus aenigmaticus* (ragfish), 17t, 22t, 327f, 330, 523t
- Idiacanthus antrostomus* (black dragon), 307, 310f, 351t, 356, 357f, 366t, 370f
- Il'inski, E. N., 361
- Ilypnys gilberti* (cheekspot goby), 95f, 154f, 155t, 159, 161t, 534t; in bays and estuaries, 126t, 134f, 136t, 137f, 145t, 303; trophic relations and, 396f, 401
- indigenous peoples, 624, 627
- Indo-Pacific fishes, 69
- Inegocia japonica*, 521t
- inland silverside (*Menidia beryllina*), 612t, 616, 619
- inner shelf (IS), 82t, 85t, 86f, 96, 109, 171, 176; analysis of, 104t; trophic relations and, 397f
- interannual variability, 140–42
- International Pacific Halibut Commission (IPHC), 525
- intertidal copepod (*Tigriopus californicus*), 38
- Isaacs, J. D., 588, 624
- Isistius brasiliensis* (cigar shark), 561, 561f
- island kelpfish (*Alloclinus holderi*), 92f, 162t, 238f, 477, 534t; cleaner behavior of, 555, 556f, 557t; feeding by, 60, 61f
- island populations. *See* California Channel Islands
- Isopsetta isolepis* (butter sole), 17t, 174, 179t, 181f, 194f, 296f
- Istophorus* spp., 523t; *I. platypterus* (sailfish), 21t, 25t, 333
- Isurus oxyrinchus* (shortfin mako), 20t, 22t, 89f, 390, 518t, 579; movement of, 527–28, 529f, 540; in pelagic zone, 326f, 331, 337t; pollution and, 601; trophic relations and, 401f, 406
- Iwama, G. K., 207
- Iwamoto, T., 374
- jack mackerel. *See* *Trachurus symmetricus*
- jacks. *See* Carangidae (jacks)
- jacksmeat. *See* *Atherinopsis californiensis*
- Jackson, J. B. C., 264, 625, 627
- Japanese sea raven (*Hemitripterus villosus*), 499, 521t
- jawfish (*Opistognathus* spp.), 485, 521t
- Jensen, Greg, 556
- Johrlandallia nigrirostris* (barberfish), 235t
- Johnson, C. E., 485
- Johnson, G. D., 62, 544
- Jones, Everett, 561
- Jones, J. A., 213, 223
- Jordan, David Starr, 253, 569, 574
- Jordania zonope* (longfin sculpin), 17t, 21t, 211, 234t, 238f, 500t
- Kathetostoma averruncus* (smooth stargazer), 20t, 23t, 179t, 185f, 197
- Katsuwonus pelamis* (skipjack tuna), 20t, 89f, 326f, 331, 337t, 573, 628
- kelp bass. *See* *Paralabrax clathratus*
- kelp bed/rocky reef (KBRF), 85t, 86f, 87, 92f, 227–52; activities of fishes in, 248–49; analysis of, 104t; assessment of abundance in, 228–32; bottom depth in, 241–43; fish assemblages and, 247–48; interannual variability in, 249–50; kelp overview, 228; latitude and, 239–41; major components of, 109f, 111; major taxa of, 232–36t; microalgae and bottom characteristics, 243–47; overfishing in, 625–26, 627; reef structure overview, 228; resource partitioning in, 454t; study localities, 82t, 84f; trophic relations in, 400f, 404–5, 625; young-of-year (YOY) and, 248
- kelp clingfish (*Rimicola muscarum*), 18t, 111f, 155t, 236t, 238f, 405
- kelpfishes. *See* Clinidae; *Gibbonsia* spp.
- kelp goby, 238f. *See also* Gobiidae
- kelp greenling. *See* *Hexagrammos decagrammus*
- kelp gunnel (*Ulvicola sanctaerosae*), 18t, 72, 72f, 238f
- kelp perch (*Brachyistius frenatus*), 19t, 111f, 162t, 422, 477, 494; adaptations of, 71f, 75; in kelp bed/rocky reef, 92f, 235, 239f, 240f; movement and activity patterns of, 533t, 544, 556f, 557t; predation and, 431f, 438–39, 440f, 442f; trophic relations and, 395, 400f, 404
- kelp pipefish (*Syngnathus californiensis*), 19t, 103, 111f, 234t, 238f, 507; in surf zone, 150t, 154f, 155t
- kelp poacher (*Agonomalus mozinoi*), 70
- kelp rockfish (*Sebastes atrovirens*), 19t, 157t, 162t, 270, 415, 536t; feeding/foraging of, 60f, 67–68; in kelp bed/rocky reef, 92f, 234t, 237f, 239f, 240f; phylogeographic structures of, 34t, 38
- kelp spp. *See* *Macrocystis* spp. (kelp)
- Kendall, Arthur, Jr., 270, 273, 306, 485
- killer whales, 429
- killifish (*Luacania parva*), 130, 131, 134f, 612t, 614. *See also* *Fundulus parvipinnis* (California killifish)
- King Harbor, 244, 249, 630
- king-of-the-salmon (*Trachipterus altivelis*), 20t, 327f, 332
- Kleppel, G. S., 305
- Kline, D. E., 371
- Kramer, S. H., 151, 541
- krill (*Euphausia pacifica*), 371
- Kroon, F. J., 531
- Kuhlia sandwichiensis*, 522t
- Kyphosidae (sea chubs), 71, 105, 106t, 235t, 534t, 557t; feeding by, 65–66. *See also* *Girella nigricans*
- Kyphosus analogus*, 20t
- Labridae (wrasses), 72, 74, 75, 106t, 235t, 393, 557t; on continental shelf, 176, 180t; feeding of, 64, 65; movement and activity patterns of, 534t, 541, 547, 548, 549; and reproduction, 496, 497, 498, 503, 507; *Thalassoma* spp., 235t, 487, 488. *See also* *Halichoeres* spp.; *Oxyjulis californica* (señorita); *Semicossyphus pulcher* (sheephead)
- Labrisomus* spp., 289, 534t, 557t; *L. xanti* (largemouth blenny), 236t. *See also* *Paraclinus integripinnis* (reef finspot)
- Lactoria diaphana*, 20t, 24t
- Lagocephalus lagocephalus* (oceanic puffer), 20t, 23t, 332
- Lamna ditropis* (salmon shark), 17t, 22t, 89f, 326f, 330, 337t, 524
- Lampadena urophaos* (sunbeam lampfish), 277, 279f, 280, 351t
- lampara nets, 571, 575, 590
- Lampetra* spp. (lamprey), 128f; *L. ayresii*, 17t; *L. tridentata* (Pacific lamprey), 18t, 130t, 133, 518t, 561–62, 561f
- lampfish. *See* *Nannobranchium* spp.; *Stenobranchius leucopsaurus*; *Triphoturus* spp.

- Lampris guttatus*, 19t, 22t, 519t
Lampris regius (opah), 89f, 321, 327f, 330, 337t
Lamprogrammus niger (paperbone cuskeel), 353t
 Lane, E. D., 135
 La Niña event, 4–5, 222, 283, 286. *See also* ENSO
 lanternfish, 351t, 370f; larval stages of, 276–80. *See* blue lanternfish; California lanternfish; *Diogenichthys* spp.; Myctophidae (lanternfish)
 Larson, Ralph J., 231, 244, 245, 531; competition studies of, 454, 455–56, 457
 larval recruitment. *See* recruitment
 larval stages. *See* ichthyoplankton assemblages
 Lasker, R., 310, 312
 lateral line, 343
 latitude, 8, 10, 110f; distribution and, 68–70, 122–23; kelp bed/rocky reef species and, 239–41; and ordination analysis, 11–14, 243
 Laur, D. R., 246, 437–38
 Laurs, R. M., 542
 Lavenberg, R. J., 348, 359, 363, 483, 539
 lavender sculpin (*Leiocottus hirundo*), 18t, 24t, 75, 76f, 532t; in kelp bed/rocky reef, 92f, 234t, 249t
 Lea, Robert N., 6, 483, 628
 LeBoeuf, B. J., 561
 Lee, W. J., 27
 Leet, W. S., 125
 left-coiling foram (*Neogloboquadrina pachyderma*), 36
 lefteye flounders. *See* Paralichthyidae
Leiocottus hirundo (lavender sculpin), 18t, 24t, 75, 76f, 532t; in kelp bed/rocky reef, 92f, 234t, 240t, 532t
Leionathus brevirostris, 521t
Leiostomus xanthurus (spot), 469, 472f
 leopard shark (*Triakis semifasciata*), 19t, 95f, 158t, 233t, 518t; in bays and estuaries, 127t, 128f, 132f; movement of, 527, 528t; in surf zone, 150t, 151, 153, 156f, 157t
Lepidocybium flavobrunneum (escolar), 20t, 22t, 327f, 330
Lepidogobius lepidus (bay goby), 18t, 98f, 155t, 158, 161t, 453t, 534t; in bays and estuaries, 126t, 128f, 132f; larval stages of, 292, 294f, 298, 300, 303, 304f
Lepidopsetta bilineata (rock sole), 17t, 22t, 605
Lepidopus caudatus, 523t
Lepidopus fitchi (Pacific scabbardfish), 7t, 19t, 23t
Lepomis macrochirus (bluegill sunfish), 487
Lepophidium microlepis (finescale cusk-eel), 177t, 183f, 196f
Lepophidium stigmatissium (Mexican cusk-eel), 177t, 182f, 195f
Leptocephalus giganteus (eel), 307, 310f
Leptocottus armatus (staghorn sculpin), 18t, 209t, 500t, 532t, 613; in bays and estuaries, 93f, 126t, 128f, 129f, 132f, 134f, 138t, 143f; on continental shelf, 178t, 181f, 194f; in surf zone, 150, 153t, 155t, 157t; trophic relations and, 396f, 402
Leptoscarus vaigiensis, 522t
Lestidiops ringens (slender barracudina), 277, 280, 351t, 355f, 370f
Lestidium pseu. sphyraenoides, 519t
Lethops connectens (goby), 18t, 23t, 418
Lethrinus miniatus, 521t
Leuresthes tenuis/sardina (grunion), 19t, 73, 145t, 149, 233t, 621; fisheries and, 567, 568, 589; genetic characteristics of, 43, 44f, 45t, 46t, 623; larval forms of, 303; movement of, 532t, 543, 549; reproduction and, 494, 495f, 520t; respiration in, 211; in surf zone, 95f, 150t, 151f
Leuroglossus stilbius (California smooth-tongue), 349t, 354, 355f, 360t, 368, 370f; larval stages of, 275, 276t, 277t, 278f, 292; swimbladder of, 365, 366t; trophic relations of, 402f, 406
 LIDAR (light detection and ranging) systems, 527–28
 Light, T., 618
 lightfishes (Phosichthyidae), 350t
 Limbaugh, Conrad, 227, 243, 555
 limpet (*Collisella limatula*), 220
 lingcod. *See* *Ophiodon elongata*
 Link, J. S., 197
Limphryne coronata, 520t
Liopropoma spp., 307, 308f
Liparis spp. (snailfishes), 175, 178t, 180t, 353t; *L. florum* (tidepool snailfish), 17t, 90f, 215t, 216f, 238f; *L. fuscus*, 17t, 521t; *L. mucosus* (slimy snailfish), 18t, 151, 153t; *L. pulchellus* (showy snailfish), 17t, 178t, 194f; *L. rutteri*, 17t; reproduction and, 485, 486
Lipolagus ochotensis (popeye blacksmelt), 349t
 lithodid crabs, 486
Lithognathus mormyrus, 521t
 lizardfishes. *See* Synodontidae
Lobianchia gemellarii (Cocco's lanternfish), 351t
Lobotes pacificus (Pacific tripletail), 7t, 20t, 25t
Lobotes surinamensis, 521t
 locomotion, 321–23. *See also* movement
Loligo opalescens (market squid), 406
 Lomolino, M. V., 621
 Long, J. A., 56
 longfin dragonfish (*Tactostoma macropus*), 351t, 359, 370f
 longfin lanternfish (*Diogenichthys atlanticus*), 351t, 370f; larval stages of, 276t, 277f, 279, 282f, 290f, 368
 longfin pearleye (*Benthalbella linguidens*), 351t
 longfin sanddab. *See* *Citharichthys xanthostigma*
 longfin sculpin (*Jordania zonope*), 17t, 21t, 211, 234t, 238f, 500t
 longfin smelt (*Spirinchus thaleichthys*), 17t, 126t, 128–29, 128f, 130t, 138t
 Longhurst, A. R., 588
 longjaw bigscale (*Scopeloberyx robustus*), 352t, 365, 366t
 longjaw mudsucker (*Gillichthys mirabilis*), 11, 19t, 88, 211, 533t, 617; in bays and estuaries, 93f, 126t, 129f, 132f, 134f, 136t, 138t, 143f; hypoxia and, 472; larval stages of, 303, 304f; phylogenetic relations and, 35t, 43, 44f, 45t, 46t; trophic relations and, 396f, 402
 longnose butterflyfish (*Forcipiger flavissimus*), 63
 longnose cat shark (*Apristurus kampae*), 353t, 372f
 longnose lancetfish (*Alepisaurus ferox*), 327f, 330, 519t
 longnose puffer (*Sphoeroides lobatus*), 7t, 20t, 25t
 longnose skate (*Raja rhina*), 18t, 101f, 177t, 184, 185f, 197f
 longspine combfish. *See* *Zaniolepis latipinnis*
 longspine thornyhead (*Sebastolobus altivelis*), 19t, 23t, 96, 101f, 311f, 334f; in bathypelagic zone, 353t, 369, 371, 372f; movement of, 537t, 540, 541
 loosetooth parrotfish (*Nicholsina denticulata*), 7t, 14, 20t, 25t, 235t
Lophiocharon trisignatus, 520t
Lophiodes spilurus, 308f
Lophiomus setigerus, 520t
Lophotus spp.: *L. capellei*, 20t, 24t; *L. cristatus* (highbrow crestfish), 327f, 332; *L. lacepede*, 519t
 Los Angeles-Long Beach harbor, 158–62
 Los Angeles Region (LAR), 40
 Losey, G. S., Jr., 489, 493, 555
 Los Penasquitos Lagoon, 144f
 lottery hypothesis of recruitment, 414–15
 louvar, 326f, 337t
 Love, Milton S., 263, 579–80, 591, 631
 lowcrest hatchetfish (*Argyropelecus sladeni*), 350t, 354, 356f, 366t; trophic relations of, 402f
 Lowe, Christopher G., 524, 539, 540
Loweina laurae, 370f
Loweina rara, 308f
 low relief substratum, 539–40
Luacania parva (rainwater killifish), 130, 131, 134f, 612t, 614
Lumpenus sagitta, 17t
 lumptail sea robin (*Prionotus stephenophrys*), 20t, 22t, 178t, 181f, 182f, 183f; foraging guild, 194f, 195f, 196f
Lutjanus spp., 309f, 521t
Luvatus imperialis, 20t, 523t
Lycenchelys crotalinus (snakehead eelpout), 17t, 22t, 354t
Lycodapus fierasfer (black mouth eelpout), 17t, 22t, 354t
Lycodapus mandibularis (pallid eelpout), 17t, 21t, 352t
Lycodes spp. (eelpouts): *L. cortezianus* (bigfin eelpout), 18t, 22t, 179t, 184, 185f, 197f; *L. diapterus* (black eelpout), 17t, 22t, 354t; *L. pacificus* (blackbelly eelpout), 18t, 98f, 179t, 183f, 192f, 196f, 538t
Lycodopsis pacificus, 522t
Lyconectes aleutensis, 17t, 21t
Lyconema barbatum (bearded eelpout), 18t, 22t, 179t, 183f, 185f; foraging guild, 196f, 197f
 Lynn, R. J., 540, 542
Lyopsetta exilis (slender sole), 18t, 403, 535t; on continental shelf, 98f, 173, 179t, 183f, 185f; foraging guild, 192f, 196f, 197f; larval stages of, 279, 296f; pollutant concentrations in, 601, 602f
Lysmata californica (cleaner shrimp), 555
Lythrypnus dalli (bluebanded goby), 19t, 23t, 61f, 92f, 236t; activity patterns of, 534t, 545; in kelp bed/rocky reef, 236t, 238f, 247; phylogenetic characteristics of, 43, 44f, 45t, 46t, 49, 623; predation and, 431–32f, 434–37, 438, 439–42, 460–61; recruitment of, 419–20, 422; reproduction and, 487; symbiotic relations of, 555, 556f, 557t
Lythrypnus spp. (gobies), 295f, 503, 523t; reproduction and, 487, 497
Lythrypnus zebra (zebra goby), 19t, 23t, 92t, 487; activity patterns of, 534t, 545; in kelp bed/rocky reef, 236t, 238f
 MacArthur, R. H., 46
 MacGinitie, G. E. and N. MacGinitie, 559
 machete (*Elops affinis*), 7t, 20t, 518
 mackerel. *See* *Trachurus symmetricus*
 macroalgae, 437–38, 474. *See also* algae
Macrocystis pyrifera (giant kelp), 48, 67, 71f, 415, 421, 459, 477; predation and, 436, 438–39
Macrocystis spp. (kelp), 228, 241, 243–46, 404–5, 628f. *See also* kelp bed/rocky reef (KBRE)
 macrophyte substratum, 111–12, 111f
Macropinna microstoma (barreleye), 349t
Macroramphosus gracilis (slender snipefish), 20t, 25t, 332, 520t
 Macrouridae (grenadiers), 101f, 106t, 579; bathypelagic, 353t, 369, 372–77, 390; on continental shelf, 175, 177t, 180t. *See also* *Coryphaenoides* spp.; *Nezumia* spp.

- Magnuson Act (1976), 578–79, 588
 Magnuson-Stevens Act (1996), 588, 591
Makaira spp. (marlin), 542, 576; *M. indica* (black marlin), 20t, 25t, 323, 333; *M. mazana*, 20t, 25t; *M. nigricans* (blue marlin), 332; *Tetrapterus audax* (striped marlin), 21t, 24t, 89f, 326f, 332, 337t, 395, 401f
Malacocephalus laevis (softhead grenadier), 369
Malacotenus hubbsi, 522t
 Malibu Creek Estuary, 129f
 Man-of-War fish (*Nomeus gronovii*), 324, 328f, 333
 Mansueti, R. J., 558
Manta spp., 337t; *M. birostris*, 20t, 25t, 333, 518t
 Mantua, N. J., 15
 Margulies, D., 274
 Marine Life Management Act (1998), 589, 591
 Marine Life Protection Act (1999), 589, 591
 mark and recapture, 525–26; and tagging, 527, 539
 marlins. *See Makaira* spp. (marlin)
 Marshall, N. J., 344, 367
 Martell, S. J. D., 229
 masked prickleback (*Ernogrammus walkeri*), 7t, 17t, 24t
 Matarese, A. C., 306
 Matern, S. A., 131, 140, 141–42, 618
 mating systems, 489, 495–98, 506. *See also* reproduction
 matrotrophy, 484, 499–500
 Matsui, T., 373
 Matthews, K. R., 527, 539, 542
 McEvoy, A. F., 587
 McFarlane, G. A., 306
 McGowen, J. S., 291, 292, 294, 295, 298
 McLusky, D. S., 119
 Mearns, A. J., 628
Medialuna californiensis (halfmoon), 19t, 35t, 92f, 157t, 162t, 400f; activity patterns of, 534t, 544; as herbivore, 65, 66f, 74, 394, 404; in kelp bed/rocky reef, 235t, 237f, 239f, 241f; symbiotic relations and, 556f, 557t, 558
 medusafish (*Ichthyos lockingtoni*), 18t, 22t, 328f, 330, 559; larval stages of, 276, 277–78f, 279, 284, 286f
 Meffert, D. J., 135
Megachasma pelagios (megamouth shark), 327f, 528, 529f
Megalops atlanticus, 518t
Melamphaes spp. (bigscale), 279f, 280, 352t, 359; *M. acanthomus* (slender bigscale), 352t; *M. lugubris* (highsnout bigscale), 277, 308f, 352t, 358f; *M. parvus* (little bigscale), 277, 352t
Melanocetus johnsonii, 520t
Melanonus zugmayeri, 519t
Melanostigma pammelas (eelpout), 18t, 22t, 348, 352t
Melichthys niger, 20t, 25t
Menidia beryllina (inland silverside), 612t, 616, 619
 Mensinger, A. F., 358
Menticirrhus undulatus (corbina), 19t, 95f, 136t, 149, 293, 536t; commercial fishing of, 587; in harbors, 157t, 158t, 160–61t; larval stages of, 293, 295f; pollutants and, 600; in surf zone, 150t, 151
Mentodus facilis (tubeshoulder), 350t
 mercury pollution, 598, 601
Merluccius spp. (hakes), 106t, 177t, 180t, 621; *M. angustimanus* (Panama hake), 177t, 197f
Merluccius productus (Pacific hake), 19t, 305, 403, 519t, 624; and commercial fishing, 578–79; on continental shelf, 177t, 182f, 183f, 185f; as deep sea species, 336t, 337t, 353t; foraging guild, 195f, 196f, 197f; habitat for, 101f, 102, 105t; larval stages of, 275, 276t, 277f, 278f, 305
 mesopelagic fishes, 343, 345t, 346, 348, 359, 364, 368, 406
 mesopelagic zone, 320, 321f, 342, 343, 348
Metridium farcimen (anemones), 220, 606
 Mexican barracuda (*Sphyræna ensis*), 7t, 20t, 25t, 236t
 Mexican cusk-eel (*Lepophidium stigmatismum*), 177t, 182f, 195f
 Mexican flashlight fish (*Phthanopanerion harveyi*), 520t, 558f
 Mexican goby (*Bathygobius ramosus*), 211
 Mexican hogfish (*Bodianus diplotaenia*), 235t
 Mexican lampfish. *See Triphoturus mexicanus*
 Mexican lookdown (*Selene berevoortii*), 7t, 20t, 308f
 Mexican Province, 3, 4f. *See also* Baja California
 Mexican rockfish (*Sebastes macdonaldi*), 19t, 24t, 44f, 45t, 46t, 258t
Microgadus proximus (Pacific tomcod), 17t, 128f, 153t, 296f; on continental shelf, 174, 177t, 181t, 182f, 183f; foraging guild, 194f, 195f
 microhabitats, in rocky intertidal zone, 208
Micrometrus aurora (reef perch), 18t, 90f, 235t; feeding of, 394, 399f, 403; in rocky intertidal zone, 213, 215t, 216f, 220f
Micrometrus minimus (dwarf perch), 18t, 77, 111f, 145t, 161t, 213, 235t, 499, 533t; in surf zone, 95f, 150t, 151f, 152, 154f, 155t
Microstoma spp., 306
Microstomus pacificus (Dover sole), 19t, 96, 188, 307, 523t, 535t; as bathopelagic species, 354t, 369; commercial fisheries and, 574, 576; on continental shelf, 172, 179t, 182f, 183f, 185f; fin erosion disease, 603–4; foraging guild, 192f, 195f, 196f, 197f; habitat of, 101f, 103, 105t; in pelagic zone, 333, 334t; phylogenetics of, 33, 34t; pollutants and, 598, 600, 605
 mid-depth reef (MDR), 99f, 100f
 mid-depth rocky reef (MDRF), 85t, 86f, 104t
 middle shelf (MS), 85t, 86f, 104t, 109, 176, 258t
 midshipman. *See Porichthys* spp.
 midwater eelpout, 355f
 migration, 48–49, 322, 361, 530, 615; in bays and estuaries, 124, 128f, 132f, 134f; coastal pelagics, 89f, 187–88; crepuscular, 546; spatial patterns and, 541–43. *See also* movement
 milk fish (*Chanos chanos*), 519t, 612t, 618
 Miller, D. J., 6, 245–46, 253, 483, 539, 542
 mimic sanddab (*Citharichthys gordae*), 179, 183f, 193, 196f
Mirounga angustirostris (elephant seals), 530, 542, 561
 mitochondrial control region (MtCR), 27
Mobula japonica, 20t, 24t
 mojarras (Gerreidae), 175, 180t, 506; spotfin mojarras, 178t, 181f, 182f, 194f, 195f
Mola mola (ocean sunfish), 20t, 77, 89f, 523t, 540, 556; in pelagic zone, 326f, 330, 336, 337t
 molecular clocks, 28–29, 49
 monkeyface prickleback. *See Cebidichthys violaceus*
 Monocentridae (pinecone fishes), 66
 monogamy, 496–98, 503
 moray eel. *See Gymnothorax mordax*
 Moridae (codlings), 175, 177t, 180t
 Morita, T., 375
Morone saxatilis (striped bass), 18t, 88, 149–50, 390, 470, 521t; as alien species, 612t, 613f, 615, 618; in bays and estuaries, 93f, 126t, 128f, 130t, 131, 138t; fisheries and, 585t, 587; pollutants and, 598, 600, 602; reproduction of, 497; trophic relations and, 395f, 402
 Morro Bay, 131, 133
 Moser, H. G., 6, 273, 298, 305, 308, 368–69
 mosquitofish (*Gambusia affinis*), 126t, 132f, 535t, 612t, 616–17
 mosshead sculpin (*Clinocottus globiceps*), 216f, 220, 390, 500t; feeding of, 390, 392, 403
 mosshead warbonnet, 91f, 238f
 mottled sculpin (*Cottus bairdi*), 501
 movement and activity patterns, 187, 524–49; circa patterns, 548–49; diel-related, 543; fishing and, 525–26; genetic markers and, 530; home ranging and, 530–31; LIDAR and, 527–28; migration and, 541–43; ontogenetic-related, 541; in open ocean, 540; remote sensing techniques and, 527–30; in rockfish, 531, 539–40; seasonal, 541–42; in situ monitoring of, 526–27; sonar and, 527–28; spatial patterns, 530–31, 539–43; telemetry and, 527–30, 539; tide-related, 542–43. *See also* migration
 Moyle, Peter B., 611, 616, 618; on bay and estuarine species, 123, 125, 130, 131
 Moynihan, M., 561
 Mozambique tilapia (*Oreochromis mossambica*), 485, 612t, 617
 Muench, K. A., 494
Mugil cephalus (striped mullet), 20t, 157t, 158t, 520t, 534t; in bays and estuaries, 95f, 126t, 129f, 134f, 145t, 398; as brephoeipipelagic fish, 333, 334f; feeding and trophic relations, 390, 394, 396f
 Mugillidae (mulletts), 329, 470, 472f
 Mullidae (goatfishes), 7t, 135t, 329, 506
 Mulligan, T. J. and H. L. Mulligan, 151
 Mullin, M., 305
Mulloidichthys dentatus (Mexican goatfish), 235t
 mummichog (*Fundulus heteroclitus*), 139, 470, 520t
 Muehara, H., 489, 507
 mussel blenny (*Hypsoblennius jenkinsi*), 19t, 92f, 136f, 238f, 247; movement patterns of, 531, 532t; phylogenetic characteristics of, 43, 44f, 45t, 46t; territoriality of, 451, 453t
 mussel (*Mytilus californianus*), 221, 555
Mustelus californicus (gray smoothhound), 19t, 95f, 127t, 132f, 233t; in surf zone, 150t, 151, 153, 156f, 157t, 158t
Mustelus henlei (brown smoothhound), 19t, 95f, 127t, 128f, 233t; in surf zone, 153, 156f, 157t, 158t
Mustelus lunulatus, 20t
 mutualism, 554–58. *See also* symbiotic relationships
Mycteroperca spp. (grouper): *M. jordani* (gulf grouper), 19t, 25t, 234t; *M. xenarcha* (broomtail grouper), 20t, 23t, 58, 234t
 Myctophidae (lanternfish), 306, 329, 406, 557–58; deep sea species, 356, 358, 359, 368, 370f; larval stages of, 276–80; swimbladders of, 366t, 367–68
Myctophum aurolaternatum (golden lanternfish), 307, 310f, 370f

- Myers, J. M., 31
- Myliobatis californica* (bat ray), 19t, 95f, 233t, 337t, 429, 518t, 535t; in bays and estuaries, 127t, 128f, 132f, 134f, 137f; feeding of, 394; and subsistence fishing, 568; in surf zone, 150t, 153, 156f, 157t, 158t, 160t
- Myoxocephalus polyacanthocephalus*, 500t
- Myrberg, A. A., Jr., 489
- Myripristis amaena*, 520t
- Myrophis vafer*, 20t, 25t
- Mytilus californianus* (mussels), 221, 555
- Myxine circifons*, 20t, 25t
- Myxinidae (hagfishes), 103, 106t, 353t, 578–79; *Eptatretus deani* (black hagfish), 18t, 101f, 353t, 372f
- Nakano, K., 207
- nakedbelly sea robin (*Bellator gymnotetus*), 178t, 183f, 196f
- naked goby (*Gobiosoma bosc*), 469
- Nannobranchium ritteri* (broadfin lampfish), 351t, 356, 357f, 359, 368, 370f; larval stages of, 276t, 277f, 279, 280, 305; swimbladder of, 366t, 367; trophic relations of, 402f
- Nannobranchium* spp. (lampfishes), 276t, 279f, 309f, 369; *N. regale* (pinpoint lanternfish), 351t, 358, 365, 366t, 370f
- Nansenia crassa*, 519t
- Naples Reef, 455, 456, 458f, 474f, 476, 546–47
- Narcine brasiliensis*, 518t
- National Marine Fisheries Service (NMFS), 230, 336, 589; continental shelf species and, 168, 169, 170, 171, 172, 175
- Natural History Museum of Los Angeles County (LACM), 275
- Naucrates ductor* (pilotfish), 20t, 23t, 324, 328t
- Naumann, E., 306
- Nautichthys oculofasciatus* (sailfin sculpin), 17t, 91f, 234t, 238f, 500t
- Navarro River Estuary, 129f
- nearshore (NS) habitat, 94–96, 94f, 95f
- nearshore soft bottom (NSB), 95f, 96, 97f, 157t, 158t
- needlefishes (Belonidae), 73, 135t
- Neighbors, Margaret A., 6, 367
- Nelson, D. R., 528, 544
- Nelson, J. S., 55, 61, 73, 77
- Nematistius pectoralis*, 20t, 25t, 521t
- Nemichthyidae, 354
- Nemichthys scolopaceus* (slender snipe eel), 349t, 519t
- Nemipterus virgatus*, 522t
- Neocirrhites armatus*, 522t
- Neoclinus* spp. (fringeheads): *N. blanchardi*, 18t; *N. stephensae* (yellowfin fringehead), 19t, 23t, 236t, 238f, 247; *N. uninotatus*, 18t
- Neogloboquadrina pachyderma* (left-coiling foram), 36
- Neoscopelidae (blackchins), 358
- Neotrypaea biffari* (ghost shrimp), 559, 560f
- Neotrypaea californiensis* (pink ghost shrimp), 559
- Nereocystis luetkeana* (bull kelp), 228, 243, 245–46
- neuromasts, 343, 359
- Newell, N. D., 57
- Newport Bay, 135, 162, 164f
- Nezumia* spp. (grenadiers): *N. kensmithi* (bluntnose grenadier), 353t; *N. liolepsis* (smooth grenadier), 353t, 369; *N. stelgidolepis* (California grenadier), 18t, 23t, 177t, 184, 185f, 197f, 353t, 369
- niche overlap, competition and, 451, 452f, 456, 457, 458f
- Nicholsina* spp. (parrotfish), 522t; *N. denticulata* (loosetooth parrotfish), 7t, 14, 20t, 25t, 235t
- Nicol, J. A. C., 73
- Nielsen, J. L., 31
- night smelt (*Spirinchus starksi*), 17t, 72, 151, 152f, 153t, 335; on continental shelf, 177, 181f, 194f
- Nile tilapia (*Oreochromis niloticus*), 393
- nocturnality, 66–68, 94, 95f, 156f, 188, 189, 433f; movement patterns and, 543, 548
- Nomeus gronovii* (Man-of-War fish), 324, 328f, 333
- non-bay species, 8, 11f, 12f
- nonmetric multidimensional scaling (NMDS), 144f
- Norris, K. S., 414, 419
- northern anchovy. *See* *Engraulis mordax*
- northern clingfish (*Gobiesox maeandricus*), 18t, 33, 34t, 90f, 238f, 523t; in rocky intertidal zone, 209, 215t, 216f, 220f, 221; trophic relations and, 399f, 403
- northern lampfish. *See* *Stenobranchius leucoparus*
- northern pearleye (*Benthalbella denatata*), 351t, 370f
- Norton, S. F., 219–20, 223, 407
- notacanthiforms, 307
- Notorhynchus* spp.: *N. cepedianus*, 518t; *N. cepedianus* (sevengill shark), 19t, 233t; *N. valdiviae* (topside lampfish), 351t, 370f
- nuclear power plant effects, 270
- nursery, bays and estuaries as, 142–43, 303–5, 468. *See also* ichthyoplankton assemblages
- oarfish, 327f, 337t
- Occella vernicosa*, 17t
- oceanic puffer (*Lagocephalus lagocephalus*), 20t, 23t, 332
- ocean sunfish (*Mola mola*), 20t, 77, 89f, 523t, 540, 556; in pelagic zone, 326f, 330, 332, 336
- ocean surface warming, 5, 6, 242, 257. *See also* temperature of water
- ocean whitefish (*Caulolatilus princeps*), 20t, 22t, 35t, 87, 311f, 521t; in kelp bed/rocky reef, 92f, 234, 237f, 239f; and recreational fishing, 586
- Odontaspis ferox*, 19t, 25t
- Odontaspis taurus*, 518t
- Odontopyxis trispinosa* (pygmy poacher), 18t, 178t, 182f, 192f, 195f
- oilfish (*Ruvettus pretiosus*), 20t, 24t, 327f, 332
- oil spills, 595, 596
- olfaction, 343
- Oligocottus* spp. (sculpins), 500; *O. rimensis* (saddleback sculpin), 17t, 21t, 90f, 209t, 215t, 216f, 399f
- Oligocottus maculosus* (tidepool sculpin), 17t, 91f, 500t, 507, 533t; in rocky intertidal zone, 207, 209t, 215t, 216f, 217f, 218
- Oligocottus rubellio* (rosy sculpin), 17t, 23t, 90f, 234t, 238f, 399f; in rocky intertidal zone, 215t, 216f, 217f, 220f
- Oligocottus snyderi* (fluffy sculpin), 18t, 90f, 399f, 500t; in rocky intertidal zone, 209t, 215t, 216f, 217f, 218, 220f
- Oligoplites saurus*, 24t
- Oligopus diagrammus*, 20t
- olive rockfish (*Sebastes serranoides*), 18t, 60f, 91f, 162t, 219t; in kelp bed/rocky reef, 234t, 237f, 240f, 243f; movement and activity of, 537t, 548; and recreational fishing, 584, 585t, 590; trophic relations and, 400f, 405
- Oncorhynchus kisutch* (coho salmon), 18t, 30t, 31, 32, 88, 233t, 330, 331f, 487; in bays and estuaries, 126t, 128f, 130t
- Oncorhynchus mykiss* (steelhead), 18t, 30t, 31, 32, 618; in bays and estuaries, 126t, 128f, 129f, 130t, 132f
- Oncorhynchus* spp. (anadromous trout), 393; *O. clarkii* (cutthroat trout), 17t, 21t, 29, 130t; *O. gorbusha* (pink salmon), 17t, 22t, 130t; *O. keta* (chum salmon), 17t, 21t, 130t, 233t; *O. nerka* (sockeye salmon), 17t, 21t, 519t, 624–25, 626f
- Oncorhynchus tshawytscha* (chinook salmon), 17t, 153t, 233t, 330, 331f, 630; in bays and estuaries, 93f, 126t, 128f, 130t, 131; fisheries and, 569, 579, 585t, 589, 628; phylogeography and, 30t, 31–32, 32t, 39
- Oneirodes* spp. (dreamers), 17t, 352t, 395, 402t; *O. acanthias* (spiny dreamer), 352t, 358, 390, 520t; *O. eschrichtii* (bulbous dreamer), 352t
- onespot fringehead, 97f
- Onuf, C. P., 140, 141
- oophagy, 484
- opah (*Lampris regius*), 89f, 321, 327f, 330, 337t
- opaleye (*Girella nigricans*), 19t, 102, 105t, 136t, 150t, 534, 585t; activity patterns of, 544–45; in harbors, 157t, 158t, 162t; as herbivore, 65, 66f, 74, 394, 400f, 403, 404; in kelp bed/rocky reef, 92f, 235t, 237f, 239f, 241f; mutualism of, 556f, 557t, 558; phylogenetic characteristics of, 35t, 41, 43, 44f, 45t, 46t; recruitment of, 414, 419; in rocky intertidal zone, 209t, 215t, 216f
- open ocean pelagics. *See* pelagic zone
- Ophichthus* spp.: *O. rufus*, 518t; *O. triserialis*, 20t; *O. zophochir*, 20t
- Ophiclinus* spp., 522t
- Ophidiidae (cusk-eels), 106, 177t, 180t, 197f, 353t, 369; *Lepophidium* spp., 177t, 182f, 183f, 195f, 196f; movement and activity of, 535t, 544. *See also* *Chilara taylori*
- Ophidion barbatum*, 519t
- Ophidion scrippsae* (basketweave cusk-eel), 18t, 163, 177t, 535t, 603; foraging guild, 194f, 195f; larval stages of, 277t, 280f, 293, 295f, 297f, 298
- Ophioblennius steindachneri* (Panamic fanged blenny), 236t
- Ophiodon elongatus* (lingcod), 18t, 100f, 103, 105t, 129f, 132f, 485; as ambusher, 59f; as brephoepelagic fish, 333, 334f; climate change and, 624, 631; on continental shelf, 178t, 181f, 182f, 183f; in deep rock habitat, 262f, 263t; fisheries and, 574, 576, 579, 589, 591; foraging guild, 194f, 195f; in kelp bed/rocky reef, 234t, 237f, 240f; larval stages of, 270, 295, 297f; and subsistence fishing, 568
- Ophiophthalmus normani* (brittle star), 371, 540
- Opisthonema libertate*, 519t
- Opisthoproctidae (spookfishes), 349t, 519t
- Opistognathus* spp. (jawfish), 485, 521t
- orange roughy (*Hoplostethus atlanticus*), 66, 520t
- orangethroat pikeblenny (*Chaenopsis alepidota*), 19t, 24t, 137f; phylogenetic characteristics of, 43, 44f, 45t, 46t
- Orbach, M. K., 590
- Oregonian Province, 3, 4f, 239
- Oreochromis mossambicus* (Mozambique tilapia), 485, 612t, 617

- Oreochromis niloticus* (Nile tilapia), 393
- Orthodon microlepidotus* (Sacramento blackfish), 393
- Orthonopias triacis* (snubnose sculpin), 18t, 23t, 60, 91f, 234t, 238t, 533t
- Orthopristis reddingi* (bronzestripe grunt), 178t, 181f, 182f, 194f, 195f
- Osborn, T. R., 309
- Osmeridae (true smelts), 72, 73, 77, 120, 128–29, 149; on continental shelf and slope, 175, 177t, 180t; larval stages of, 296, 304f
- osmoregulation, 212
- Ostracion meleagris*, 523t
- otter (*Enhydra lutris*), 624, 627, 628
- otter trawl, 575
- outer shelf (OS), 85t, 86f, 109, 176, 179–80; analysis of, 104t; trophic relations and, 398f
- oval flounder (*Syacium ovale*), 179t, 194f
- overfishing, 567, 572, 573, 576, 580, 588; climate change and, 625–26, 627, 633. *See also* fishing and fisheries
- oviparity (egg laying), 483–84, 486–87, 499; egg size and, 500–501
- oxygen depletion. *See* hypoxia
- Oxyjulis californica* (señorita), 18t, 58, 105t, 136t, 259, 522t; activity patterns of, 534t, 544, 546, 548; cleaner behavior of, 555, 556f, 557t; feeding of, 65, 389, 390, 400f; in harbors, 159, 162t; in kelp bed/rocky reef, 87, 92f, 235t, 237f, 239f, 240f, 241f, 404–5; larval stages of, 293, 295f; macrophyte substratum and, 111; predation and, 431f, 436; reproduction and, 497, 498
- Oxylebius pictus* (painted greenling), 18t, 162t, 295, 420, 521t, 538t; in deep rock habitat, 258, 263t; in kelp bed/rocky reef, 70, 71f, 91f, 234t, 237f, 238f, 404; phylogeography and, 33, 34t, 38, 38f; reproduction and, 489, 490, 492f, 495, 496, 501; symbiotic relations of, 560–61, 561f; trophic relations and, 400f
- Oxymonacanthus longirostris*, 523t
- Pacific argentine (*Argentina sialis*), 19t, 22t, 100f, 177t, 182f, 195f, 519t
- Pacific baracuda. *See* *Sphyrna argentea*
- Pacific blackchin (*Scopelogadus trisus*), 351t, 357f, 358, 366t
- Pacific blackdragon (*Idianthanthus antrastomus*), 307, 310f, 351t, 356, 357f, 366t, 370f
- Pacific blacksmelt (*Bathylagus pacificus*), 310, 349t, 354, 355f, 370f; swimbladder of, 365, 366t
- Pacific bonito. *See* *Sarda chiliensis*
- Pacific butterflyfish, 337t
- Pacific cod (*Gadus macrocephalus*), 17t, 21t, 605
- Pacific crevalle jack (*Caranx caninus*), 7t, 20t, 25t
- Pacific Decadal Oscillations. *See* PDO
- Pacific dreamer. *See* spiny dreamer (*Oneirodes acanthias*)
- Pacific electric ray (*Torpedo californica*), 18t, 95f, 156f, 233t, 348, 405, 518; on continental shelf, 177t, 183f, 196f; hunting behavior of, 433f; movement of, 538t, 540, 542, 549; in pelagic zone, 335, 337t
- Pacific Fisheries Management Council (PFMC), 588, 589
- Pacific flatnose (*Antimora microlepis*), 18t, 21t, 353t, 372f
- Pacific grenadier (*Coryphaenoides acrolepis*), 18t, 22t, 353t, 369, 372f, 373, 374
- Pacific hake. *See* *Merluccius productus*
- Pacific halibut (*Hippoglossus stenolepis*), 17t, 525, 580
- Pacific herring. *See* *Clupea pallasii*
- Pacific lamprey (*Lampetra tridentata*), 18t, 518t, 561–62, 561f
- Pacific lightfish (*Vinciguerria lucetia*), 305, 350t, 368, 370f; larval stages of, 276t, 277f, 279f, 280, 282f, 286, 292f
- Pacific line sole (*Achirus mazatlanus*), 137f, 179t, 181f, 194f, 307, 523t
- Pacific mackerel. *See* *Scomber japonicus*
- Pacific pomfret, 326f, 337t. *See also* *Taractes longipinnis*
- Pacific pompano (*Peprilus simillimus*), 19t, 91, 94f, 103, 305; feeding of, 393; in harbors, 159f, 160–61t; in pelagic zone, 156, 330, 332f
- Pacific porgy (*Calamus brachysomus*), 20, 137f, 235t
- Pacific sanddab. *See* *Citharichthys sordidus*
- Pacific sand lance (*Ammodytes hexapterus*), 17t, 75, 153t, 296f, 494–95, 522t
- Pacific sand perch (*Diplectrum pacificum*), 178t, 181f, 194f, 521t
- Pacific sardine. *See* *Sardinops sagax*
- Pacific saury (*Cololabis saira*), 326f, 330, 335, 336t, 401f, 405
- Pacific scabbardfish (*Lepidopus fitchi*), 7t, 19t, 23t
- Pacific seahorse (*Hippocampus ingens*), 20t, 25t, 135t, 137f, 484
- Pacific sleeper shark (*Somniosus pacificus*), 17t, 21t, 353t, 518t
- Pacific tomcod (*Microgadus proximus*), 17t, 128f, 153t, 181t, 296f; on continental shelf, 174, 177t, 181t, 182f, 183f; foraging guild, 194f, 195f
- Pacific Transition Conception, 40, 240
- Pacific tripletail (*Lobotes pacificus*), 7t, 20t, 25t
- Pacific viperfish (*Chauliodus macouni*), 351t, 356, 357f, 365, 366t, 370f; trophic relations, 395, 402f, 406
- Paine, R. T., 442, 554
- painted greenling. *See* *Oxylebius pictus*
- pallid eelpout (*Lycodapus mandibularis*), 17t, 21t, 352t
- Pallisina barbata* (tubenose poacher), 17t, 153t
- Palos Verdes, 244, 249, 630
- Palos Verdes Shelf: fin erosion disease in, 603, 604f, 607f; pollution in, 596, 598, 600, 601, 607–8
- Panama hake (*Merluccius angustimanus*), 177t, 197f
- Panama lightfish. *See* Pacific lightfish (*Vinciguerria lucetia*)
- Panama snaggletooth (*Borostomias panamensis*), 351t, 357f, 366t
- Panamic fanged blennie (*Ophioblennius steindachneri*), 236t
- Panamic Province, 3, 4f
- Panamic sergeant major (*Abudefduf troschelli*), 92f, 235t, 237f, 241f
- Pangea, breakup of, 621, 622f
- paperbone cusk-eel (*Lamprogrammus niger*), 353t
- Paraclinus integripinnis* (reef finspot), 19t, 92f, 236t, 403, 534t, 628; in bays and estuaries, 136t, 137f; hypoxia and, 468; in rocky intertidal zone, 211, 215t, 216f, 222f
- Paralabrax* spp. (sand bass), 395, 506; larval stages, 293, 295f, 298, 300, 301–2; *P. auroguttatus* (golden spotted rock bass), 234t, 497; *P. humeralis*, 497
- Paralabrax clathratus* (kelp bass), 19t, 35t, 58, 111f, 260, 506, 521t; in bays and estuaries, 136t, 137f, 145t; commercial fishing of, 576, 591; ENSO and, 631; feeding of, 395; growth of, 62; habitat range of, 92f, 102, 103, 105t; in harbors, 157t, 159, 162t; in kelp bed/rocky reef, 234t, 237f, 239f, 241f; movement and activity patterns of, 525, 526f, 531, 537t, 539, 545, 549; pollution and, 600, 605; predation and, 430, 431f, 437, 438–39, 441, 444; recreational fishing of, 581, 584, 585t, 586; recruitment of, 414, 418, 419, 421; reproduction and, 497; storm disturbance and, 477, 478f; subsistence fishing and, 568; in surf zone, 152, 154f, 155t; symbiotic relationships of, 555, 556f, 557t; trophic relations and, 395, 400f, 405
- Paralabrax maculatofasciatus* (spotted sand bass), 19t, 111f, 506, 521t, 538t, 591, 631; in bays and estuaries, 95f, 126t, 134f, 136t, 137f, 145t; on continental shelf, 178t, 181f, 194f; phylogenetic characteristics of, 43, 44f, 45t, 46t; trophic relations and, 395, 396f, 402
- Paralabrax nebulifer* (barred sand bass), 19t, 92f, 102, 103, 105t, 111f, 506, 631; in bays and estuaries, 127t, 136t, 137f, 145t; on continental shelf, 178t, 181f, 194f; fisheries and, 576, 582, 584, 585t, 591; in harbors, 157t, 160t, 162t; in kelp bed/rocky reef, 234t, 237f, 239f, 241f; movement of, 531, 538t; predation and, 431; reproduction and, 497; trophic relations and, 395, 397f
- Paralepididae (barracudinas), 279f, 351t
- Paralichthyidae (lefteye flounders), 70, 71, 106t, 236t, 289, 301, 535t; on continental shelf, 172, 176, 179t, 180t, 184. *See also* *Citharichthys* spp.
- Paralichthys californicus* (California halibut), 19t, 34t, 95f, 160–61t, 236t, 305; ambush feeding of, 59f, 60, 395, 432; in bays and estuaries, 127t, 129f, 132f, 136t, 137f, 138t, 145t; commercial fisheries and, 573–74, 576, 579, 587, 591; on continental shelf, 172, 179t, 181f, 194f; habitat of, 102, 105t; larval stages of, 291, 294, 297f, 298, 299f; movement of, 535t, 539, 541; pollution and, 600, 606; recreational fishing and, 582, 583, 585t; recruitment of, 187, 420; in surf zone, 151–52, 153t, 154f, 155t, 157t, 158t; trophic relationships and, 396f, 397f, 402, 403
- Paraliparis cephalus* (swellhead snailfish), 353t
- Paramonacanthus japonicus*, 523t
- Paranthias colonus* (Pacific creolefish), 234t, 309f
- Parapercis snyderi*, 522t
- parasitism, 486, 554, 555, 561–62, 605
- parental care, 484–85, 495, 499–500, 502–4, 506; egg size and, 500, 501t; viviparity and, 484, 502, 503, 504, 507
- Paricelinus hoplitis*, 18t, 22t
- Parin, N. V., 324, 325, 329, 331, 333
- Parmaturus xanurus* (fletail cat shark), 19t, 23t, 101f, 353t, 518t
- Parulirus interruptus* (spiny lobster), 555
- Parophrys vetulus*. *See* *Pleuronectes/Parophrys vetulus*
- Parques viola* (rock croaker), 235t
- Parrish, J. K., 333, 573
- Parrish, R. H., 76
- parrotfishes (Scaridae), 394, 506; *Nicholsina* spp., 7t, 14, 20t, 25t, 235t, 522t

- Parvilux ingens* (giant lampfish), 351t, 366t, 370f
 patchwork lanternfish, 370f
 Patterson, C., 66
 Patton, M. L., 240, 244
 Paxton, J. R., 363
 PCBs and DDT contamination, 595–603, 605, 607f. *See also* pollution
 PDO (Pacific Decadal Oscillations), 5, 15, 628–31; ichthyoplankton assemblages and, 283, 284, 285–93f, 299–302f, 303
 peacock wrasse (*Symphodus tinca*), 502
 Pearcy, William G., 270, 295, 296, 298, 361, 539
 pearleyes (*Benthalbella* spp.), 351t, 370f, 519t
 Pearson product-moment correlation matrix, 6
 pebbled butterflyfish (*Chaetodon multicinctus*), 63
Pelagia spp. (sea jellies), 328f
 pelagic egg, 77. *See also* oviparity
 pelagic larval duration, 33, 49
 pelagic (PEL) zone, 82t, 85t, 86t, 89f, 104t, 106, 320–41; adaptations to, 321–24; boreal region, 330; brephoepipelagic fish, 325, 328–29, 333, 334; California Current system and, 333–36; epheroepipelagic fish, 325, 328, 331f; holoeipelagic fish, 324–25, 326–28f, 330t, 331; locomotor adaptations, 321–23; meroepipelagic fish, 325, 328t; nyctoepipelagic fishes, 325, 329; prominent world-wide groups in, 320–21; schooling, 323–24; temperate region, 330; tropical region, 331–33; xenoepipelagic fishes, 325, 329, 330t, 332f
 pelagic stingray (*Dasyatis violacea*), 326f, 331, 337t
Pelagophycus porra (elk kelp), 228
Pempheris vanicolensis, 522t
 penpoint gunnel (*Apodichthys flavidus*), 17t, 90f, 111f, 129f, 151, 153t, 485; in rocky intertidal zone, 209t, 215t, 216f, 221
Pepilus simillimus (Pacific pompano), 19t, 91, 94f, 103, 305; feeding of, 393; in harbors, 159f, 160–61t; in pelagic zone, 156, 330, 332f
 Pequegnat, W. E., 227, 229
 perciforms, 71–72, 75, 105, 178, 306
 Percoidae, 57–58, 321
Perissias taeniopterus (flag flounder), 179t, 183f, 193, 196f
 Perkins, H. C., 359, 361
 Pescadero Creek Lagoon, 129f
 Peterson, C. W., 499
 petrale sole (*Eopsetta jordani*), 18t, 179t, 182f, 183f, 185f, 193; foraging guild, 193, 195f, 196f, 197f
Petromyzontidae, 561–62. *See also* *Lampetra* spp. (lamprey)
 Pfister, C. A., 214, 223, 421
Phanerodon atripes (sharpnose seaperch), 18t, 23t, 75, 235t, 555; cleaner behavior of, 556f, 557t
Phanerodon furcatus (white seaperch), 18t, 75, 150t, 155t, 258, 631; cleaner behavior of, 556f, 557t; on continental shelf, 178t, 181f; foraging guilds, 192f, 194f; habitat range of, 92f, 102, 103, 105t, 127t, 132f; in harbors, 157t, 160–61t, 162t; in kelp bed/rocky reef, 235t, 240f; movement of, 533t, 542; reproduction of, 494; trophic relations and, 397f, 403
 pheromones, 491, 493, 494
Pholis spp. (gunnels), 9f, 10f, 72, 90f, 209t, 217f; *P. clemensi*, 17t, 21t; *P. laeta* (crescent gunnel), 17t, 21t, 209t, 485, 522t; *P. ornata* (saddleback gunnel), 17t, 153t, 399f; *P. schultzi*, 17t, 21t; *Ulvicola sanctaerosae* (kelp gunnel), 18t, 24t, 72, 72f, 236t. *See also* penpoint gunnel; rockweed gunnel
 Phosichthyidae (lightfishes), 350t
Photonectes margarita, 351t
Phragmatopoma californica (tube worm), 228
Phthanophaneron harveyi (Mexican flashlight fish), 520t, 558f
Phtheichthys lineatus (slender sucker), 20t, 25t, 324
Phyllospadix spp. (surf grass), 228
 phylogeny, 68, 195
 phylogeography, 26–50; anadromous fishes and, 29–32; Baja California, 40–41, 44–46t; biogeography and, 9, 26; Channel Islands and, 33, 43, 47–49; characteristics of coastal fishes, 34–35t; climate change and, 36–38; comparative, 26–27; gene flow and, 27, 33, 38, 48, 49–50; glaciations and, 32–33, 36, 37, 41; Gulf of California, 41–43; pelagic larval duration and, 33; subregions, 27; tectonism and oceanographic patterns, 33, 36; zoographic barriers and, 26–27
Physalia spp. (siphonophores), 333
Physiculus rastrelliger (hundred-fathom codling), 20t, 23t, 98f, 177t, 185f, 197f; larval stages of, 309f
Phytichthys chirus, 17t, 21t
 Pickett, S. T. A., 466
 Pihl, L., 143
 pile perch. *See* *Rhacochilus vacca*
 pilotfish (*Naucrates ductor*), 20t, 23t, 324, 328f
 pinecone fishes (Monocentridae), 66
 pinfish (*Lagodon rhomboids*), 469, 472f
 pink cardinalfish (*Apogon pacificus*), 7t, 20t, 25t, 234t
 pinknose rockfish (*Sebastes simulator*), 18t, 24t, 100f, 255, 262f
 pink rockfish (*Sebastes eos*), 18t, 23t, 100f, 258t
 pink salmon (*Oncorhynchus gorbuscha*), 17t, 22t, 130t
 pink seaperch (*Zalembius rosaceus*), 19t, 23t, 105t, 258, 263t, 533t; on continental shelf, 98f, 176, 178t, 182f, 183f; foraging guild, 192f, 195f, 196f; trophic relations and, 398f, 403
 pinpoint lanternfish (*Nannobranchium regale*), 351t, 358, 365, 366t, 370f
 pipefishes. *See* *Syngnathus* spp.
 pipeline assemblages, 264
 piscivores, 131, 330, 395, 402, 430, 435, 548; filial cannibalism, 436, 500–502. *See also* predators
 Pittenger, G. G., 540
Plagiogrammus hopkinsii (crisscross prickly-back), 17t, 24t, 91f, 236t, 238f
Plagiotremus azaleus (sabertooth blenny), 7t, 20t, 25t, 236t
 plaice (*Pleuronectes platessa*), 539, 604
 plainfin midshipman. *See* *Porichthys notatus*
 planktivores, 63, 75, 131, 189, 420, 547
 planktonic larval duration, 26, 27f
Platichthys stellatus (starry flounder), 17t, 296f, 600, 605, 606; in bays and estuaries, 93f, 127f, 128f, 129f, 132f, 138t
Platycephalus speculator, 521t
 Platyrrhinidae (thornbacks), 175, 177t, 180t
Platyrrhinoides triseriatus (thornback), 18t, 97f, 157t, 158t, 233t; in pelagic zone, 177t, 181f, 194f
 Platyroctidae (tubeshoulders), 350t, 354, 355f, 369, 519t
Plecoglossus altivelis (ayu), 612t, 618
Plectobranchius evides, 17t, 22t
 Pleistocene epoch, 623, 624
Plesiops nigricans, 521t
Pleurogrammus monopterygius, 17t, 21t
Pleuronectes/Parophrys vetulus (English sole), 19t, 98f, 105t, 296f, 398f, 535t; in bays and estuaries, 127t, 128f, 132f; on continental shelf, 172, 179t, 181f, 182f; foraging guilds, 192f, 194f, 195f; pollutant-related disease in, 600, 604, 605, 606; in surf zone, 152f, 153t, 155t
Pleuronectes platessa (plaice), 539, 604
 Pleuronectidae (righteye flounders), 9f, 10f, 70–71, 105, 106t, 535t; pelagic, 329, 333
 pleuronectiforms (flatfishes), 59, 96, 307, 539; commercial fishing of, 569, 574, 575, 579; on continental shelf, 172, 179t, 180t, 184, 186, 189; larval stages of, 289, 306; movement of, 541; pollutants and, 600, 606
Pleuronichthys coenosus (C-O sole), 18t, 60, 61f, 70, 236t, 535t
Pleuronichthys decurrens (curlfin sole/turbot), 18t, 98f, 398f, 535t, 606; on continental shelf, 179t, 181f; foraging guild, 192f, 194f
Pleuronichthys guttulatus. *See* *Hypsopsetta guttulata*
Pleuronichthys ritteri (spotted turbot), 19f, 95f, 151, 155t, 157t; on continental shelf, 179t, 181f, 182f; foraging guild, 194f, 195f; larval stages of, 292, 293, 294f, 300; pollution and, 600, 603; trophic relations and, 397f
Pleuronichthys verticalis (hornyhead turbot), 19t, 98f, 294f, 398f, 535t, 606; on continental shelf, 179t, 181f, 182f; foraging guild, 192f, 193, 194f, 195f; larval stages of, 291, 299; trophic relations and, 397f, 398f
 Pliocene epoch, 622f, 623
Plotosus lineatus, 519t
 poachers, 18t, 70, 98f, 153t, 178t; bigeye poacher (*Bathyagonus pentacanthus*), 178t, 197f; pricklebreast poacher (*Stellerina xyosterna*), 18t, 152f, 153t, 296; pygmy poacher (*Odontopyxis trispinosa*), 18t, 178t, 182f, 192f, 195f; tubenose poacher (*Pallasina barbata*), 17t, 153t. *See also* Agonidae; *Xeneretmus* spp.
Podothecus sachi, 521t
Poecilia latipinna (sailfin molly), 612t, 613f, 616
 Poeciliidae spp., 494, 616–17
 Point Conception, 3, 5, 6, 75, 76
 Polivka, K. M., 221
 pollution, 121, 145, 595–608; bioaccumulation in fish and, 598–601; biomarkers and sublethal effects, 605–6; contamination effects, 601–8; DDT and PCBs, 595–603, 605, 607f; effects on populations and assemblages, 606–8; external disease, anomalies and parasites, 603–5; fin erosion disease and, 603, 604f; human health risk, 601–2; sediment contamination, 596, 598, 600f, 604, 607f, 608; sources of contaminants, 595–96; in water column, 596; wild-life health risk and, 602–3
 polychaete worm (*Cirriiformia luxuriosa*), 220
Polydactylus approximans, 20t, 23t
Polydactylus operaculis, 20t, 25t
Polydactylus sexfilis, 522t
 polygamy, 483, 496, 502–3
 polygyny, 497

- Polyplepium cruentum* (bleeding wrasse), 179t, 183f, 196f
- Polyprion americanus* (Atlantic wreckfish), 497, 521t
- Pomacanthus* spp., 235t, 522t
- Pomacentridae (damselfishes), 65, 74, 76, 106t, 547; in kelp bed/rocky reef, 235t, 237f; movement patterns of, 535t, 548; reproduction and, 488, 489, 491, 493, 495, 503, 506. *See also* *Chromis* spp.; *garibaldi* (*Hypsypops rubicundus*)
- Pomatomus saltatrix* (bluefish), 149, 521t
- Pomatoschistus*, 498
- Pommeranz, T., 305
- pompano, 305. *See also* Pacific pompano
- Pondella, D. J., II, 153
- Pontinus sierra*, 311f
- popeye blacksmelt (*Bathylagus ochotensis*), 349t, 365, 366t, 370f; larval stages of, 276t, 277f, 368
- porcupinefish (*Diodon hystrix*), 21t, 236t
- Porichthys* spp. (midshipmen), 129f, 189, 557; *P. analis* (darkedge midshipman), 178t, 182f, 195f
- Porichthys myriaster* (specklefin midshipman), 19t, 97f, 136t, 145t, 157t, 178t, 233t, 532t; foraging guild, 192f, 194f, 195f
- Porichthys notatus* (plainfin midshipman), 19t, 38, 136t, 233t, 335, 520t; on continental shelf, 98f, 178t, 181f, 182f, 183f; foraging guild, 192f, 194f, 195f, 196f; migration of, 188, 532t; reproduction and, 486f, 487, 489, 491, 493; in rocky intertidal zone, 209t, 210, 213, 219t; trophic relations and, 398f, 403
- Poroclinus rothrocki*, 17t, 22t
- Poromitra crassiceps* (crested bigscale), 352t, 358f, 365
- Posner, M., 539
- predation, 312, 323, 343, 428–45; adaptations of predators, 432–33; anti-predatory adaptations, 428–32; behavioral responses to, 430–32; community structure and, 442–43; density-dependent, 438–41, 442, 444; evolutionary influences on, 428–33; extinction and, 625; filial cannibalism, 501; future research on, 444–45; group living and, 429–30; growth of prey and, 434–35; habitat structural complexity and, 441–42, 444; hatching and, 495; monogamy and, 497; mortality and, 434, 437–38; movement of, 542; pollutants and, 598; prey demography and, 434–41; recruitment and, 418, 420, 421, 435–37, 442; regional and geographic comparisons, 443–44; size-selective, 445. *See also* feeding divergence; trophic interactions
- Priacanthidae, 66
- pricklebacks (Stichaeidae), 7t, 9f, 10f, 72, 186, 329; *Plagiogrammus hopkinsii* (criss-cross prickletback), 17t, 24t, 91f, 236t, 238f; reproduction and, 498; in rocky intertidal zone, 90f, 207, 209t, 210, 213, 216f, 217f. *See also* monkeyface prickletback; *Xiphister* spp.
- pricklebreast poacher (*Stellerina xyosterna*), 18t, 152f, 153t, 296
- prickly sculpin (*Cottus asper*), 126t, 128f, 129f, 132f, 296
- Priede, I. G., 375–76, 377, 540
- Prionace glauca* (blue shark), 20t, 89f, 326f, 330, 337t, 389, 518t; movement of, 524, 532t, 540, 542, 543; telemetry and, 527–28; trophic relations and, 401f
- Prionotus* spp., 277f, 280f; *P. evolans*, 520t; *P. ruscarius* (rough sea robin), 178t, 181f, 194f
- Prionotus stephanophrys* (lumptail sea robin), 20t, 22t, 178t, 181f, 182f, 277; foraging guild, 194f, 195f, 196f
- Pristigenys serrula*, 20t, 24t, 311f
- productivity, 138–39, 249. *See also* reproduction
- Prognathodes falcifer* (scythe butterflyfish), 20t, 24t, 220, 235t
- promiscuity, 496, 497, 503
- Pronotogrammus multifasciatus* (threadfin bass), 7t, 178t, 183f, 196f, 255
- Proposition 132 (California), 590–91
- Protomyctophum crockeri* (California flashlightfish), 351t, 357f, 359, 370f; larval stages of, 276t, 277f, 279f, 280, 306; swim bladder of, 366t, 367
- Psenes* spp. (driftfish), 333; *P. pellucidus* (blackrag), 328f, 330, 559, 900
- Psettichthys melanostictus* (sand sole), 17t, 97f, 296f, 605; on continental shelf, 179t, 181f, 194f; in surf zone, 152f, 153t, 155t
- Pseudobalistes flavimarginatus*, 523t
- Pseudobathylagus ochotensis* (robust blacksmelt), 349t, 365
- Pseudochromis olivaceus*, 521t
- Pseudopeneus grandisquamis*, 21t, 25t
- Pseudopeneus maculatus*, 522t
- Pseudopentaceros wheeleri*, 20t, 22t, 522t
- Psychrolutes phrictus* (blob sculpin), 353t, 372f, 521t
- Pteraclis aesticola*, 20t, 24t
- Pterocaesio diagramma*, 521t
- Pteroplatytrygon violacea*, 20t, 24t
- Pterygophora californica* (brown algae), 228, 246, 437–38, 474
- Ptilichthys goodei*, 522t
- Pudget Sound sculpin (*Ruscarius meanyi*), 7t
- pudgy cuskeel (*Spectrunculus grandis*), 353t, 369
- puffers (Tetraodontidae), 77, 137f, 149, 321, 394; *Canthigaster* spp., 503, 523t; *Lagocephalus lagocephalus*, 20t, 23t, 332. *See also* *Sphoeroides* spp.
- purse seines, 571, 572, 573, 575, 577, 588
- pygmy poacher (*Odontopyxis trispinosa*), 18t, 178t, 182f, 192f, 195f
- pygmy rockfish (*Sebastes wilsoni*), 17t, 255, 259, 260f, 263t
- pygmy shark (*Euprotomiscus bispinnatus*), 17t, 331
- Pyrosoma* sp. (colonial salp), 328f
- Quammen, M. L., 140, 141
- Quast, J. C., 232, 243–44, 404
- queenfish. *See* *Seriphus politus*
- Quietula y-cauda* (shadow goby), 19t, 88, 303, 534t; in bays and estuaries, 93f, 126t, 132f, 134f, 145t; trophic relations and, 396f, 401
- quillback rockfish (*Sebastes maliger*), 17t, 21t, 255, 537t, 539, 542
- quinaldine (ichthyocide), 231, 232
- Rachycentron canadensis*, 521t
- Radovich, J., 14, 628
- Radulinus* spp.: *R. asprellus* (slim sculpin), 17t, 22t, 178t, 195f; *R. boleoides*, 17t, 21t; *R. vinculus*, 17t, 24t
- ragfish (*Icosteus aenigmaticus*), 17t, 22t, 327f, 330, 523t
- rainbow runner (*Elegatis bipinnulatus*), 324
- rainbow scorpionfish (*Scorpaenodes xyris*), 20t, 25t, 70, 234t
- rainbow seaperch (*Hypsurus caryi*), 18t, 91f, 235t, 239f, 457; cleaner behavior of, 556f, 557t; movement of, 533t, 542, 544–45; in surf zone, 152, 155t, 162t
- rainbow surfperch, 240f
- rainbow trout, 31. *See also* *Oncorhynchus* spp.
- Rainwater, C. I., 364
- rainwater killifish (*Luacania parva*), 130, 131, 134f, 612t, 614
- Raja binoculata* (big skate), 18t, 177t, 181f, 182f, 183f, 518t; foraging guild, 194f, 195f, 196f
- Raja inornata* (California skate), 18t, 98f, 177t, 181f, 182f, 183f; foraging guild, 194f, 195f, 196f
- Raja rhina* (longnose skate), 18t, 101f, 177t, 184, 185f, 197f
- Raja stellata*, 17t, 22t
- Rajidae (skates), 106t, 175, 180t, 185f, 189, 353t; and commercial fisheries, 574; and reproduction, 483. *See also* *Bathyraja* spp.
- Ralston, S., 173, 417, 530
- range end points, 12f, 14
- Ranzania laevis* (slender mola), 21t, 24t, 311f, 332
- ratfish. *See* spotted ratfish (*Hydrolagus colliiei*)
- Rathbunella* spp. (ronquils): *R. allenii*, 235t; *R. hypoplecta* (stripedfin ronquill), 18t, 72, 72f, 522t, 532t; *R. jordani* (northern ronquill), 235t
- rays. *See* Batoidae (rays)
- Reavis, R. H., 527
- recreational fishing, 567, 580–92; (1850–1940), 580–83; (1941–2001), 583–84; barge fishing and, 582–83, 583f, 588; competition with commercial, 589–92; fishing methods, 582, 583–84; regulation of, 586–89
- recruitment, 187, 240–41, 388, 411–23, 631, 632; benthic environment and, 420–22; community structure and dynamics, 414–15; defined, 411; density-dependence and, 421–22; directions for research, 423; disturbance and, 472, 477; importance of, 411–12; larval dispersal and, 417–18; larval production and, 415–17; management and conservation in, 422–23; population structures and distribution, 413–14; predation and, 418, 420, 421, 435–37, 442; in rocky intertidal zone, 214; settlement phase of, 418–20, 435–37; spatial and temporal variation in, 415–22
- redbanded rockfish (*Sebastes babcocki*), 18t, 22t, 255
- redbelly tilapia (*Tilapia zilli*), 612t, 613f, 617
- red brotula (*Brosomphycis marginata*), 18t, 186, 233t, 238f, 519t
- red drum (*Sciaenops ocellatus*), 150
- red goatfish (*Pseudopeneus grandisquamosus*), 135t
- red hake (*Urophycis chuss*), 472
- red Irish lord (*Hemilepidotus hemilepidotus*), 17t, 485, 486f, 498, 500t, 521t
- redtail surfperch, 128f, 585t
- reduction fisheries, 570–74
- Reebs, S., 549
- reef finspot (*Paraclinus intergripinnis*), 19t, 92f, 236t, 238f, 403, 628; in bays and estuaries, 136t, 137t; hypoxia and, 468; in rocky intertidal zone, 211, 215t, 216f, 222f
- reef perch (*Micrometrus aurora*), 18t, 90f, 235t; feeding of, 394, 399f, 403; in rocky intertidal zone, 213, 215, 216f, 220f
- Regalecus glesne*, 21t, 25t, 519t

- Reinhardtius hippoglossoides*, 17t, 22t
Remilegia australis (whalesucker), 324
Remora spp., 324–25; *R. albescens*, 21t, 23t; *R. australis*, 20t, 22t; *R. brachyptera* (gray remora), 21t, 25t, 325; *R. osteochir*, 20t, 521t; *R. remora*, 21t, 23t, 324, 328f, 329f, 337t
Remorina albescens (white suckerfish), 325
remote sensing technology, 527–30, 528f
reproduction, 73, 76, 483–507; in bathopelagic fish, 368–69; behavioral ecology and, 495–502; body size and, 483, 493, 497, 498; coloration and, 488, 489, 490, 493, 494, 498; courtship and, 487–95; diel patterns of, 495, 507; evolutionary ecology and, 495–502; filial cannibalism and, 500–502; future research in, 505–7, 506t; gender allocation and, 487, 488t, 495–98, 502–4, 505f, 506; gonochorism and, 483, 497, 498, 502–3, 504, 507; hermaphroditism and, 487, 488t, 496–98, 502–3, 504, 506; hypoxia and, 471–72; life histories and, 187, 486–87; mating systems, 489, 495–99, 506; modes of, 483–86; monogamy, 496–98, 503; morphology and, 491–92; motor patterns and, 491; oviparity (egg laying) and, 483–84, 486–87, 499, 500–501; parental care and, 484, 485t, 486, 487, 495, 496, 499–500, 502–4, 506; pheromones in, 491, 493, 494; pollution and, 602; polygamy, 483, 496, 502–3; promiscuity and, 503; in rocky intertidal fishes, 213–14; seasonal, 498; sexual selection and, 498–99; in soft-bottom species, 186–87; spawning and, 488–89, 494, 495, 507, 587; spawning types, 483–86; species tables, 518–23t; sperm competition and, 499, 505, 506; temperate-tropical dichotomies in, 504; territoriality and, 489, 493, 495, 502; upwelling systems and, 503–4; viviparity, 483–84, 486, 502, 503, 504, 505, 507. *See also* ichthyoplankton
respiration, 323; hypoxia and, 207, 210, 211; in intertidal fishes, 208, 210–12, 222
rex sole. *See* *Glyptocephalus zachirus*
Rhacochilus toxotes (rubberlip seaperch), 19t, 75, 92f, 103, 494; activity patterns of, 533t, 546; competition and, 456–57, 458f; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 235t, 239f, 240f; prey dissemblance and, 64; trophic relations and, 400f, 405
Rhacochilus vacca (pile perch), 18t, 75, 92f, 111f, 260, 494; as benthivore, 65; habitat range of, 102, 103, 105t; in harbors, 157t, 159, 161t, 162t; in kelp bed/rocky reef, 235t, 239f, 240t, 243f; movement and activity patterns of, 533t, 542, 544–45; symbiotic relations of, 556f, 557t; trophic relations and, 400f, 405
Rhamphocottus richardsonii (grunt sculpins), 17t, 21t, 213, 500t, 521t
Rhinobatidae (guitarfishes), 176, 177t, 180t, 183f, 189, 233t. *See also* *Zapteryx exasperata*
Rhinobatis productus, 518t
Rhinobatis productus (shovelnose guitarfish), 19t, 137f, 233t, 397f, 568; as nearshore soft bottom fish, 97f, 157t, 158t
Rhinocodon typus (whale shark), 20t, 25t, 325, 331f, 333, 518t; movement of, 540
Rhinogobiops nicholsii (blackeye goby), 19t, 60, 61f, 76, 162t, 420, 506; habitat of, 102, 105t; in kelp bed/rocky reef, 236t, 238f, 239f, 240f, 247; in mid-depth rocky reef, 99f, 255, 260f, 263t; movement and activity patterns of, 523t, 531, 534t, 545; predation study of, 431–32f, 434–37f, 438, 439–42, 460–61; reproduction and, 494, 497; territoriality of, 455
Rhinoliparis barbulifer, 353t
Rhinomuraena quasita, 518t
Rhizoprionodon longurio, 20t, 25t
Rhombochirus osteochir (hardfin marlinsucker), 25t, 325
ribbon halfbeak (*Euleptorhampus viridis*), 331f, 333
Richardson, Sally L., 270, 295, 296, 298
Riegle, K. C., 212
Rimicola spp. (clingfish): *R. cabrilla* (Channel Islands clingfish), 7t, 17t, 24t; *R. dimorpha*, 18t, 24t; *R. eigenmanni* (slender clingfish), 19t, 25t, 92f, 111f, 236t, 238f; *R. muscarum* (kelp clingfish), 18t, 111f, 155t, 236t, 238f, 405
Ripley, W. E., 574
RIT. *See* rocky intertidal (RIT)
Roach, S. W., 231
Roberts, D. A., 245, 246
Robinson, C. J., 543
Robison, B. H., 543
robust black smelt (*Pseudobathylagus ochotensis*), 365, 370f
rockfish. *See* *Sebastes* spp. (rockfish)
rockhead (*Bothragonus swanii*), 17t, 70
rockpool blenny (*Hypoblennius gilberti*), 19t, 236t, 236t, 403; movement patterns of, 531, 532t, 543; in rocky intertidal zone, 209t, 210, 211, 215t, 216f
rock prickleback (*Xiphister mucosus*), 17t, 90f, 236t, 498, 522; in rocky intertidal zone, 207, 209t, 212, 215t, 216f, 219, 220f; trophic relations and, 394, 399f, 403
rock sole (*Lepidopsetta bilineata*), 17t, 22t, 605
rockweed gunnel (*Xerperes fucorum*), 18t, 90f, 111f; feeding of, 399f, 403; in rocky intertidal zone, 207, 208, 209t, 215t, 216f, 220
rock wrasse. *See* *Halichoeres semicinctus*
rocky intertidal (RIT) zone, 82–83t, 84f, 85t, 86f, 87, 205–23; aerial and aquatic respiration, 208, 210–12; age composition, 218; amphibious fishes in, 208, 209, 213; analysis of, 104t; behavioral and physiological traits, 208, 210–13; climate change and, 205, 221–22, 630; distribution patterns, 206–8; fish habitat, 205–6; horizonatal zonation, 208; osmoregulation and desiccation risk in, 212; recruitment, 214; reproduction in, 213–14; residents and visitors, 217–18; seasonal changes, 207–8; species abundance, 215, 217, 221; species groups in, 90f; stability and persistence in, 220–21; taxonomic composition, 214–15; trophic interactions in, 218–20, 399f, 403; vertical zonation, 206–8
rocky reef. *See* kelp bed/rocky reef (KBRR)
rocky subtidal (RST), 83t, 84f, 85t, 86f, 91f; analysis of, 104t; species groups in, 90f, 238f
Roedel, P. M., 588
Roncador stearnsii (spotfin croaker), 19t, 95f, 150t, 151f, 603
Rondeletia loricata, 520t
ronquils (Bathymasteridae), 100f, 186, 238f; *Rathbunella* spp., 18t, 72, 72f, 235t, 522t, 532t
Ronquilus jordani, 17t, 21t
Rosales-Casian, J. A., 139
Rosen, D. E., 55, 63
Rosenblatt, R. H., 6, 621, 628
rosethorn rockfish (*Sebastes helvomaculatus*), 17t, 22t, 255, 262
rosy lip sculpin (*Ascelichthys rhodorus*), 17t, 91f, 209t, 499, 500t, 507
rosy rockfish (*Sebastes rosaceus*), 19t, 22t, 100f, 255, 260f, 263t
rosy sculpin (*Oligocottus rubellio*), 17t, 23t, 90f, 234t, 238f, 399f; in rocky subtidal zone, 215, 216f, 217f, 220f
Rothschild, B. J., 309
rough abyssal grenadier (*Coryphaenoides yaquinae*), 353t, 369, 374–77, 540
roughback sculpin. *See* *Chitonotus pugetensis*
roughcheek sculpin: *Artedius creaseri*, 75, 92f, 238f; *Ruscarius creaseri*, 19t, 23t, 234t, 533t
roughies (Trachichthyidae), 66
rough sea robin (*Prionotus rescarius*), 178t, 181f, 194f
roundfishes, 189
round stingrays. *See* *Urolophus/Urobatis halleri*
rubberlip seaperch. *See* *Rhacochilus toxotes*
rubynose brotula (*Cataetx rubrirostris*), 186
Ruscarius creaseri (roughcheek sculpin), 19t, 23t, 234t, 533t
Ruvettus pretiosus (oilfish), 20t, 24t, 327f, 332
Rypticus spp., 521t
sabertooth blenny (*Plagiotremus azuleus*), 7t, 20t, 25t, 236t
sablefish (*Anoplopoma fimbria*), 18t, 101f, 106t, 306, 521t, 532t; in bathypelagic zone, 353t, 369, 372f; commercial fisheries and, 574, 575, 578, 580; on continental shelf, 172, 178t, 185f, 187; foraging guild, 192f, 197f; pollutants and, 598
Saccopharynx lavenbergi, 519t
Sacramento blackfish (*Orthodon microlepidotus*), 393
Sacramento sucker, 129f
saddleback gunnel (*Pholis ornata*), 17t, 153t, 399f
saddleback sculpin (*Oligocottus rimensis*), 17t, 21t, 90f, 209t, 215t, 216f, 399f
Sagamichthys abei (shining tubeshoulder), 350t, 354, 355f, 369, 519t
sailfin molly (*Poecilia latipinna*), 612t, 613f, 616
sailfin sculpin (*Nautichthys oculofasciatus*), 17t, 91f, 234t, 238f, 500t
sailfish (*Istiophorus platypterus*), 21t, 25t, 333
Sakuma, K. M., 305
salema. *See* *Xenistius californica*
Salmonidae (salmon), 29–32, 33, 72–73, 120, 128; climate change and, 632, 633; commercial fishing of, 570, 574, 575, 579, 580, 586; recreational fishing of, 580, 589; reproduction and, 487. *See also* *Oncorhynchus* spp.
salmon shark (*Lamna ditropis*), 17t, 22t, 89f, 326f, 330, 337t; movement of, 524
Salmo salar (Atlantic salmon), 612t, 618, 619
Salmo trutta (brown trout), 612t, 616
sampling techniques, 170–71, 229–32. *See also* CalCOFI samplings
Sanchez, C., 305
San Clemente Islands, 47, 48
sand bass. *See* *Paralabrax* spp.
sanddabs. *See* *Citharichthys* spp.
Sanderson, S. L., 393, 407
San Diego Province, 3, 4f, 239
San Diego Bay, 120, 135, 135t, 141f, 145t
sandpaper skate, 185f

- sand perch. *See* *Diplectrum* spp.
- sand sole (*Psettichthys melanostictus*), 17t, 97f, 296f, 605; on continental shelf, 179t, 181f, 194f; in surf zone, 152f, 153t, 155t
- San Francisco Bay, 120, 129, 131, 598; alien species in, 612, 613–18
- Santa Barbara Basin, 359
- Santa Barbara Channel, 261, 263, 458–59
- Santa Barbara Island, 48
- Santa Catalina Basin, 371
- Santa Catalina Island, 47, 48, 74, 422, 587, 589; Naples Reef compared to, 546–47; pollution off, 600, 601; predation study at, 430–31, 432f, 434, 441, 443
- Santa Cruz Island, 455, 457f, 458–59
- Sarda chiliensis* (Pacific bonito), 20t, 87, 89f, 236t, 585t; in coastal pelagic zone, 154, 157t, 158t, 159f, 160t; commercial fisheries and, 574, 575; feeding of, 406; in pelagic zone, 330, 332f, 336t, 337t; pollution and, 600, 601; recreational fishing and, 581, 582, 584; reproduction and, 494; seasonal movement of, 541–42; trophic relations and, 401f
- sardine and anchovy regimes, 15, 89, 281–82, 283–84f, 289; climate change and, 624, 625f, 633; competition and, 450; PDO and, 629–30, 629f, 630f, 631; trophic relations and, 403, 405
- sardines (Clupeidae), 72, 89, 91; commercial fishing of, 569, 570–73, 578–79, 588, 590; and subsistence fishing, 567, 568
- Sardinops sagax* (Pacific sardine), 19t, 72, 103, 145t, 233t, 260, 429; abundance of, 89, 91, 336t; commercial fisheries and, 570–73, 579; feeding of, 393, 394; larval stages of, 276t, 277–78f, 279, 293, 294f, 298; in pelagic zone, 94f, 330, 332f, 335t, 337t; and recreational fishing, 585t, 586; in surf zone, 150t, 153t, 157t, 160–61t; trophic relations and, 401f
- sargassum blenny, 137f
- sargassum pipefish (*Syngnathus peligius*), 324
- Sargassum* spp. (algae), 228, 324, 419; *S. palmeri*, 430, 436f
- sargo (*Anisotremus davidsoni*), 19t, 66, 67f, 74, 89, 94f, 136t, 534t; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 235t, 239f, 241f; phylogenetic relations and, 43, 44f, 45t, 46t
- Sargocentron suborbitalis*, 311f
- Sars, G. O., 269
- sawtooth eel (*Serrivomer sector*), 348, 349t, 355f
- scalychad sculpin (*Artedius harringtoni*), 17t, 91f, 213, 234t, 238f; reproduction and, 498, 500t, 502
- Scaridae (parrotfishes), 394, 506; *Nicholsina* spp., 7t, 14, 20t, 25t, 235t, 522t
- SCCWRP (Southern California Coastal Water Research Project), 168, 631
- Schaeffer, B., 55, 63
- Schmitt, Russell, 307, 429, 430–31, 459–60
- schooling species, 186, 323–24, 494; predation and, 429–30
- Schultz, S. A., 245
- Sciaenidae (croakers), 9f, 69, 105, 105t, 235t, 393, 581; on continental shelf, 175, 178t, 180t; larval stages of, 276t, 289, 294; movement and activity of, 535t–536t, 539–40, 548; pollution and, 602; species richness, 10f; in surf zone, 149, 150, 153. *See also* *Genyonemus lineatus* (white croaker); *Seriphus politus* (queenfish); white sea bass (*Atractoscion nobilis*)
- Sciaenops ocellatus* (red drum), 150
- scissortail chromis (*Chromis atrilobata*), 92f, 235t, 241f
- scissortail damselfish, 237f
- scleratinian coral reefs, 56–57
- Scleroplax granulata* (pea crab), 559f
- Scofield, W. L., 570, 587, 588
- Scolopsis* spp., 522t
- Scomber japonicus* (Pacific chub mackerel), 20t, 96f, 160t, 236t, 261, 279, 429; abundance of, 336t; in coastal pelagic zone, 154, 156, 157t, 159f, 330, 332f, 335t; commercial fishing of, 337t, 574–75, 576, 579, 585t; decline of, 588; feeding of, 406; historical fluctuation in, 624; larval stages of, 276t, 277–78f, 305, 311; recreational fishing of, 582, 583, 584; seasonal movement of, 541–42; trophic relations and, 401f, 406
- Scomberomorus concolor*, 19t, 24t
- Scomberomorus sierra*, 20t, 25t
- Scombridae (chub mackerels), 9f, 10f, 61, 105, 321, 329; genetic divergence in, 621; and reproduction, 494, 496
- Scopelogys tristic* (Pacific blackchin), 351t, 357f, 358, 519t
- Scopelogadus mizolepis*, 520t
- Scopeloberyx robustus* (longjaw bigscale), 352t, 365
- Scopelogadus mizolepis bispinosus* (twospine bigscale), 352t, 361, 365, 366t
- Scopelosaurus* spp., 519t
- Scorpaena guttata* (California/spotted scorpionfish), 19t, 70f, 136t, 255, 520t; fisheries and, 579, 585t, 586; foraging guild, 192f, 195f; habitat of, 99f, 102, 105t; in harbors, 157t, 160t, 162t; in kelp bed/rocky reef, 234t, 239f; in pelagic zone, 178t, 182f; pollutants and, 598; seasonal movement of, 536t, 542; trophic relations and, 395, 398f, 403
- Scorpaena histrio* (player scorpionfish), 234t
- Scorpaenichthys marmoratus* (cabezon), 19t, 70f, 162t, 270, 500t, 521t, 533t; in bays and estuaries, 132f; as brephoeipipelagic fish, 333, 334f; commercial fisheries and, 574, 579; ENSO and, 631; feeding of, 399t, 403; in kelp bed/rocky reef, 111f, 234t, 238f, 240f; in rocky intertidal zone, 91f, 215t, 216f, 217f, 218, 219t; subsistence fishing and, 568; in surf zone, 151, 152f, 153t, 155t
- Scorpaenidae (scorpionfishes and rockfishes), 59–60, 68, 77, 217f, 234t, 500; on continental shelf, 172, 175, 180t, 186, 258t; habitat for, 105, 106t; movement of, 536t–537t; in pelagic zone, 329, 333; pollution and, 601; and reproduction, 491, 499, 504. *See also* Cottidae; Hexagrammidae; *Sebastes* spp.
- Scorpaenodes xyris* (rainbow scorpionfish), 20t, 25t, 70, 234t
- Scorpi lineolata*, 522t
- Scripps Institution of Oceanography (SIO), 269
- scuba diving, 526–27
- sculpins. *See* *Artedius* spp.; *Clinocottus* spp.; Cottidae; *Icelinus* spp.; *Oligocottus* spp.
- Scyliorhinidae (cat sharks), 175, 177t, 180t, 485; *Parmaturus xanthurus* (filetail), 19t, 23t, 101f, 353t, 518t. *See also* *Apristurus* spp.
- Scytalina cerdale* (graveldiver), 17t, 90f, 215t, 216f, 220f, 522t; feeding of, 399f, 403
- scythe butterflyfish (*Prognathodes falcifer*), 20t, 24t, 235t
- sea basses. *See* *Atractoscion nobilis*; *Paralabrax* spp.; Serranidae
- sea chubs (Kyphosidae), 7t, 105, 106t, 235t, 534t, 557t; feeding of, 65–66. *See also* *Girella nigricans*; *Medialuna californiensis*
- seahorse (*Hippocampus ingens*), 20t, 25t, 135t, 137f, 484
- sea jellies (*Pelagia* spp.), 328f
- sea otter (*Enhydra lutris*), 624, 627, 628f
- seaperch. *See* *Phanerodon* spp.; *Rhacochilus* spp.
- sea robins (Triglidae), 176, 178t, 181f, 189. *See also* *Prionotus* spp.
- searsids (*Holthymia* spp.), 350t
- seasonality, 139–40, 248, 249; larval distribution, 296–98; in rocky intertidal zone, 207–8
- sea trout (*Cynoscion* spp.), 19t, 24t, 150
- sea urchins, 404, 441, 455, 476; climate change and, 625, 626, 627, 628f
- Sebastes* (rockfishes), 9f, 10f, 70, 75, 76, 77, 234t; climate change and, 624, 632, 633; commercial fishing of, 573–74, 576, 579, 590, 591; in deep rock habitat, 255–56, 257f, 258t, 259–64; feeding of, 59–60; habitat for, 91f, 96, 99f, 100, 102–3, 105; larval stages of, 276t, 277, 278f, 292, 294f, 296f, 299, 305; movement and activities of, 536t–537t, 539, 541, 544; recreational fishing of, 580, 581, 584, 589; recruitment of, 417; reproduction and, 484, 486, 493, 499, 504, 505; species richness of, 10, 255; subsistence fishing and, 568; in surf zone, 152; wastewater pollution and, 606; YOY abundance, 83, 260, 261, 262f, 263t, 440–41
- Sebastes* spp. (rockfish): *S. aleutianus*, 18t, 22t; *S. alutus* (Pacific perch), 17t, 22t, 258t; *S. aurora* (aurora rockfish), 18t, 22t, 101f, 278; *S. babcocki* (redbanded rockfish), 18t, 22t, 255; *S. borealis*, 17t, 21t; *S. brevispinis*, 17t, 21t; *S. constellatus* (starry rockfish), 19t, 23t, 99f, 234t, 260f, 263t; *S. cramerii* (darkblotched rockfish), 17t, 21t, 257, 589; *S. dallii* (calico rockfish), 19t, 98f, 187, 234t, 398f, 416, 536t, 542; *S. ensifer* (swordspine rockfish), 18t, 23t, 99f, 102, 105t, 255, 260f, 263t; *S. entomelas*, 17t, 22t; *S. eos* (pink rockfish), 18t, 23t, 100f, 258t; *S. gilli* (bronzespotted rockfish), 18t, 24t, 258t; *S. helvomaculatus* (rosethorn), 17t, 22t, 255, 262; *S. hopkinsi* (squarespot rockfish), 18t, 23t, 99f, 234t, 255, 257, 260f, 263t, 591; *S. inermis*, 493; *S. lentiginosus* (freckled rockfish), 18t, 25t, 255; *S. macdonaldi* (Mexican rockfish), 19t, 24t, 44f, 45t, 46t, 258t; *S. maliger* (quillback rockfish), 17t, 21t, 255, 537t, 539, 542; *S. melanostomus* (semaphore rockfish), 18t, 22t, 258; *S. moseri* (whitespeckled rockfish), 100f, 255; *S. nebulosus* (china rockfish), 17t, 76, 76f, 234t, 240t; *S. nigrocinctus*, 17t, 21t; *S. ovalis* (speckled rockfish), 18t, 23t, 99f, 255, 260f; *S. phillipsi* (chameleon rockfish), 17t, 23t, 258t; *S. proriger*, 17t, 22t; *S. reedi*, 17t, 21t; *S. rosaceus* (rosy rockfish), 19t, 22t, 100f, 255, 260f, 263t; *S. ruberrimus* (yelloweye rockfish), 17t, 22t, 255, 260f; *S. rubrivinctus* (flag rockfish), 7t, 18t, 99f, 255, 260f, 263t; *S. rufus* (bank rockfish), 18t, 23t, 100f, 102, 105t, 262f; *S. schlegelii*, 520t; *S. simulator* (pinknose rockfish), 18t, 24t, 100f, 255, 262f; *S. umbrosus* (honeycomb rockfish), 19t, 24t, 99f, 255, 260f, 586; *S. wilsoni* (pygmy rockfish), 17t, 255, 259, 260f, 263t; *S. zacentrus*, 18t, 23t

- Sebastes atrovirens* (kelp rockfish), 19t, 157t, 162t, 270, 415, 536t; feeding/foraging of, 60f, 67–68; in kelp bed/rocky reef, 92f, 234t, 237f, 239f, 240f; phylogeographic structures of, 34t, 38
- Sebastes auriculatus* (brown rockfish), 19t, 22t, 111f, 255, 270; in kelp bed/rocky reef, 92f, 234t, 240f; movement of, 536t, 542; in surf zone, 152, 155t
- Sebastes carnatus* (gopher rockfish), 19t, 23t, 219t, 258, 270, 585t; in kelp bed/rocky reef, 91f, 234t, 237f, 240f; movement of, 531, 536t; recruitment and, 415, 420; territoriality of, 453t, 455–56f, 457f
- Sebastes caurinus* (copper rockfish), 17t, 111f, 255, 260f, 270, 585t; on continental shelf, 178t, 182f, 195f; in kelp bed/rocky reef, 92f, 234t, 240t; recruitment of, 415; seasonal movement of, 536t, 539, 542; in surf zone, 152, 155t
- Sebastes chlorostichus* (green-spotted rockfish), 18t, 22t, 178t, 183f, 196f; in deep rock habitat, 100f, 255, 262f, 263t; movement of, 528, 536t, 539
- Sebastes chrysomelas* (black-and-yellow rockfish), 19t, 67f, 91f, 162t, 215t, 219t; in kelp bed/rocky reef, 234t, 237t, 240f; movement of, 531, 536t; recruitment of, 415, 420; territoriality of, 453t, 455–56f, 457f; trophic relations of, 399, 403
- Sebastes diploproa* (splinose rockfish), 18t, 22t, 101f, 258t, 279, 333; on continental shelf, 172, 178t, 185f; foraging guild, 192f, 197f; as generalist, 103, 105t; movement of, 536t, 541
- Sebastes elongatus* (green-striped rockfish), 18t, 22t, 102, 105t, 537t; in deep rock habitat, 100f, 255, 262f, 263t
- Sebastes flavidus* (yellowtail rockfish), 17t, 99f, 234t, 520t, 589; in deep rock habitat, 260f, 263t; movement of, 537t, 539
- Sebastes goodei* (chilepepper), 19t, 100f, 172, 262f; fisheries and, 576, 585t, 590
- Sebastes jordani* (shortbelly rockfish), 18t, 22t, 59, 100f, 178t, 183f, 192f, 537t; in deep rock habitat, 255, 262f; larval stages of, 276t, 285, 287f
- Sebastes levis* (cow cod), 18t, 23t, 100f, 103, 537t, 589; in deep rock habitat, 255, 262f, 263t; larval stages of, 285, 288f
- Sebastes melanops* (black rockfish), 17t, 70, 70f, 155t, 234t, 255, 475; rocky intertidal zone, 215t, 216f, 217t, 219t
- Sebastes miniatus* (vermillion rockfish), 18t, 22t, 99f, 234t, 493, 631; in deep rock habitat, 255, 260f, 263t; fisheries and, 585t, 586
- Sebastes mystinus* (blue rockfish), 18t, 59f, 91f, 215t, 416; in deep rock habitat, 255, 260f; fishing and, 584, 585t, 590; in kelp bed/rocky reef, 234t, 240f, 242, 243f; reproduction and, 494; seasonal movement of, 537t, 542; trophic relations and, 394, 404
- Sebastes paucipinnis* (bocaccio), 18t, 59, 234t, 334f, 484f, 632; commercial fishing of, 576, 589, 590; in deep rock habitat, 256, 260f, 261f, 263t; epibenthic, 539; habitat for, 100f, 103; larval stages of, 276t, 309f; movement of, 528, 537t, 541; recreational fishing of, 584, 585t; young-of-year (YOY), 111f, 261, 262f
- Sebastes pinniger* (canary rockfish), 18t, 99f, 255, 260f, 263t; fisheries and, 585t, 589, 591
- Sebastes rastrelliger* (grass rockfish), 18t, 59f, 90f, 111f, 234t, 403, 430; in rocky intertidal zone, 90f, 215t, 216f, 217f, 219t; in surf zone, 152, 153t, 155t, 157t
- Sebastes rosenblatti* (greenblotched rockfish), 18t, 23t, 178t, 192f, 196f, 537t; in deep rock habitat, 100f, 255, 262f, 263t
- Sebastes saxicola* (stripetail rockfish), 19t, 23t, 172, 178t, 258, 537t, 631; on continental shelf, 172, 178, 182f, 183f; foraging guild, 192f, 195f, 196f; recruitment of, 416; trophic relations, 398f, 403
- Sebastes semicinctus* (halfbanded rockfish), 19t, 23t, 99f, 178t, 183f, 196f, 631; in deep rock habitat, 255, 257, 260f, 263t
- Sebastes serranoides* (olive rockfish), 18t, 60f, 91f, 162t, 219t; in kelp bed/rocky reef, 234t, 237f, 240f, 243f; movement and activity of, 537t, 548; and recreational fishing, 584, 585t, 590; trophic relations and, 400f, 405
- Sebastes serriceps* (treefish), 18t, 23t, 67f, 162t, 537t; in kelp bed/rocky reef, 92f, 234t, 237f, 239f; trophic relations and, 400f, 405
- Sebastolobus* spp. (thornyheads), 34t, 333
- Sebastolobus alascanus* (shortspine thornyhead), 18t, 23t, 33, 353t, 369, 520t; on continental shelf, 172, 178t, 185f, 197f; habitat of, 101f, 102, 105t; movement of, 537t, 540
- Sebastolobus altivelis* (longspine thornyhead), 19t, 23t, 96, 101f, 311f, 334f; in bathypelagic zone, 353t, 369, 371, 372f; movement of, 537t, 540, 541
- sediment contamination, 596, 598, 600f, 604, 607f, 608. *See also* pollution
- Selene brevoorti* (Mexican lookdown), 7t, 20t, 308f
- Selene peruviana*, 20t, 25t
- semaphore rockfish (*Sebastes melanostomus*), 18t, 22t, 258
- Semicossyphus pulcher* (California sheephead), 19t, 23t, 92f, 235t, 259, 522t; cleaner behavior of, 556f, 557t; and commercial fishing, 574, 579; feeding of, 65, 394; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 237f, 239f, 241f, 243f; larval stages of, 270, 293, 295f, 297f, 302; movement and activity patterns of, 534t, 541, 546, 548; overfishing and, 625, 627, 628f; phylogenetic relations and, 35t, 44f, 45t, 46t, 49, 624; recruitment of, 414, 417; reproduction and, 487, 490, 491f, 496, 497; subsistence fishing and, 568; telemetry and, 527; trophic relations and, 400f, 405
- señorita (*Oxyjulis californica*), 18t, 58, 105t, 136t, 259, 522; activity patterns of, 534t, 544, 546, 548; cleaner behavior of, 555, 556f, 557t; feeding of, 65, 389, 390, 400f; in harbors, 159, 162t; in kelp bed/rocky reef, 87, 92f, 235t, 237f, 239f, 240f, 241f, 404–5; larval stages of, 293, 295f; macrophyte substratum and, 111; predation and, 431f, 436; reproduction and, 497, 498
- sensory systems, 213, 223; vision, 307, 343
- Seriola lalandi* (yellowtail), 18t, 23t, 87, 89f, 330, 332f; abundance of, 336t, 337t, 628; commercial fisheries and, 574, 575, 588; hunting behavior of, 62, 432f; in kelp bed/rocky reef, 234t, 241f; recreational fishing of, 580, 581f, 582, 583, 584–86; trophic relations and, 401f, 406
- Seriola rivoliana* (almaco jack), 20t, 25t, 62
- Seriphus politus* (queenfish), 19t, 91, 136t, 145t, 522t, 585t, 631; activity patterns of, 536t, 548; in coastal pelagic zone, 155t, 156, 157t, 159f, 160–61t; on continental shelf, 178t, 181f; foraging guilds, 192f, 194f; larval forms of, 305; nocturnality of, 66, 67f; in surf zone, 95f, 150–51, 150t, 154f; trophic relations and, 397f
- Serranidae (sea basses and groupers), 87, 105, 106t, 234t, 506, 537t–538t; commercial fishing and, 579, 591; on continental shelf, 175, 178t, 180t; *Mycteroperca* spp. (grouper), 19t, 20t, 23t, 25t, 58, 234t; reproduction and, 487, 502, 503. *See also* *Stereolepis gigas* (giant sea bass); *Paralabrax* spp.; white sea bass (*Atractoscion nobilis*)
- Serranus* spp. (serrano), 58, 496; *S. aequidens* (deepwater serrano), 178t, 180t, 196f; *S. fasciatus*, 521t; *S. psittacinus* (banded serrano), 234t; *S. tabacarius*, 521t
- Serrivomer sector* (sawtooth eel), 348, 349t, 519t
- Setran, A. C., 214
- Sette, Oscar E., 269
- sexual selection, 495, 498–99. *See also* hermaphroditism; reproduction
- shad. *See* American shad; threadfin shad
- shadow goby (*Quietus y-cauda*), 19t, 88, 303, 534t; in bays and estuaries, 93f, 126t, 132f, 134f, 145t; trophic relations and, 396f, 401
- shallow slope (SSLP), 85t, 86f
- Shanks, A. L., 414
- Shannon-Wiener (H') diversity index, 83, 85, 107–8
- sharks, 233t, 324, 326f, 632; basking shark (*Cetorhinus maximus*), 19t, 326f, 336t, 390, 518t; commercial fishing and, 574, 576, 579; hammerheads, 89f, 337t, 538t; megamouth (*Megachasma pelagios*), 327f, 528, 529f; movement of, 527–28, 530, 542; nocturnals, 95f; pollutant concentrations in, 602; reproduction and, 483, 484; in surf zone, 153; whale shark (*Rhinocodon typus*), 20t, 25t, 325, 331f, 333, 518t, 540. *See also* Carcharhinidae; *Heterodontes francisi*; Scyliorhinidae (cat sharks); Squalidae (dogfish sharks); Triakidae (hound sharks)
- sharksucker (*Echeneis naucrates*), 20t, 24t, 324, 329f
- sharpchin rockfish, 100f, 263t
- sharpchin slickhead (*Bajacalifornia barragei*), 349t
- sharpnose seaperch (*Phanerodon atripes*), 75, 555, 556f, 557f
- sheephead. *See* *Semicossyphus pulcher*
- shelf habitat, 83t, 84f, 100f. *See also* continental shelf
- shell mound assemblage, 264
- shimofuri goby (*Tridentiger bifasciatus*), 130t, 131, 142f, 612t, 614
- shiner perch. *See* *Cymatogaster aggregata*
- shining loosejaw (*Aristostomias scintillans*), 356, 357f, 366t, 370f
- shining tubeshoulder (*Sagamicthys abei*), 350t, 354, 355f, 369, 519t
- shokihaze goby (*Tridentiger barbatus*), 130t, 131, 612t, 614
- shortbelly rockfish (*Sebastes jordani*), 59, 100f, 183f
- shortfin corvina (*Cynoscion parvipinnis*), 135t
- shortfin mako (*Isurus oxyrinchus*), 20t, 22t, 89f, 390, 518t, 579; movement of, 527–28, 529f, 540; in pelagic zone, 326f, 331, 337t; pollution and, 601; trophic relations and, 401f, 406

- shortspine combfish. *See Zaniolepis frenata*
- shortspine thornyhead (*Sebastolobus alascanus*), 18t, 23t, 33, 353t 369, 520t; on continental shelf, 172, 178t, 185f, 197f; habitat of, 101f, 102, 105t; movement of, 537t, 540
- shoulderspot (*Caelorinchus scaphopsis*), 369
- shovelnose guitarfish (*Rhinobatis productus*), 19t, 137f, 233t, 397f, 568; as nearshore softbottom fish, 97f, 157t, 158t
- showy bristlemouth (*Cyclothone signata*), 276t, 350t, 354, 356f, 360t, 362–63, 366t, 370f; trophic relations of, 402f
- shrimp spp., 555, 558f, 559–60, 560f
- Sidera* spp., 518t
- Siebenaller, J. F., 374
- Siganus canaliculatus*, 523t
- signal blenny (*Emblemaria hypacanthus*), 432, 503, 522t
- silversides. *See* Atherinopsidae
- silverspotted sculpin (*Blepsias cirrhosus*), 17t, 151, 152f, 153t; reproduction in, 486, 499, 500t
- silver surfperch (*Hyperprosopon ellipticum*), 18t, 91f, 235, 240f, 585t; in surf zone, 152f, 153t
- Simberloff, D., 611
- sixspot pricklyback (*Kasatkia seigeli*), 7t
- skates. *See* *Bathyraxia* spp.; Rajidae (skates)
- skipjack tuna (*Katsuwonus pelamis*), 20t, 89f, 326f, 331, 337t, 573, 628
- skipper fishes, 211, 213
- sleep patterns, 549
- slender barracudina (*Lestidiops ringens*), 277, 280, 351t, 355f, 370f
- slender bigscale (*Melamphaes acanthomus*), 352t
- slender bristlemouth (*Cyclothone pseudopallida*), 276, 280
- slender clingfish (*Rimicola eigenmanni*), 19t, 25t, 92f, 111f, 236t, 238f
- slender cockscomb (*Anoplarchus insignis*), 17t, 21t, 238f
- slender hatchetfish (*Argyropelecus affinis*), 354, 356f, 359, 366t
- slender mola (*Ranzania laevis*), 21t, 24t, 311f, 332
- slender snipefish (*Macroramphosus gracilis*), 20t, 25t, 332, 520t
- slender sole. *See* *Lyopsetta exilis*
- slender sucker (*Phtheichthys lineatus*), 20t, 25t, 324
- slendertail lanternfish (*Goniichthys tenuiculus*), 277, 279f, 280, 370f
- slickheads. *See* Alepocephalidae
- slim sculpin (*Radulinus asprellus*), 17t, 22t, 178t, 195f
- slimy snailfish (*Liparis mucosus*), 18t, 151, 153t
- slipskin snailfish, 238f
- slope habitat, 83t, 84f, 258t. *See also* continental shelf and slope
- slough anchovy (*Anchoa delicatissima*), 19t, 396f, 401–2; in bays and estuaries, 93f, 126t, 134f, 138t, 145t
- smalldisk snailfish (*Careproctus gilberti*), 353t
- smallhead flyingfish, 331f, 401f
- small squaretail (*Tetragonurus cuvieri*), 18t, 23t, 328f, 330
- smelts, fisheries and, 573, 574, 581. *See* Atherinidae; Bathylagidae; Osmeridae
- Smith, C. I., 497
- Smith, D. R., 404
- Smith, G. B., 174, 197
- Smith, K. L., Jr., 371, 373, 376
- Smith, P. E., 305, 323
- Smith, R. W., 6
- Smith-Beasley, I., 348
- smooth grenadier (*Nezumia liolepsis*), 353t, 369
- smooth hammerhead, 89f, 337t
- smoothhead sculpin (*Artedius lateralis*), 18t, 90f, 209t, 216f, 238f, 399f; feeding of, 392, 394
- smooth puffer, 137f
- smooth stargazer (*Kathetostoma avertuncus*), 20t, 179t, 185f, 197
- smoothtongue. *See* *Leuroglossus stilbius*
- snailfishes. *See* *Careproctus* spp.; *Liparis* spp.
- snakehead eelpout (*Lycenchelys crotalinus*), 17t, 22t, 354t
- snake mackerel (*Gempylus serpens*), 21t, 25t, 327f, 329, 332
- snipe eels (*Avocettina* spp.), 349t, 354, 355f
- snubnose blacksmelt (*Bathylagus wesethi*), 349, 354, 355f, 360t, 370f; larval stages of, 276t, 277f, 279f, 280, 285–86, 289f, 368; swimbladder of, 365, 366t
- snubnose pipefish (*Cosmocampus arctus*), 19t, 92f, 234t, 238f
- snubnose sculpin (*Orthonopias triacis*), 18t, 23t, 60, 91f, 532t; in kelp bed/rocky reef, 234t, 238f
- sockeye salmon (*Oncorhynchus nerka*), 17t, 21t, 519t, 624–25, 626f
- soft bottom species, 111, 167, 601. *See also* continental shelf and upper slope
- softhead grenadier (*Malacocephalus laevis*), 369
- soldierfishes, 547, 548
- sole. *See* Pleuronectidae (righteye flounders)
- Solenostomus* spp., 520t
- Somniosus pacificus* (Pacific sleeper shark), 17t, 21t, 353t, 518t
- sonar systems, 527–28
- Sorenson, P. W., 493
- soupfin shark (*Galeorhinus galeus*), 19t, 89f, 233t, 337t, 576
- Soutar, A., 624
- Southern California Bight (SCB), 504, 630, 631; contaminants in, 595, 596
- Southern California Coastal Water Research Project (SCCWRP), 168, 631
- southern spearnose poacher (*Agonopsis sterletus*), 19t, 24t, 178t, 183f, 196f
- spatial patterns, 530–31, 539–43. *See also* movement and activity patterns
- spawning, 488–89, 494, 495, 507, 587. *See also* reproduction
- species abundance. *See* abundance
- species ranges, 8–9, 12f, 13f, 28f
- speckled rockfish (*Sebastes ovalis*), 18t, 23t, 99f, 255, 260f
- speckled sanddab (*Citharichthys stigmaeus*), 19t, 60, 61f, 236t, 258, 631; in bays and estuaries, 127t, 128f, 132f; on continental shelf, 96, 97f, 103, 176, 179t, 181f; foraging guilds, 192f, 194f; in harbors, 158, 161t; larval stages of, 276t, 277–78f, 280, 298, 299; movement of, 535, 543; pollution and, 601, 602f, 606; in surf zone, 152f, 153t, 155t; trophic relations and, 397f, 403
- specklefin midshipman (*Porichthys myriaster*), 97f, 136t, 145t, 157t, 158t; on continental shelf, 181f, 182f; trophic relations and, 397f, 403
- speckletail flounder (*Engyophrys sanctilaurentii*), 7t, 20t, 25t, 179t, 182f, 195f
- Spectrunculus grandis* (pudgy cusk-eel), 353t, 369
- Sphoeroides annulatus* (bullseye puffer), 20t, 25t, 95f, 96, 137f, 236t, 237f, 241f
- Sphoeroides lobatus* (longnose puffer), 7t, 20t, 25t
- Sphyræna argentea* (California barracuda), 19t, 236t, 330, 390, 523t, commercial fishing of, 574, 575f, 587; larval stages of, 295f, 297f, 300f, 302, 311; recreational fishing of, 581, 582, 583, 584, 585t; seasonal movement of, 541–42; in surf zone, 150t, 154, 155t, 158t
- Sphyræna ensis* (Mexican barracuda), 7t, 20t, 25t, 236t
- Sphyrna* spp. (hammerheads): *lewini* (scallop hammerhead), 538t; *S. tiburo*, 20t, 25t; *S. zygaena*, 18t
- spines, and predation, 429
- spiny dogfish. *See* *Squalus acanthias*
- spiny dreamer (*Oneirodes acanthias*), 352t, 358, 390, 395, 520t; trophic relations of, 402f
- Spirinchus starksi* (night smelt), 17t, 72, 151, 152f, 153t, 33t; on continental shelf, 177t, 181f, 194f
- Spirinchus thaleichthys* (longfin smelt), 17t
- splitnose rockfish (*Sebastes diploproa*), 18t, 21t, 101f, 258t, 279, 333; on continental shelf, 172, 178t, 185f; foraging guild, 192f, 197f; as generalist, 103, 105t; movement of, 536t, 541
- sponges, 486
- spotfin croaker (*Roncador stearnsii*), 95f, 150t, 151f, 603; and commercial fishing, 587
- spotfin mojarra (*Eucinostomus argenteus*), 178t, 181f, 182f, 194f, 195f
- spotfin surfperch (*Hyperprosopon anale*), 18t, 97f, 152f, 153t, 155t
- spot (*Leiostomus xanthurus*), 469, 472f
- spotted cabrilla (*Epinephelus analogus*), 20t, 24t, 234t
- spotted cusk-eel (*Chilara taylori*), 19t, 98f, 163, 176, 177t, 398f; on continental shelf, 182f, 183f, 185f; foraging guild, 192f, 194f, 195f, 196f, 197f; movement patterns of, 535t, 546
- spotted flounder (*Hippoglossina bollmani*), 179t, 182f, 195f
- spotted kelpfish (*Gibbonsia elegans*), 19t, 60, 61f, 92f, 111f, 155t, 532t; hypoxia and, 468; in kelp bed/rocky reef, 236t, 238f; in rocky intertidal zone, 207, 215t, 216f, 217, 220f; in surf zone, 145t, 154f, 155t, 159
- spotted lizardfish (*Synodus evermanni*), 177t, 182f, 183f, 195f, 196f
- spotted ratfish (*Hydrolagus colliiei*), 18t, 177t, 183f, 185f, 394, 532t; foraging guild, 196f, 197f; habitat of, 101f, 102, 103, 105t
- spotted sand bass. *See* *Paralabrax maculatofasciatus*
- spotted scorpionfish. *See* *Scorpaena guttata*
- spotted turbot (*Pleuronichthys ritleri*), 95f, 151, 155t, 157t, 181f; in bays and estuaries, 127t, 133, 136t, 145t; larval stages of, 292, 293, 294f, 300; pollutants and, 600, 603; trophic relations and, 397f
- Squalidae (dogfish sharks), 106t, 175, 177t, 180t. *See also* *Squatina californica*
- Squalus acanthias* (spiny dogfish), 19t, 136t, 157t, 158t, 160t, 429, 484, 518t; and commercial fisheries, 574; on continental shelf, 172, 177t, 182f, 183f, 195f, 196f; habitat of, 101f, 103; pollutants and, 598, 600
- squarespot rockfish (*Sebastes hopkinsi*), 18t, 23t, 99f, 234t, 591; in deep rock reef, 255, 257, 260f, 263t

- squaretails (*Tetragonurus* spp.), 324, 333, 523t;
T. cuvieri (small squaretail), 18t, 23t, 328f, 330
- Squatina californica* (angel shark), 19t, 95f, 184, 233t, 390, 518t; commercial fishing and, 579, 591; movement of, 527, 538t, 540; in surf zone, 153, 156f, 157t, 158t
- squid fisheries, 401f, 406, 575
- squirrel fishes, 547
- Stacey, N. E., 493
- staghorn sculpin (*Leptocottus armatus*), 18t, 209t, 500t, 532t, 613; in bays and estuaries, 93f, 126t, 128f, 129f, 132f, 134f, 138t, 143f; on continental shelf, 178t, 181f, 194f; in surf zone, 150, 153t, 155t, 157t; trophic relations and, 396f, 402
- stargazers (Uranoscopidae), 175, 180t, 522t, 604
- Starks, E. C., 574
- Starksia hoessii*, 522t
- Sarr, M. P., 539, 554
- starry flounder (*Platichthys stellatus*), 17t, 296f, 600, 605, 606; in bays and estuaries, 93f, 127f, 128f, 129f, 132f, 138t
- starry rockfish (*Sebastes constellatus*), 19t, 23t, 99f, 234t, 260f, 263t
- State Water Resources Control Board (SWRCB), 168, 169, 170
- Steele, M. A., 247, 419–20, 421, 460, 531; predation study of, 430, 431–32, 434–37, 438, 439–41, 442
- steelhead (*Oncorhynchus mykiss*), 29, 30t, 31, 32, 618; in bays and estuaries, 126t, 128f, 129f, 130t, 132f
- Stegastes partitus* (bicolor damselfish), 488
- Stegastes rectifraenum* (Cortez damselfish), 235t
- Stein, D. L., 374
- Stellerina xyosterna* (pricklebreast poacher), 18t, 152f, 153t, 296
- Steneck, R. S., 5–6
- Stenobranchius leucoparus* (northern lampfish), 351t, 357f, 359, 361–62, 363, 519t; bioluminescence in, 356; larval stages of, 276t, 277–78f, 282f, 284, 285f, 292, 294f, 299, 305; reproduction and, 368; swim bladder of, 366t, 367; trophic relations of, 402f
- Stenobranchius* spp., 298, 299
- Stephen, C. A., 621
- Stephens, John S., Jr., 158, 163, 531, 542, 630; on kelp bed/rocky reef species, 227, 230, 231, 241, 244, 246
- Stephenson, W., 295
- Stereolepis gigas* (giant sea bass), 18t, 87, 555, 624; in kelp bed/rocky reef, 92f, 234t, 237f, 239f, 241f; recreational fishing and, 583, 589; reproduction of, 494, 497; trophic relations, 400f, 405
- Sternopyx* spp. (hatchetfishes), 350t, 365, 366t, 406; *S. diaphana* (diaphanous hatchetfish), 350t, 354, 356f, 402f, 519t; *S. obscura*, 350t
- Stichaeidae (pricklebacks), 9f, 10f, 72, 186, 329; reproduction and, 498; in rocky intertidal zone, 90f, 207, 209t, 210, 213, 216f, 217f. *See also* monkeyface prickleback; *Xiphister* spp.
- sticklebacks, 120, 142f. *See also* three-spine stickleback
- stingray. *See* *Dasyatis* spp.; *Urolophus halleri*
- Stomias atriventer* (blackbelly dragonfish), 351t, 356, 357f, 366t, 370f, 395; larval stages of, 277, 279f, 280; trophic relations of, 402f, 406
- Stomiidae (dragonfishes), 307, 329, 350–51t, 354, 356
- storms, disturbance and, 474–77, 478f
- Strasburg, D. W., 555
- streaklight tubeshoulder (*Holothyrina latifrons*), 369
- striped bass. *See* *Morone saxatilis*
- stripedfin ronquil (*Rathbunella hypoplecta*), 18t, 72, 72f, 235t, 522t, 532t
- striped kelpfish (*Gibbonsia metzi*), 19t, 90f, 111f, 155t, 236t, 238f; in rocky intertidal zone, 215t, 216f; trophic relations of, 399f, 403
- striped marlin (*Tetrapterus audax*), 21t, 24t, 89f, 395, 401f, 433; in pelagic habitat, 326f, 332, 337t
- striped mullet (*Mugil cephalus*), 20t, 157t, 158t, 520t, 534t; in bays and estuaries, 95f, 126t, 129f, 134f, 145t, 398; as brephoepipelagic fish, 333, 334f; feeding and trophic relations, 390, 394, 396f
- striped sea chub (*Kyphosus analogus*), 7t
- striped seaperch (*Embiotoca lateralis*), 18t, 75, 92f, 585t; activity patterns of, 544–45; competition and, 450, 453t, 457–60, 461–62f, 463f; in kelp bed/rocky reef, 235t, 237f, 240f, 404; phylogenetic structures of, 34t, 38; recruitment of, 415, 416; in surf zone, 111f, 151, 152f, 153t
- stripefin poacher (*Xeneretmus ritteri*), 18t, 24t, 98f, 178t, 183, 185f, 197f
- stripetail rockfish (*Sebastes saxicola*), 19t, 23t, 100f, 103, 258, 537t, 631; on continental shelf, 172, 178, 182f, 183f; foraging guild, 192f, 195f, 196f; recruitment of, 416; trophic relations, 398f, 403
- Stromateoid fishes, 321, 324, 333
- Strongylocentrotus* spp. *See* sea urchins
- Strongylura exilis*, 20t, 520t
- sturgeon (*Acipenser* spp.), 125, 587, 602; *A. medirostris* (green sturgeon), 18t, 125, 126t, 128f, 130t; *A. transmontanus* (white sturgeon), 18t, 125, 126t, 128f, 518t
- subsistence fisheries, 524, 567–68, 580
- subtidal and reef species, 91f
- suckerfishes, 324–25. *See also* *Remora* spp.
- Sudis atrox*, 311f
- Suflamen verres* (orange-side triggerfish), 236t
- sunbeam lampfish (*Lampadena urophaos*), 277, 279f, 280, 351t
- superfoetation, 486–87, 500
- surf grass (*Phyllospadix* spp.), 228
- surfperches. *See* *Amphistichus* spp.; *Embiotocidae* (surfperches); *Hyperprosopon* spp.
- surf smelt (*Hypomesus pretiosus*), 17t, 72, 494–95, 519; in bays and estuaries, 93f, 127t, 128f, 129f, 132f; in surf zone, 151, 152f, 153t
- surf zone (SZ), 83t, 85t, 86f, 94, 95f, 96, 97f, 149–53; analysis of, 104t; drift algal beds and, 151–52, 154f
- surgeonfish (*Acanthurus* spp.), 329, 523t
- surveys, 253–54. *See also* CalCOFI
- Sustainable Fisheries Act (1996), 588
- swallowtail damsel (*Azurina hirundo*), 235t
- Swanson, C., 616
- Sweetwater Marsh, 144f
- swellhead snailfish (*Paraliparis cephalus*), 353t
- swell shark (*Cephaloscyllium ventriosum*), 21t, 23t, 95f, 153, 156f, 157t, 158t; activity patterns of, 536t, 544, 549; in kelp bed/rocky reef, 233t, 239f
- swimbladders and buoyancy, 365–68
- swordfish. *See* *Xiphias gladius*
- swordspine rockfish (*Sebastes ensifer*), 18t, 23t, 99f, 102, 105t; in deep rock reef, 255, 260f, 263t
- Syacium ovale* (oval flounder), 179t, 194f
- symbiotic relationships, 554–63, 558f; bioluminescence and, 557–58; cleaner behavior, 555–57, 557t; commensalism, 554, 558–61; mutualism, 554–58; parasitism, 486, 554, 555, 561–62, 605; shrimp and, 555, 558f, 559–60, 560f
- Symbolophorus californiensis* (California lanternfish), 351t, 357f, 358, 359, 402f; larval stages of, 276t, 277f, 279f, 280, 282f; swimbladder of, 366t, 367
- Symbolophorus* spp. (lightfish), 286
- Symphodus ocellatus*, 522t
- Symphodus tinca* (peacock wrasse), 502
- Symphurus atriacaudus* (tonguefish), 19t, 98f, 155t, 179t, 475, 523t, 533t; in bays and estuaries, 132f, 136t; on continental shelf, 181f, 182f, 183f, 194; foraging guild, 192f, 194f, 195f, 196f; in harbors, 158, 161t; larval stages of, 277, 293, 295f, 301, 310f; pollutants and, 603; trophic relations and, 398f
- Symphurus* spp. (tonguefish), 277f, 280f; *S. atrimentatus* (halfspotted tonguefish), 179t, 182f, 195f; *S. oligomerus* (whitetail tonguefish), 179t, 183f, 196f
- Synchiropus atrilabiatus* (blacklip dragonet), 7t, 20t, 25t, 179t, 183f, 196f
- Synchirus gilli*, 17t, 21t, 500t
- Syngnathus* spp. (pipefishes), 103, 106t, 234t, 507; and reproduction, 484; *S. exilis* (barcheek pipefish), 111t, 151f, 154f, 155t, 598; *S. fuscus*, 520t; *S. pelagius*, 324
- Syngnathus auliscus* (barred pipefish), 20t, 88, 97f, 111f, 152; in bays and estuaries, 93f, 126t, 134f, 145t
- Syngnathus californiensis* (kelp pipefish), 19t, 103, 111f, 234t, 238f, 507; in surf zone, 150t, 154f, 155t
- Syngnathus leptorhynchus* (bay pipefish), 18t, 95f, 111f, 429, 479; in bays and estuaries, 126t, 128f, 129f, 132f, 134f, 136t, 137f, 145t; in surf zone, 151, 152f, 153t; trophic relations and, 396f, 402
- Synodus* spp. (lizardfishes), 277f, 280, 294, 298, 329, 353t, 506, 519t; on continental shelf, 175, 177t, 180t; *S. evermanni* (spotted lizardfish), 177t, 182f, 183f, 195f, 196f; *S. scituliaceps* (lance lizardfish), 177t, 181f, 194f
- Synodus lucioceps* (California lizardfish), 18t, 98f, 155t, 538t; on continental shelf, 173, 177t, 181f, 182f, 183f; foraging guilds, 192f, 194f, 195f, 196f; larval stages of, 277, 293, 295f, 301, 302f; trophic relations and, 395, 397f, 398f
- Taaningichthys bathyphilus*, 351t, 366t
- Tactostoma macropus* (longfin dragonfish), 351t, 359, 370
- Talismania bifurcata* (threadfin slickhead), 349t, 353t
- tan bristlemouth (*Cyclothone pallida*), 350t
- Taractes longipinnis* (bigscale pomfret), 327f, 330
- Taraichthys steindachneri*, 20t, 25t
- Targett, Tim, 307
- Tarletonbeania crenularis* (blue lanternfish), 351t, 357f, 358, 359, 360t, 370f; larval stages of, 276t, 277–78f, 279, 306
- Tautoga unitis* (tautog), 498, 612t, 618
- Tautoglabrus adspersus* (cunner), 419
- Tavolga, W. N., 493
- taxonomic identifications, 231
- tectonism, 33, 36
- telemetry, 527–30, 539

- teleosts, 501, 504; evolution and, 55, 56–57, 68; reproduction and, 484, 488t
- temperate region, 330, 497; tropical region compared, 483, 502–4, 547
- temperature of water, 495, 499; distribution and, 76; fisheries and, 567, 579, 584, 586; movement patterns and, 542; ocean surface warming, 5, 6, 242; submergence and, 96
- Tenogobius sagittula*, 18t, 25t
- Teo, S. L. H., 139
- Terapon jarbua*, 522t
- territoriality, 531f, 539; competition and, 450, 451, 453t, 454, 455–56
- Terry, C., 230, 231
- Tetragonurus* spp. (squaretails), 324, 333, 523t; *T. cuvieri* (small squaretail), 18t, 23t, 328f, 330
- Tetraodontidae (puffers), 77, 137f, 149, 321, 394. *See also* *Sphoeroides* spp.
- Tetrapterus angustirostris*, 21t, 23t
- Tetrapterus audax* (striped marlin), 21t, 24t, 89f, 395, 401f, 433; in pelagic habitat, 326f, 332, 337t
- Thalassoma bifasciatum* (bluehead wrasse), 487, 488
- Thalassoma lucasanum* (Cortez rainbow wrasse), 235t
- Thaleichthys pacificus*, 17t
- Theragra chalcogramma*, 17t, 21t, 519t
- thermogeography, 5–6
- The Theory of Island Geography* (MacArthur & Wilson), 43
- thickhead lanternfish (*Hygophum atratum*), 277, 279f, 280, 370f
- thicklip conger (*Chilconger dentatus*), 177t, 183f, 195f, 196f
- Thomas, C. G., Jr and C. G. Thomas III, 589–90
- Thompson, W. F., 572, 587
- Thomson, C. J., 494, 567
- thornback (*Platyrhinoides triseriatus*), 18t, 97f, 157t, 158t, 233t; on continental shelf, 177t, 181f, 194f
- thornyheads. *See* *Sebastolobus* spp.
- threadfin bass (*Pronotogrammus multifasciatus*), 7t, 178t, 183f, 196f, 255
- threadfin dragonfish (*Bathophilus filifer*), 350t
- threadfin grenadier (*Coryphaenoides filifer*), 353t, 369
- threadfin sculpin (*Icelinus filamentosus*), 18t, 22t, 178t, 183f, 196f, 258t
- threadfin shad (*Dorosoma petenense*), 161t, 612t, 613f, 617
- threadfin slickhead (*Talismaania bifurcata*), 349t, 353t
- threebanded butterflyfish (*Chaetodon humeralis*), 20t, 24t, 235t
- three-spine stickleback (*Gasterosteus aculeatus*), 18t, 153t, 520t; in bays and estuaries, 126t, 128f, 129f, 132f, 142f; reproduction and, 488, 489, 491–92
- thresher shark (*Alopias vulpinus*), 19t, 233t, 542; bigeye thresher, 89f, 326f, 337t; and commercial fishing, 579, 591
- Thunnus* spp. (tuna), 322–23, 335t; bigeye tuna, 89f, 326f, 337t; fisheries and, 333f, 337t, 570–77, 579, 581; seasonal movement of, 530, 542; *T. obesus*, 20t, 23t; *T. orientalis*, 20t, 23t
- Thunnus alalunga* (albacore), 18t, 23t, 89f, 389, 401f; fisheries and, 573–74, 575, 584; in pelagic zone, 326f, 327f, 330, 336t, 337t; seasonal movement of, 536t, 542
- Thunnus albacares* (yellowfin tuna), 21t, 24t, 89f, 523t, 540, 573; in pelagic zone, 326f, 337t
- Thunnus thynnus* (bluefin tuna), 89f, 401f, 542; commercial fishing of, 573, 578; feeding behavior of, 389, 390; in pelagic zone, 326f, 330, 336t, 337t
- Thyrsites atune*, 523t
- tidpool blennie, 334f
- tidpool sculpin (*Oligocottus maculosus*), 17t, 91f, 500t, 507, 533t; in rocky intertidal zone, 207, 209t, 215t, 216f, 217f, 218
- tidpool snailfish (*Liparis flarae*), 17t, 90f, 215t, 216f, 238f
- tidewater goby (*Eucyclogobius newberryi*), 5, 18t, 132f, 614; in bays and estuaries, 126t, 128f, 129f, 130t, 134f; phylogeographic structure, 34t, 38, 39f
- Tijuana Estuary, 135–36, 144f, 400
- Tilapia* spp. (cichlids): *Oreochromis mossambicus* (Mozambique tilapia), 485, 612t, 617; *Oreochromis niloticus* (Nile tilapia), 393; *T. zilli* (redbelly tilapia), 485, 522t, 612t, 613f, 617
- timed transects, for density estimate, 230–31
- Tinbergen, N., 489
- toadfishes (Batrachoididae), 106t, 178t, 180t, 210, 532t; reproduction and, 493, 503, 504, 506
- Todd, E. S., 472
- tomcod (*Microgadus proximus*), 17t, 128f, 153t, 296f; on continental shelf, 174, 177t, 181t, 182f, 183f; foraging guild, 194f, 195f
- tonguefishes (Cynoglossidae), 298, 329; on continental shelf, 175, 179t, 180t, 189. *See also* *Symphurus* spp.
- topside lampfish (*Notorhynchus valdiviae*), 351t, 370f
- topsmelt. *See* *Atherinops affinis*
- Torpedinidae (torpedo electric rays), 175, 177t, 180t
- Torpedo californica* (Pacific electric ray), 18t, 95f, 156f, 233t, 348, 405, 518t; on continental shelf, 177t, 183f, 196f; hunting behavior of, 433f; movement patterns of, 538t, 540, 542, 549; in pelagic zone, 335, 337f
- Trachichthyidae, 66
- Trachinotus paitensis*, 20t, 25t
- Trachinotus rhodopus*, 20t, 24t
- Trachinus vipera*, 522t
- Trachipterus altivelis* (king-of-the-salmon), 20t, 327f, 332
- Trachipterus fukuzaki*, 21t, 25t
- Trachurus symmetricus* (jack mackerel), 19t, 23t, 77, 96f, 172, 432f, 521t; abundance of, 336t; in coastal pelagic zones, 154, 157t, 159f, 160t; commercial fishing of, 570, 571, 573, 574, 576, 578, 579, 591; DDT levels in, 601; feeding of, 406; genetic divergence in, 621; in kelp bed/rocky reef, 234t, 239f, 241f; larval stages of, 276t, 277f, 279f, 305; as pelagic species, 330, 332f, 335t, 337t; recreational fishing of, 581, 582; schooling by, 324; trophic relations and, 401f
- treefish (*Sebastes serriceps*), 18t, 23t, 67, 67f, 162t, 537t; in kelp bed/rocky reef, 92f, 234t, 237f, 239f; trophic relations and, 400f, 405
- triacylglycerols, deep water buoyancy and, 365–68
- Triakidae (hound sharks), 233t; *Galeorhinus galeus* (soupfin), 19t, 89f, 233t, 337t, 576. *See also* *Mustelus* spp.
- Triakis semifasciata* (leopard shark), 19t, 95f, 158t, 233t, 518t; in bays and estuaries, 127t, 128f, 132f; movement of, 527, 538t; in surf zone, 150t, 151, 153, 156f, 157t
- Trichiurus nitens* (cutlassfish), 20t, 25t, 327f, 332
- Trichodon trichodon*, 17t, 522t
- Trichonotus filamentosus*, 522t
- Tridentiger* spp. (gobies), 131, 614; *T. barbatus* (shokihaze goby), 130t, 131, 612t, 614; *T. bifasciatus* (shimofuri goby), 130t, 131, 142f, 612t, 614; *T. trigonocephalus* (chameleon goby), 612–13, 612t
- triggerfishes (*Balistes* spp.), 175, 179t, 180t, 236t; *B. polyplepis* (finescale triggerfish), 21t, 23t, 179t, 194f, 236t, 309f, 485, 523t
- Triglidae (sea robins), 176, 178t, 180t, 181f. *See also* *Prionotus* spp.
- Triphoturus mexicanus* (Mexican lampfish), 351t, 357f, 359, 362, 366t, 368, 370f; larval forms of, 276t, 277f, 280, 286, 290, 291f, 293–94, 298, 305; trophic relations of, 402f
- Triphoturus nigrescens* (highseas lampfish), 277
- Triphoturus oculus* (lampfish), 277
- trophic interactions, 394–407; in bays and estuaries, 396–98, 400–402; categories, 394–95; continental shelf, 397f–398f, 402–3; decline in, 631–32; deep midwater, 402f, 406; disturbance and, 471, 476; epipelagic fishes, 401f, 405–6; ichthyoplankton, 307–13; in kelp bed/rocky reef, 400f, 404–5; pollutants and, 598, 601; rocky intertidal, 399f, 403
- tropical hatchetfish (*Argyropelecus lychnus*), 350t, 356f, 359, 366t
- tropical region, 331–33; temperate region compared, 483, 502–4, 547
- true smelts (Osmeridae), 72, 73, 77, 120, 128–29, 149; on continental shelf and slope, 175, 177t, 180t; larval stages of, 296, 304f
- tubenose poacher (*Pallasina barbata*), 17t, 153t
- tubeshoulders (Platytroutidae), 350t, 354, 355f, 369, 519t
- tubesnout (*Autorhynchus flavidus*), 18t, 91f, 111f, 504, 520t; in kelp bed/rocky reef, 233t, 240f, 243f
- tuna. *See* *Thunnus* spp.
- turbots, 193. *See also* *Peuronichthys* spp.
- Turner, Charles H., 227
- two-line eelpout (*Bothrocara brunneum*), 18t, 22t, 354t, 372f
- twospine bigscale (*Scopelogadus mizolepis bispinosus*), 352t, 361, 365, 366t
- Typhlogobius californiensis* (blind goby), 19t, 24t, 211, 497, 523t; symbiotic relations of, 559–60, 560f
- Ugoretz, J., 230
- Ulvicola sanctaerosae* (kelp gunnel), 18t, 24t, 72, 72f, 236t
- umbrellamouth gulper (*Eurypharynx pelicanoides*), 348, 519t
- Umbrina roncadore* (yellowfin croaker), 19t, 95f, 127t, 134f, 145t; fisheries and, 581, 585t, 587; movement patterns of, 536t, 546; in surf zone, 150t, 151f, 154, 157t, 160t; trophic relations and, 396f, 402
- U. S. Bureau of Fisheries, 572
- United States Environmental Protection Agency (USEPA), 168, 170, 602
- Upogebia pugettensis* (mud shrimp), 559
- upwelling systems, 503–4
- Uranoscopidae (stargazers), 175, 180t, 522t, 604
- Uraspis secunda*, 20t, 25t

- Urechis caupo* (fat inkeeper worm), 559, 559f
 Urolophidae (round stingrays), 175, 177t, 180t
Urolophus/Urobatis halleri (round stingray), 20t, 233t, 397f, 429, 518t, 540; in bays and estuaries, 95f, 127t, 129f, 136t, 137f, 145t; on continental shelf, 173, 177t, 181f; foraging guilds, 194f, 195f; in surf zone, 150t, 151, 157t, 158t
Urophycis chuss (red hake), 472
Uticina lofotensis (white-spotted rose anemone), 561f
 Utter, E., 31
- Van Holle, D., 611
 Vehrencamp, S. I., 488
Venefica tentaculata, 519t
 vermilion rockfish (*Sebastes miniatus*), 18t, 22t, 99f, 234t, 493, 631; in deep rock habitat, 255, 260f, 263t; fisheries and, 585t, 586
 vertical distributions, 206–8, 305–6, 363
 Vetter, R. D., 540
 vicariance, gene flow and, 50
Vinciguerria lucetia (Pacific lightfish), 305, 350t, 368, 370f; larval stages of, 276t, 277f, 279f, 280, 282f, 286, 292f
Vinciguerria nimbaria, 519t
 vision, 307, 343
 viviparity, 483–84, 486, 502, 503, 504, 507
 Volpe, J. P., 618
- wahoo (*Acanthocybium solandri*), 333, 337t
 wakasagi (*Hypomesus nipponensis*), 130t, 131, 612t, 615–16
 Wakefield, W. W., 371
 Walker, Boyd W., 76, 152
 Walker, H. J., 297, 298
 wall-eye surfperch (*Hyperprosopon argenteum*), 18t, 67f, 95f, 186, 397f, 585t, 631; activity patterns of, 533t, 544, 546; in coastal pelagic zone, 155t, 157t, 159f, 160–61t; in kelp bed/rocky reef, 235t, 239f; in surf zone, 150–51, 150t, 152f, 153t, 154f
 Waples, Robin S., 31, 375
 warbonnets (*Chirolophus* spp.), 7t, 17t, 236t
 Warrant, E., 343
 Watanabe, H., 361
 Watson, W., 6, 291, 297–98, 369
 wax esters, buoyancy and, 367–68
 Weber, M. L., 589
 Weitzman, S. H., 348
 western mosquitofish (*Gambusia affinis*), 126t, 535t, 612t, 613f, 616–17
 whale shark (*Rhinocodon typus*), 20t, 25t, 325, 331f, 333, 518t; movement of, 540
 whalesucker (*Remilegia australis*), 324
 whip nose anglerfish (*Gigantactis* sp.), 358
 White, B. N., 621
 White, P. S., 466
 white croaker. *See* *Genyonemus lineatus*
 white sea bass (*Atractoscion nobilis*), 19t, 96f, 111f, 336t, 337t, 432, 603; and commercial fishing, 574, 579, 587; in kelp bed/rocky reef, 235t, 239f, 241f; larval stages of, 274, 275f; pollutants and, 598, 603; and recreational fishing, 580, 582, 584; and reproduction, 494; in surf zone, 151, 154f, 155t, 156f, 157t, 158t
 white seaperch. *See* *Phanerodon furcatus*
 white shark (*Carcharodon carcharias*), 20t, 22t, 89f, 337t, 390; movement of, 530, 534t, 542; pollution and, 601
 whitespeckled rockfish (*Sebastes moseri*), 100f, 255
- white sturgeon (*Acipenser transmontanus*), 18t, 125, 126t, 128f, 518t
 white suckerfish (*Remorina albescentis*), 325
 whitetail tonguefish (*Symphurus oligomerus*), 179t, 183f, 196f
 whitetip shark (*Carcharhinus longimanus*), 19t, 25t, 328f, 331
 widow rockfish, 99f, 257, 260f, 263t, 589
 Williams, E. H., 173
 Williams, G. C., 207, 543
 Wilson, Allan, 27
 Wilson, E. O., 46
 Wilson, Raymond R., Jr., 6, 369, 373, 374, 375, 377
 wolf-eel (*Anarrhichthys ocellatus*), 17t, 75, 75f, 91f, 255; in kelp bed/rocky reef, 236t, 240t; reproduction and, 485, 497, 498, 522t
 Wood, R., 56
 wooly sculpin (*Clinocottus analis*), 35t, 90f, 399f, 403, 628; hypoxia and, 468; parental care in, 500t, 507; in rocky intertidal zone, 215, 216f, 217f, 219, 222f
 wrasses, 437; bleeding wrasse (*Polylepium cruentum*), 179t, 183f, 196f; *Decodon melasma* (blackspot wrasse), 7t, 20t, 24t; *Thalassoma* spp., 235t, 487, 488. *See also* *Halichoeres* spp.; Labridae (wrasses)
- Xanthichthys mento*, 19t, 24t
Xeneretmus spp. (poachers): *X. latifons* (black-tip poacher), 18t, 23t, 98f, 178t, 183f, 192f, 196f, 532t; *X. leiops*, 17t, 21t; *X. ritteri* (stripefin poacher), 18t, 24t, 98f, 178t, 183, 185f, 197f; *X. triacanthus*, 18t, 23t
Xenistius californica (salema), 20t, 74, 111f, 136t, 145t, 154; activity patterns of, 534t, 548; cleaner behavior, 556f, 557t; in coastal pelagic zone, 157t, 159f, 160t; on continental shelf, 178t, 181f; foraging guilds, 194f, 195f; in kelp bed/rocky reef, 235t, 237f, 239f, 241f; as nocturnal predator, 66, 67f, 95f
Xerperes fucorum (rockweed gunnel), 18t, 90f, 111f; feeding of, 399f, 403; in rocky intertidal zone, 207, 208, 209t, 215t, 216f, 220f
Xiphias gladius (swordfish), 20t, 23t, 89f, 523t, 534; and commercial fisheries, 574, 579; diel movements of, 543; feeding of, 395, 433; in pelagic zone, 323, 326f, 330, 332, 337f; trophic relations and, 401f, 405, 406
Xiphister atropurpureus (black prickleback), 18t, 90f, 236t, 498, 522t; in rocky intertidal zone, 209t, 210, 211f, 215t, 216f; trophic relations and, 394, 399f, 403
Xiphister mucosus (rock prickleback), 18t, 90f, 236t, 498, 522; in rocky intertidal zone, 207, 209t, 212, 215t, 216f, 219, 220f; trophic relations and, 394, 399f, 403
Xystreus liolepis (fantail sole), 19t, 145t, 155t, 158t; on continental shelf, 179t, 181f, 182f; foraging guilds, 192f, 194f, 195f; larval stages of, 293, 295f, 300, 301f, 302
- yellowchin sculpin (*Icelinus quadriseriatus*), 19t, 23t, 398f, 403, 532t; on continental shelf, 98f, 173, 178t, 182f; foraging guilds, 192f, 195f
 yelloweye rockfish (*Sebastes ruberrimus*), 17t, 255, 260f, 263t
- yellowfin croaker (*Umbrina roncadore*), 19t, 95f, 127t, 134f, 145t; fisheries and, 581, 585t, 587; movement patterns of, 536t, 546; in surf zone, 150t, 151f, 154, 157t, 160t; trophic relations and, 396f, 402
 yellowfin fringehead (*Neoclinus stephensae*), 19t, 23t, 236t, 238f, 247
 yellowfin goby (*Acanthogobius flavimanus*), 95f, 533t, 612t, 613–14, 618; in bays and estuaries, 126t, 130t, 131, 132f, 142f
 yellowfin tuna (*Thunnus albacares*), 21t, 24t, 89f, 326f, 337t, 523t, 540, 573
 yellowstripe grunt (*Haemulopsis axillaris*), 178t, 182f, 194f, 195f
 yellowtail rockfish (*Sebastes flavidus*), 17t, 99f, 234t, 520t, 589; in deep rock habitat, 260f, 263t; movement of, 537t, 539
 yellowtail (*Seriola lalandi*), 18t, 23t, 87, 89f, 330, 332f; abundance of, 336t, 337t, 628; commercial fisheries and, 574, 575, 587; hunting behavior of, 62, 432f; in kelp bed/rocky reef, 234t, 241f; recreational fishing of, 580, 581f, 582, 583, 584, 585t, 586; trophic relations and, 401f, 406
 Yoklavich, M. M., 123, 133, 264
 Yoshiyama, R. M., 212, 214, 217, 218, 220, 221, 395
 Young, P. H., 497
 young-of-year (YOY), 83, 187, 248, 432, 474–75; rockfishes, 83, 260, 261, 262f, 263t, 440–41
- Zalemmbius rosaceus* (pink sea perch), 19t, 23t, 258, 263t, 533t; on continental shelf, 98f, 176, 178t, 182f, 183f; foraging guild, 192f, 195f, 196f; trophic relations and, 398f, 403
Zalieutes elater, 20t, 24t, 520t
Zanclus cornutus, 523t
Zaniolepis frenata (shortspine combfish), 19t, 23t, 100f, 102, 105t, 263t, 538t; on continental shelf, 178t, 183f, 258; foraging guild, 192f, 196f
Zaniolepis latipinnis (longspine combfish), 19t, 23t, 98f, 195f, 263t, 398f, 538t; on continental shelf, 176, 178t, 182f, 192f, 258
Zaniolepis spp. (combfish), 186, 255
Zaprora silenus, 17t, 21t, 522t
Zapteryx exasperata (banded guitarfish), 20t, 135t, 137f, 233t; on continental shelf, 177f, 183f, 194, 196f
 zebra goby (*Lythrypnus zebra*), 19t, 23t, 92t; activity patterns of, 534t, 545; in kelp bed/rocky reef, 236t, 238f
 zebraperch (*Hermosilla azurea*), 15, 19t, 87, 92f, 544, 630; genetic characteristics of, 44f, 45t, 46t; in harbors, 157t, 158t, 162t; in kelp bed/rocky reef, 92f, 235t, 237f, 241f, 404; symbiotic relations and, 556f, 558; trophic relations and, 394, 400f
 Zedler, Joy B., 135–36, 397
Zenopsis nebulosa, 17t, 23t, 520t
 Zerba, K. E., 241, 244
Zesticelus profundorum, 18t, 23t
 Ziegler, A. M., 68
 Zoarcidae (eelpouts), 9f, 10f, 72, 352t, 354t, 355f, 505; on continental shelf, 172, 175, 179t, 180t; reproduction and, 485, 498, 522t. *See also* *Lycodapus*; *Lycodes* spp.
 zoogeographic barriers, 26–27
 zooplanktivores, 73, 394–95, 548
Zostera marinus (eelgrass), 111, 134, 397, 475
Zu cristatus, 20t, 25t, 308f, 519t