

- Adams, S.L., and L.G. Allen. 2021. Seasonal distribution and abundance of tidepool fishes at six locations within the Southern California Bight, 2004-2005. Bull. S. Calif. Acad. Sci., 120(1):1-25.
- Alcaraz, G., C. Vanegas, and X. Chiappa-Carrara. 2002. Metabolic rate of *Bathygobius ramosus* upon a natural daytime tidal cycle. Oceanogr. Eastern Pacif., 11:60-66.
- Andrades, R., S. González-Murcia, T.J. Buser, et al. 2023. Ecology, evolution and conservation of tidepool fishes of the Americas. Rev. Fish Biol. and Fish., 33:1263-1290.
- Arora, H.L. 1948. Observations on the habits and early life history of the batrachoid fish, *Porichthys notatus* Girard. Copeia, 1948(2):89-93.
- Barber, W.E., L.M. McDonald., W.P. Erickson, and M. Vallarino. 1995. Effect of the *Exxon Valdez* oil spill on intertidal fish: a field study. Trans. Amer. Fish. Soc., 124:461-475.
- Barton, M.G. 1982. Intertidal vertical distribution and diets of five species of central California stichaeoid fishes. Calif. Fish Game, 68:174-182.
- Beck M.W., K.L. Heck Jr., K.W. Able, D.L. Childers, D.B. Eggleston, B.M. Gillanders, B. Halpern, C.G. Hays, K. Hoshino, T.J. Minello, R.J. Orth, P.F. Sheridan, and M.P. Weinstein. 2001. The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. BioScience, 51:633-641.
- Benitez, S., C. Duarte, T. Opitz, N. Lagos, and J. Pulgar. 2017. Intertidal pool fish *Girella laevifrons* (Kyphosidae) shown strong physiological homeostasis but shy personality: The cost of living in hypercapnic habitats. Mar. Pollut. Bull., 118(1-2):57-63.
- Benson, K.R. 2002. The study of vertical zonation on rocky intertidal shores – A historical perspective. Integr. Comp. Biol., 42:776-779.
- Blaxter, J.H.S., and J.R. Hunter. 1982. The biology of clupeoid fishes. Adv. Mar. Biol., 20:1-123.
- Bose, A.P., G.B. McClelland, and S. Balshine. 2015. Cannibalism, competition, and costly care in the plainfin midshipman fish, *Porichthys notatus*. Behav. Ecol., 27(2):628-636.
- Bose, A.P.H., B.G. Borowiec, G.R .Scott, and S. Balshine. 2019. Nesting on high: reproductive and physiological consequences of breeding across an intertidal gradient. Evol. Ecol., 33(1):21-36.
- Boyle, K.S. 2004. Comparison of feeding guild structure and ecomorphology of intertidal fish assemblages from central California and central Chile. Master's Thesis. Calif. State Univ., Fullerton.
- Boyle, K.S., and M.H. Horn. 2006. Comparison of feeding guild structure and ecomorphology of intertidal fish assemblages from central California and central Chile. MEPS, 319:65-84.
- Brantley, R.K., and A.H. Bass. 1994. Alternative male spawning tactics and acoustic signals in the plainfin midshipman fish *Porichthys notatus* Girard (Family, Batrachoididae). Ethology, 96:213-232.
- Breder, C.M. 1948. Observations on coloration in reference to behavior in tide-pool and other marine shore fishes. Bull. Am. Mus. Nat. Hist., 92:285-311.
- Bressman, N.R., A.C. Gibb, and S.C. Farina. 2018. A walking behavior generates functional overland movements in the tidepool sculpin, *Oligocottus maculosus*. Zoology (Jena), 131:20-28.
- Bridges, C.R. 1993. Ecophysiology of intertidal fish. Pp. 375-400. In: Fish ecophysiology. JC Rankin and FB Jensen (eds.). Chapman and Hall.

- Brown, C.R., M.S. Gordon, and K.L.M. Martin. 1992. Aerial and aquatic oxygen uptake in the amphibious Red Sea rockskipper fish *Alticus kirki* (Family Blenniidae). *Copeia*, 1992:1007-1013.
- Brown, N.A.W., N.S.B. Houpt, N.L. Yee, J.E.M. Curtis, B.M. Bolker, F. Juanes, and S. Balshine. 2021. Consequences of nest site selection vary along a tidal gradient. *J. Anim. Ecol.*, 90:528-541.
- Burgess, T.J. 1978. The comparative ecology of two sympatric polychromatic populations of *Xererpес fucorum* Jordan & Gilbert (Pisces: Pholididae) from the rocky intertidal zone of central California. *J. Exp. Mar. Biol. Ecol.*, 35(1):43-58.
- Carbia, P.S., and C. Brown. 2019. Environmental enrichment influences spatial learning ability in captive-reared intertidal gobies (*Bathygobius cocosensis*). *Anim. Cogn.*, 22(1):89-98.
- Chadwick, E.M.P. 1976. A comparison of growth and abundance for tidal pool fishes in California and British Columbia. *J. Fish Biol.*, 8:27-34.
- Chan, A.S., M.H. Horn, K.A. Dickson, and A. Gawlicka. 2004. Digestive enzyme activities in carnivores and herbivores: comparisons among four closely related prickleback fishes (Teleostei: Stichaeidae) from a California rocky intertidal habitat. *J. Fish Biol.*, 65(3):848-858.
- Clardy, T.R., E.J. Hilton, and W.K. Vogelbein. 2015. Morphology and ontogeny of multiple lateral-line canals in the rock prickleback, *Xiphister mucosus* (Cottiformes: Zoarcoidae: Stichaeidae). *J. Morphol.*, 276(10):1218-1229.
- Clayton, D.A., 1993. Mudskippers. *Oceanogr. Mar. Biol.: an annual review*, 31:507-577.
- Cogliati, K.M., Neff, B.D., and S. Balshine. 2013. High degree of paternity loss in a species with alternative reproductive tactics. *Behav. Ecol., Sociobiol.*, 67(3):399-408.
- Cogliati, K.M., S. Balshine, and B. Neff. 2014. Competition and cuckoldry: estimating fitness of alternative reproductive tactics in plainfin midshipman. *Behaviour*, 151(8):1209-1227.
- Coleman, R. 1992. Reproductive biology and female parental care in the cockscomb prickleback, *Anoplarchus purpurescens* (Pisces: Stichaeidae). *Environ. Biol. Fish.*, 41:177-186.
- Coleman, R.M. 1999. Parental care in intertidal fishes. Pp. 165-180. In: *intertidal fishes: life in two worlds*. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Colleye, O., B.J. Vetter, R.A. Mohr, L.H. Seeley, and J.A. Sisneros. 2019. Sexually dimorphic swim bladder extensions enhance the auditory sensitivity of female plainfin midshipman fish, *Porichthys notatus*. *J. Exp. Biol.*, 222(14):jeb204552.
- Compaire, J.C., P. Casademont, C. Gomez-Cama, and M.C. Soriguer. 2018. Reproduction and recruitment of sympatric fish species on an intertidal rocky shore. *J. Fish Biol.*, 92(2):308-329.
- Compaire, J.C., J. Gómez-Enri, C. Gómez-Cama, P. Casademont, O.V. Sáez, F. Pastoriza-Martin, C. Díaz-Gil, R. Cabrera, A. Cosin, and M.C. Soriguer. 2019. Micro-and macroscale factors affecting fish assemblage structure in the rocky intertidal zone. *MEPS*, 610:175-189.
- Congleton, J.L. 1974. The respiratory response to asphyxia of *Typhlogobius californiensis* (Family: Gobiidae) and some related gobies. *Biol. Bull.*, 146:186-205.
- Congleton, J.L. 1980. Observations on the responses of some southern California tidepool fishes to nocturnal hypoxic stress. *Comp. Biochem. Physiol.*, 66A:719-722.
- Crane, J.M. 1981. Feeding and growth by the sessile larvae of the teleost *Porichthys notatus*. *Copeia*, 1981(4):895-897.

- Craig, M.T., and Pondella, D.J., 2006. A survey of the fishes of the Cabrillo National Monument, San Diego, California. *Calif. Fish Game*, 92(4):172-183.
- Cross, J.N. 1981. Structure of a rocky intertidal fish assemblage. PhD Diss, Univ. Washington.
- Dabruzzi, T.F., M.L. Wygoda, J.E. Wright, J. Eme, and W.A. Bennett. 2011. Direct evidence of cutaneous resistance to evaporative water loss in amphibious mudskipper (family Gobiidae), and rockskipper (family Blenniidae) fishes from Palau Hoga, southeast Sulawesi, Indonesia. *J. Exp. Mar. Biol. Ecol.*, 406(1-2):125-129.
- Dall, W., and N.E. Milward. 1969. Water intake, gut absorption and sodium fluxes in amphibious and aquatic fishes. *Comp. Biochem. Physiol.*, 30:247-260.
- Davenport, J., A.D. Woolmington. 1981. Behavioural responses of some rocky shore fish exposed to adverse environmental conditions. *Mar. Freshw. Behav. Physiol.*, 8:1-12.
- Davis, J.L.D. 2000a. Changes in a tidepool fish assemblage on two scales of environmental variation: seasonal and El Niño Southern Oscillation. *Limnol. Oceanogr.*, 45:1368-1379.
- Davis, J.L.D. 2000b. Spatial and seasonal patterns of habitat partitioning in a guild of southern California tidepool fishes. *MEPS*, 196:253-268.
- Davis, J.L.D. 2001. Diel changes in habitat use by two tidepool fishes. *Copeia*, 2001:835-841.
- Davis, T.R., and M. Nimbs. 2022. Susceptibility of tidal pool fish assemblages to climate change. *Ecologies*, 3:510-520.
- Daxboeck, C., and T.A. Heming. 1982. Bimodal respiration in the intertidal fish, *Xiphister atropurpureus* (Kittlitz). *Mar. Behav. Physiol.*, 9:23-33.
- Dejours, P. 1994. Environmental factors as determinants in bimodal breathing: an introductory overview. *Am. Zool.*, 34:178-183.
- DeMartini, E.E. 1978. Spatial aspects of reproduction in the buffalo sculpin, *Enophryss bison*. *Environ. Biol. Fish.*, 3:331-336.
- Edwards, D.G., and J.J. Cech Jr. 1990. Aquatic and aerial metabolism of the juvenile monkeyface prickleback, *Cebidichthys violaceus*, an intertidal fish of California. *Comp. Biochem. Physiol.*, 96A:61-65.
- Eger, W.H. 1971. Ecological and physiological adaptations of intertidal clingfishes (Teleostei: Gobiesocidae) in the northern Gulf of California. PhD Diss., Univ. Arizona.
- Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A Field Guide to Pacific Coast fishes of North America. Houghton Mifflin.
- Evans, D.H., P.M. Piermarini, and K.P. Choe. 2005. The multifunctional fish gill: dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiol. Rev.*, 85(1):97-177.
- Evans, D.H., J.B. Claiborne, and G.A. Kormanik. 1999. Osmoregulation, acid-base regulation, and nitrogen excretion. Pp. 79-96, In: intertidal fishes: life in two worlds. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Fauth, J.E., J. Bernardo, M. Camara, W.J. Resetarits Jr., J. Van Buskirk, and S.A. McCollum. 1996. Simplifying the jargon of community ecology: a conceptual approach. *Am. Nat.*, 147:282-286.
- Feder, M.E., and W.W. Burggren. 1985. Cutaneous gas exchange in vertebrates: design, patterns, control and implications. *Biol. Rev.*, 60:1-45.
- Freeman, M.C., N. Neally, and G.D. Grossman. 1985. Aspects of the life history of the fluffy sculpin, *Oligocottus snyderi*. *Fish. Bull.*, 83:645-655.
- Fris, M.B., and M.H. Horn. 1993. Effects of diets of different protein content on food consumption, gut retention, protein conversion, and growth of *Cebidichthys violaceus*

- (Girard), an herbivorous fish of temperate zone marine waters. *J. Exp. Mar. Biol. Ecol.*, 166:185-202.
- Gaines, S.D., and J. Lubchenco. 1982. A unified approach to marine plant-herbivore interactions. II. Biogeography. *Annu. Rev. Ecol. Syst.*, 13:111-138.
- García-Huidobro, M.R., M. Aldana, C. Duarte, C. Galbán-Malagón, and J. Pulgar. 2017. Seawater-temperature and UV-radiation interaction modifies oxygen consumption, digestive process and growth of an intertidal fish. *Mar. Environ. Res.*, 129:408-412.
- Garey, W.F. 1962. Cardiac responses of fishes in asphyxic environments. *Biol. Bull.*, 122:362-368.
- German, D.P., M.H. Horn, and A. Gawlicka. 2004. Digestive enzyme activities in herbivorous and carnivorous prickleback fishes (Teleostei: Stichaeidae): ontogenetic, dietary, and phylogenetic effects. *Physiol. Biochem. Zool.*, 77(5):789-804.
- Giacomello, E., F.C. Neat, and M.B. Rasotto. 2008. Mechanisms enabling sperm economy in blenniid fishes. *Behav. Ecol. Sociobiol.*, 62(5):671-680.
- Gibb, A.C., M.A. Ashley-Ross, and S.T. Hsieh. 2013. Thrash, flip, or jump: the behavioral and functional continuum of terrestrial locomotion in teleost fishes. *Integr. Comp. Biol.*, 53(2):295-306.
- Gibson, R.N. 1969. The biology and behaviour of littoral fish. *Oceanogr. Mar. Biol. Annu. Rev.*, 7:367-410.
- Gibson, R.N. 1982. Recent studies on the biology of intertidal fishes. *Oceanogr. Mar. Biol. Annu. Rev.*, 20:363-414.
- Gibson, R.N. 1999. Movement and homing in intertidal fishes. Pp. 97-125. In: *Intertidal fishes: life in two worlds*. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Gibson, R.N., and R.M. Yoshiyama. 1999. Intertidal fish communities. Pp. 264-296. In: *Intertidal fishes: life in two worlds*. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Gonzales, T.T., M. Katoh, M.A. Ghaffar, and A. Ishimatsu. 2011. Gross and fine anatomy of the respiratory vasculature of the mudskipper, *Periophthalmodon schlosseri* (Gobiidae: Oxudercinae). *J. Morphol.*, 272(5):629-40.
- Graham, J.B. 1971. Aerial vision in amphibious fishes. *Fauna*, 1971:14-23.
- Graham, J.B. 1997. Air-breathing fishes: evolution, diversity and adaptation. Academic Press.
- Graham, J.B., and RH Rosenblatt. 1970. Aerial vision: unique adaptation in an intertidal fish. *Science*, 168:386-388.
- Graham, J.B., R.H. Rosenblatt, and C. Gans. 1978. Vertebrate air breathing arose in fresh waters and not in the oceans. *Evolution*, 32:459-463.
- Graham, J.B., C.B Jones, and I. Rubinoff. 1985. Behavioural, physiological, and ecological aspects of the amphibious life of the pearl blenny *Entomacrodus nigricans* Gill. *J. Exp. Mar. Biol. Ecol.*, 89:255-268.
- Grebel, J.M. 2003. Age, growth, and maturity of cabezon, *Scorpaenichthys marmoratus*, in California. Master's Thesis. Calif State Univ., Hayward.
- Green J.M. 1971. Local distribution of *Oligocottus maculosus* Girard and other tidepool cottids of the west coast of Vancouver Island, British Columbia. *Can. J. Zool.*, 49:1111-1128.
- Grier, H.J., W.F. Porak, J. Carroll, and L.R. Parenti. 2018. Oocyte development and staging in the Florida bass, *Micropterus floridanus* (LeSueur, 1822), with comments on the evolution of pelagic and demersal eggs in bony fishes. *Copeia*, 2018(2):329-345.

- Grossman, G.D. 1982. Dynamics and organization of a rocky intertidal fish assemblage: the persistence and resilience of taxocene structure. *Am. Nat.*, 119:611-637.
- Grossman, G.D. 1986a. Food resource partitioning in a rocky intertidal fish assemblage. *J. Zool. Lond. B*, 1:317-355.
- Grossman, G.D. 1986b. Long-term persistence in a rocky intertidal fish assemblage. *Environ. Biol. Fish.*, 15:315-317.
- Halpin, P.M., and K.L.M Martin. 1999. Aerial respiration in the salt marsh fish *Fundulus heteroclitus* (Fundulidae). *Copeia*, 1999:743-748.
- Haynes, T.B., E. Phillips-Mentzos, and D.E. Facey. 2009. A comparison of the hypersaline tolerances of black prickleback (*Xiphister atropurpureus*) and penpoint gunnel (*Apodichthys flavidus*). *Northwest Sci.*, 83(4):361-366.
- Helmuth, B.S.T., and G.E. Hofman. 2001. Microhabitats, thermal heterogeneity and physiological gradients of stress in the rocky intertidal zone. *Biol. Bull.*, 201:374-384.
- Heras, J., M. Chakraborty, J.J. Emerson, and D.P. German. 2020. Genomic and biochemical evidence of dietary adaptation in a marine herbivorous fish. *P. Roy. Soc. B*, 287:20192327.
- Herrera M.J., J. Heras, and D.P. German. 2022. Comparative transcriptomics reveal tissue level specialization in prickleback fishes. *J. Comp. Physiol. B*, 192:275-295.
- Hillman, S.S., and P.C. Withers. 1987. Oxygen consumption during aerial activity in aquatic and amphibious fish. *Copeia*, 1987:232-234.
- Horn, M.H. 1989. Biology of marine herbivorous fishes. *Oceanogr. Mar. Biol. Annu. Rev.*, 27:167-272.
- Horn, M.H., and L.G. Allen. 1978. A distributional analysis of California coastal marine fishes. *J. Biogeogr.*, 5(1):23-42.
- Horn, M.H., and R.N. Gibson. 1988. Intertidal fishes. *Sci. Am.*, 258(1):64-71.
- Horn, M.H., and F.P. Ojeda. 1999. Herbivory. Pp. 197-222. In: *intertidal fishes: life in two worlds*. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Horn, M.H., and K.C. Riegle. 1981. Evaporative water loss and intertidal vertical distribution in relation to body size and morphology of stichaeoid fishes from California. *J. Exp. Mar. Biol. Ecol.*, 50:273-288.
- Horn, M.H., K.F. Mailhiot, M.B. Fris, and L.L. McClanahan. 1995. Growth, consumption, assimilation and excretion in the marine herbivorous fish *Cebidichthys violaceus* (Girard) fed natural and high protein diets. *J. Exp. Mar. Biol. Ecol.*, 190:97-108.
- Horn, M.H., K.L.M. Martin, and M.A. Chotkowski. 1999. Introduction. Pp. 1-6. In: *intertidal fishes: life in two worlds*. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Horn, M.H., S.N. Murray, and T.W. Edwards. 1982. Dietary selectivity in the field and food preferences in the laboratory for two herbivorous fishes (*Cebidichthys violaceus* and *Xiphister mucosus*) from a temperate intertidal zone. *Mar. Biol.*, 67:237-246.
- Horn, M.H., S.N. Murray, and R.R. Seapy. 1983. Seasonal structure of a central California rocky intertidal community in relation to environmental variables. *Bull. S. Calif. Acad. Sci.*, 82:79-94.
- Horn, M.H., M.A. Neighbors, and S.N. Murray. 1986. Herbivore responses to a seasonally fluctuating food supply: growth potential of two temperate intertidal fishes based on the protein and energy assimilated from their macroalgal diets. *J. Exp. Mar. Biol. Ecol.*, 103:217-234.

- Houpt, N., B.G. Borowiec, A.P.H. Bose, N.A.W Brown, G.R. Scott, and S. Balshine. 2020. Parental males of the plainfin midshipman are physiologically resilient to the challenges of the intertidal zone. *Physiol. Biochem. Zool.*, 93(2):111-128.
- Hubbs, C.L. 1921. The ecology and life history of *Amphigonopterus aurora* and other viviparous perches of California. *Biol. Bull.*, 40:181-209.
- Hughes, G.M., C. Albers, D. Muster, and W.H. Goetz. 2013. Respiration of the carp *Cyprinus carpio* L., at 10 and 20°C and the effects of hypoxia. *J. Fish Biol.*, 22:613-628.
- Ip, Y.K., and S.F. Chew. 2018. Air-breathing and excretory nitrogen metabolism in fishes. *Acta Histochem.*, 120(7):680-690.
- Ip, Y.K., S.F. Chew, and J. Randall. 2002. Five tropical fishes, six different strategies to defend against ammonia toxicity on land. *Comp. Biochem. Physiol. A*, 134:S113-114.
- Irelan, C.D., and M.H. Horn. 1991. Effects of macrophyte secondary chemicals on food choice and digestive efficiency of *Cebidichthys violaceus* (Girard), an herbivorous fish of temperate marine waters. *J. Exp. Mar. Biol. Ecol.*, 153:179-194.
- Ishimatsu, A., H.V. Mai, and K.L.M Martin. 2018. Patterns of fish reproduction at the interface between air and water. *Integr. Comp. Biol.*, 58(6):1064-1085.
- Ito T., B.N. Goldman, S. Awata, and K.D. Crow. 2024. Mating season, egg-laying season, and internal gametic association in the sympatrically occurring fluffy sculpin (*Oligocottus snyderi*) and Rosy Sculpin (*O. rubellio*). *Ichthy. Herpet.*, 112(1):10-20.
- Jones, B.C. 1972. Effect of intertidal exposure on survival and embryonic development of Pacific herring spawn. *J. Fish. Res Bd. Can.*, 29:1119-1124.
- Jones, J.A. 1981. Competition for substrates in laboratory experiments between *Anoplarchus purpurescens* (Pisces, Stichaeidae) and three related species from the central California rocky intertidal zone. Master's Thesis. Calif State Univ., Fullerton.
- Kaplanis, N.J., C.B. Edwards, Y. Eynaud, and J.E. Smith. 2020. Future sea-level rise drives rocky intertidal habitat loss and benthic community change. *Peer J*, 8:e9186.
- Kim, K.H., M.H. Horn, A.E. Sosa, and D.P. German. 2014. Sequence and expression of an α -amylase gene in four related species of prickleback fishes (Teleostei: Stichaeidae): ontogenetic, dietary, and species-level effects. *J. Comp. Physiol. B*, 184:221-234.
- Klein, A., H. Münz, and H. Bleckmann. 2013. The functional significance of lateral line canal morphology on the trunk of the marine teleost *Xiphister atropurpureus* (Stichaeidae). *J. Comp. Physiol. A*, 199(9):735-749.
- Knopf, M.L., K.A. Tice, and D.C. Rypkema. 2017. Site fidelity and homing behaviour of intertidal sculpins revisited. *J. Fish Biol.*, 90(1):341-355.
- Lefevre, S., M. Bayley, D.J. McKenzie, and J.F. Craig. 2014. Air-breathing fishes. *J. Fish Biol.*, 84(3):547-553.
- Leigh, E.G. Jr, R.T. Paine, J.F. Quinn, and T.H. Suchanek. 1987. Wave energy and intertidal productivity. *PNAS*, 84:1314-1318.
- LeMoine, CM., C. Bucking, P.M. Craig, and P.J. Walsh. 2014. Divergent hypoxia tolerance in adult males and females of the plainfin midshipman (*Porichthys notatus*). *Physiol. Biochem. Zool.*, 87(2):325-333.
- Love, M.S., and J.K. Passarelli. 2020. Miller and Lea's guide to the coastal marine fishes of California. 2nd ed. Univ. Calif. Agric. Natur. Resourc. Publ. 3556.
- Low, W.P., Y.K. Ip, and D.J.W. Lane. 1990. A comparative study of the gill morphometry in the mudskippers – *Periophthalmus chrysospilos*, *Boleophthalmus boddaerti* and *Periophthalmodon schlosseri*. *Zool. Sci.*, 7:29-39.

- Luck, A., and K.L.M. Martin. 1999. Tolerance of forced air emergence by a fish with a broad vertical vertical distribution, the rockpool blenny, *Hypsoblennius gilberti* (Blenniidae). Environ. Biol. Fish., 54:295-301.
- Mandic, M., M.L. Ramon, A.Y. Gracey, and J.G. Richards. 2014. Divergent transcriptional patterns are related to differences in hypoxia tolerance between the intertidal and the subtidal sculpins. Mol. Ecol., 23(24):6091-6103.
- Marliave, J.B. 1977. Substratum preferences of settling larvae of marine fishes reared in the laboratory. J. Exp. Mar. Biol. Ecol., 27:47-60.
- Marliave, J.B. 1981. High intertidal spawning under rockweed, *Fucus distichus*, by the sharpnose *Clinocottus acuticeps*. Can. J. Zool., 59:1122-1125.
- Marliave, J.B. 1986. Lack of planktonic dispersal of rocky intertidal fish larvae. Trans. Amer. Fish. Soc., 115:149-154.
- Marliave, J.B., and E.E. DeMartini. 1977. Parental behavior of intertidal fishes of the stichaeid genus *Xiphister*. Can. J. Zool., 55:60-63.
- Marshall, W.H., and T.W. Echeverria. 1992. Age, length, weight, reproductive cycle and fecundity of the monkeyface prickleback (*Cebidichthys violaceus*). Calif. Fish Game, 78:57-64.
- Martin, K.L. 1991. Facultative aerial respiration in an intertidal sculpin, *Clinocottus analis*. Physiol. Zool., 64:1341-1355.
- Martin, K.L. 1993. Aerial release of CO₂ and respiratory exchange ratio in intertidal fishes out of water. Environ. Biol. Fish., 37:189-196.
- Martin, K.L. 1995. Time and tide wait for no fish: intertidal fishes out of water. Environ. Biol. Fish., 44:165-181.
- Martin, K.L. 1996. An ecological gradient in air-breathing ability among marine cottid fishes. Physiol. Zool., 69:1096-1113.
- Martin, K.L. 2014. Theme and variations: amphibious air-breathing intertidal fishes. J. Fish Biol., 84(3):577-602.
- Martin, K.L.M. 2015. Beach-spawning fishes: reproduction in an endanctered ecosystem. Taylor & Francis Group, CRC Press.
- Martin, K.L.M., and C.R. Bridges. 1999. Respiration in water and air. Pp. 54-78. In: intertidal fishes: life in two worlds. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Martin, K.L., and J.R.B. Lighton. 1989. Aerial CO₂ and O₂ exchange during terrestrial activity in an amphibious fish, *Alticus kirki* (Blenniidae). Copeia, 1989:723-727.
- Martin, K.L.M., R.C. Van Winkle, J.E. Drais, and H. Lakisic. 2004. Beach-spawning fishes, terrestrial eggs, and air breathing. Physiol. Biochem. Zool., 77(5):750-759.
- Martins, J., F. Almada, A. Goncalves, P. Duarte-Coelho, and P.E. Jorge. 2017. Home sweet home: evidence for nest-fidelity in the rocky intertidal fish, the shanny *Lipophrys pholis*. J. Fish Biol., 90(1):156-166.
- Matson, R.H., C.B. Crabtree, and T.R. Haglund. 1986. Ichthyofaunal composition and recolonization in a central California tidepool. Calif. Fish Game, 72:227-231.
- McArley, T.J., A.J.R. Hickey, and N.A. Herbert. 2018. Hyperoxia increases maximum oxygen consumption and aerobic scope of intertidal fish facing acutely high temperatures. J. Exp. Biol., 221(22):jeb189993.
- McArley, T.J., A.J.R. Hickey, L. Wallace, A. Kunzmann, and N.A. Herbert. 2019. Intertidal triplefin fishes have a lower critical oxygen tension (Pcrit), higher maximal aerobic

- capacity, and higher tissue glycogen stores than their subtidal counterparts. *J. Comp. Physiol. B*, 189(3-4):399-411.
- Metaxas, A., and R.E Scheibling. 1993. Community structure and organization of tidepools. *MEPS*, 98:187-198.
- Mgaya, Y.N. 1992. Density and production of *Clinocottus globiceps* and *Oligocottus maculosus* (Cottidae) in tidepools at Helby Island, British Columbia. *MEPS*, 85:219-225.
- Miller, E.C. 2023. Historical biogeography supports Point Conception as the site of turnover between temperate East Pacific ichthyofaunas. *PLoS ONE*, 18(9):e0291776.
- Miller, D.J., and R.N. Lea. 1972. Guide to the coastal marine fishes of California. Calif. Dept. Fish Game Fish Bull. 157.
- Miller, K.A., and W.H. Marshall. 1987. Food habits of large monkeyface prickleback, *Cebidichthys violaceus*. *Calif. Fish Game*, 73:37-44.
- Miranda, J.A., R.F. Oliveira, L.A. Carneiro, R.S. Santos, and M.S. Grober. 2003. Neurochemical correlates of male polymorphism and alternative reproductive tactics in the Azorean rock-pool blenny, *Parablennius parvicornis*. *Gen. Comp. Endocrinol.*, 132(2):183-189.
- Montgomery, W.L. 1977. Diet and gut morphology in fishes, with special reference to the monkeyface prickleback, *Cebidichthys violaceus* (Stichaeidae: Blennioidei). *Copeia*, 1977:178-182.
- Moring, J.R. 1976. Estimates of population size for tidepool sculpins, *Oligocottus maculosus*, and other intertidal fishes, Trinidad Bay, Humboldt County, California. *Calif. Fish Game*, 62:65-72.
- Moring, J.R. 1979. Age structure of a tidepool sculpin, *Oligocottus maculosus*, population in northern California. *Calif. Fish Game*, 65:111-113.
- Moring, J.R. 1983. Human factors affecting rocky intertidal fishes Pp. 1572-1585. In: Proceedings of the third symposium on coastal and ocean management, Coastal Zone '83. O.T. Magooon, H. Converse (eds.). Am. Soc. Civil Engineers.
- Moring, J.R. 1986. Seasonal presence of tidepool fishes in a rocky intertidal zone of northern California, USA. *Hydrobiologia*, 134:21-27.
- Moring, J.R. 1990. Seasonal abundance of fishes in tidepools of a boreal environment (Maine, USA). *Hydrobiologia*, 194:163-168.
- Morris, R.W. 1952. Spawning behavior of the cottid fish *Clinocottus recalvus* (Greeley). *Pacif. Sci.*, 6:256-258.
- Morris, R.W. 1956. Clasping mechanism of the cottid fish *Oligocottus snyderi* Greeley. *Pacif. Sci.*, 10:314-317.
- Murray, S.N., and M.H. Horn. 1989. Seasonal dynamics of macrophyte populations from an eastern North Pacific rocky-intertidal habitat. *Bot. Mar.*, 32:457-473.
- Nakamura, R. 1976. Temperature and the vertical distribution of two tidepool fishes (*Oligocottus maculosus*, *O. snyderi*). *Copeia*, 1976:143-152.
- Nakano, K., and G.K. Iwama. 2002. The 70-kDa heat shock protein response in two intertidal sculpins, *Oligocottus maculosus* and *O. snyderi*: relationship of hsp 70 and thermal tolerance. *Comp. Biochem. Physiol., A* 133:79-94.
- Neat, F.C. 2001. Male parasitic spawning in two species of triplefin blenny (Tripterygiidae): contrasts in demography, behaviour and gonadal characteristics. *Environ. Biol. Fish.*, 61(1):57-64.

- Norton, S.F., and A.E. Cook. 1999. Predation by fishes in the intertidal. Pp. 223-263. In: intertidal fishes: life in two worlds. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- O'Connell, C.P. 1953. The life history of the cabezon *Scorpaenichthys marmoratus* (Ayres). Calif. Fish Game, 93:1-76.
- Ojeda, F.P., and A.A. Muñoz. 1999. Feeding selectivity of the herbivorous fish *Scartichthys viridis*: effects on macroalgal community structure in a temperate rocky intertidal coastal zone. MEPS, 184:219-229.
- Ong, K., P. Stevens, and P. Wright. 2007. Gill morphology of the mangrove killifish (*Kryptolebius marmoratus*) is plastic and changes in response to terrestrial air exposure. J. Exp. Biol., 210(7):1109-1115.
- Ord, T.J., and G.M. Cooke. 2016. Repeated evolution of amphibious behavior in fish and its implications for the colonization of novel environments. Evolution, 70(8):1747-1759.
- Ord, T.J., T.C. Summers, M.M. Noble, and C.J. Fulton. 2017. Ecological release from aquatic predation is associated with the emergence of marine blenny fishes onto land. Am. Nat., 189(5):570-579.
- Osland, M.J., P.W. Stevens, M.M. Lamont, R.C. Brusca, K.M. Hart, J.H. Waddle, C.A. Langtimm, et al. 2021. Tropicalization of temperate ecosystems in North America: The northward range expansion of tropical organisms in response to warming winter temperatures. Glob. Change Biol., 27(13):3009–3034.
- Pace, C., and A.C. Gibb. 2014. Sustained periodic terrestrial locomotion in air-breathing fishes. J. Fish Biol., 84(3):639-660.
- Paul, N., S.C. Novais, M.F.L. Lemos, and A. Kunzmann. 2018. Chemical predator signals induce metabolic suppression in rock goby (*Gobius paganellus*). PLOS One, 13(12):e0209286.
- Pawlik, J.R., K.F. Albizati, and D.J. Faulkner. 1986. Evidence of a defensive role for limatulone, a novel triterpene from the intertidal limpet *Collisella limatula*. MEPS, 30:251-260.
- Pelster, B., C.R. Bridges, and M.K. Grieshaber. 1988. Physiological adaptations of the intertidal rockpool teleost *Blennius pholis* L., to aerial exposure. Respir. Physiol., 71:355-374.
- Petersen, C.W., C. Mazzoldi, K.A. Zarrella, and R.E. Hale. 2005. Fertilization mode, sperm characteristics, mate choice and parental care patterns in *Artedius* spp. (Cottidae). J. Fish Biol., 67(1):239-254.
- Petersen, C.W., K.A. Zarrella, C.A. Ruben, and C. Mazzoldi. 2004. Reproductive biology of the rosylip sculpin, an intertidal spawner. J. Fish Biol., 64(4):863-875.
- Pfister, C.A. 1992. Sculpin diversity in tidepools. Northwest Environ., 8:156-157.
- Pfister, C.A. 1995. Estimating competition coefficients from census data: a test with field manipulations of tidepool fishes. Am. Nat., 146:271-291.
- Pfister, C.A. 1998. Extinction, colonization, and species occupancy in tidepool fishes. Oecologia, 114(1):118-126.
- Pfister, C.A. 1999. Recruitment of intertidal fishes, p. 181-196. In: Intertidal fishes: life in two worlds. (MH Horn, KLM Martin, MA Chotkowski (eds.). Academic Press.
- Polivka, K.M., and M.A. Chotkowski. 1998. Recolonization of experimentally defaunated tidepools by northeast Pacific intertidal fishes. Copeia, 1998:456-462.
- Prochazka, K. 1996. Seasonal patterns in a temperate intertidal fish community on the west coast of South Africa. Environ. Biol. Fish., 45:133-140.
- Rafaelli, D., and S. Hawkins. 1996. Intertidal ecology. Chapman and Hall, London.

- Ragland, H.C., and E.A. Fisher. 1987. Internal fertilization and male parental care in the scalyhead sculpin, *Artedius harringtoni*. Copeia, 1987:1059-1062.
- Ralston, S.L., and M.H. Horn. 1986. High tide movements of the temperate-zone herbivorous fish *Cebidichthys violaceus* (Girard) as determined by ultrasonic telemetry. J. Exp. Mar. Biol. Ecol., 98:35-50.
- Randall, D.J., and Y.K. Ip. 2006. Ammonia as a respiratory gas in water and air-breathing fishes. Respir. Physiol. Neurobiol., 154(1-2):216-225.
- Randall, D.J., and W.W. Burggren, A.P. Farrell, and M.S. Haswell. 1981. The evolution of air-breathing in vertebrates. Cambridge Univ. Press.
- Randall, D.J., Y.K. Ip, S.F. Chew, and J.M. Wilson. 2004. Air breathing and ammonia excretion in the giant mudskipper, *Periophthalmodon schlosseri*. Physiol. Biochem. Zool., 77(5):783-788.
- Richards, J.G. 2011. Physiological, behavioral and biochemical adaptations of intertidal fishes to hypoxia. J. Exp. Biol., 214(2):191-199.
- Richkus, W.A. 1978. A quantitative study of intertidepool movement of the wooly sculpin, *Clinocottus analis*. Mar. Biol., 49:227-284.
- Richkus, W.A. 1981. Laboratory studies of intraspecific behavioral interactions and factors influencing tidepool selection of the wooly sculpin, *Clinocottus analis*. Calif. Fish Game, 67:187-195.
- Ricklefs, R.E., and G.L. Miller. 2000. Ecology. Fourth edition. W.H. Freeman.
- Riegler, K.C. 1976. Oxygen consumption, heart rates, whole body lactate levels, and evaporative water loss in the monkeyface eel *Cebidichthys violaceus* (Family: Stichaeidae), an amphibious marine fish from California. Master's Thesis. Calif State Univ, Fullerton.
- Sayer, M.D.J., and J. Davenport. 1991. Amphibious fish: why do they leave water? Rev. Fish Biol. Fish., 1:159-181.
- Setran, A.C., and D.W. Behrens. 1993. Transitional ecological requirements for early juveniles of two sympatric fishes, *Cebidichthys violaceus* and *Xiphister mucosus*. Environ. Biol. Fish., 37:381-395.
- Shanks, A.L., and C.A. Pfister. 2009. Annual recruitment of three species of tide-pool fishes is driven by variation in springtime coastal hydrodynamics. Limnol. Oceanogr. 54(5):1481-1487.
- Shimizu, N., Y. Sakai, H. Hashimoto, and K. Gushima. 2006. Terrestrial reproduction by the air-breathing fish Andamia tetractyla (Pisces; Blenniidae) on supralittoral reefs. J. Zool., 269(3):357-364.
- Sloman, K.A., M. Mandic, A.E. Todgham, N. Fangue, P. Subrt, and J.G. Richards. 2008. The response of the tidepool sculpin, *Oligocottus maculosus*, to hypoxia in laboratory, mesocosm and field environments. Comp. Biochem. Physiol., A, 149(3):284-292.
- Smyder, E.A., and K.L. Martin. 2002. Temperature effects on egg survival and hatching during the extended incubation period of California grunion, *Leuresthes tenuis*. Copeia, 2002:313-320.
- Souza-Bastos, L.R., M.I. Pascoa, C.A. Freire, and J.M. Wilson. 2014. Ammonia excretion and expression of transport proteins in the gills and skin of the intertidal fish *Lipophrys pholis*. Comp. Biochem. Physiol. A, 167:15-24.
- Stephens, J.S., Jr., R.K. Johnson, G.S. Key, and J.E. McCosker. 1970. The comparative ecology of three sympatric species of California blennies of the genus *Hypsoblennius* Gill (Teleostomi, Blenniidae). Ecol. Monogr., 40:213-233.

- Stepien, C.A. 1987. Color pattern and habitat differences between male, female, and juvenile giant kelpfish. Bull. Mar. Sci., 41:45-58.
- Stepien, C.A. 1990. Population structure, diets and biogeographic relationships of a rocky intertidal fish assemblage in central Chile: high levels of herbivory in a temperate system. Bull. Mar. Sci., 47:598-612.
- Stepien, C.A., H. Phillips, J.A. Adler, and P.J. Mangold. 1991. Biogeographic relationships of a rocky intertidal fish assemblage in an area of cold water upwelling off Baja California, Mexico. Pacif. Sci., 45:63-71.
- Stepien, C.A., and R.H. Rosenblatt. 1991. Patterns of gene flow and genetic divergence in the northeastern Pacific Clinidae (Teleostei: Blennioidei), based on allozyme and morphological data. Copeia, 1991:873-896.
- Stevenson, T.A., and A. Stevenson. 1972. Life Between Tidemarks on Rocky Shores. W.H. Freeman.
- Strathmann, R.R., and H.C. Hess. 1999. Two designs of marine egg masses and their divergent consequences for oxygen supply and desiccation in air. Am. Zool., 39:253-260.
- Takeda, T., A. Ishimatsu, S. Oikawa, T. Kanda, Y. Hishida, and K.H. Khoo. 1999. Mudskipper *Periophthalmus schlosseri* can repay oxygen debts in air but not in water. J. Exp. Zool., 284(3):265-270.
- Takiyama, Y., S. Hamasaki, and M. Yoshida. 2016. Comparison of the visual capabilities of an amphibious and an aquatic goby that inhabit tidal mudflats. Brain Behav. Evol., 87:39-50.
- Taylor, M.H. 1984. Lunar synchronization of fish reproduction. Trans. Amer. Fish. Soc., 113(4):484-493.
- Thomson, D.S., and C.E. Lehner. 1976. Resilience of a rocky intertidal fish community in a physically unstable environment. J. Exp. Mar. Biol. Ecol., 22:1-29.
- Tiralongo, F., D. Tibullo, M.V. Brundo, F.P. De Mendoza, C. Melchiorri, and M. Marcelli. 2016. Habitat preference of combtooth blennies (Actinopterygii: Perciformes: Blenniidae) in very shallow waters of the Ionian Sea, South-Eastern Sicily, Italy. Acta Ichthyol. Piscat., 46(2):65-75.
- Todgham, A.E., G.K. Iwama, and P.M. Schulte. 2006. Effects of the natural tidal cycle and artificial temperature cycling on Hsp levels in the tidepool sculpin *Oligocottus maculosus*. Physiol. Biochem. Zool., 79(6):1033-1045.
- Tomanek, L., and G.N. Somero. 1999. Evolutionary and acclimation-induced variation in the heat-shock responses of congeneric marine snails (genus *Tegula*) from different thermal habitats: implications for limits of thermotolerance and biogeography. J. Exp. Biol., 202:2925-2936.
- Truchot, J.P., and A. Duhamel-Jouve. 1980. Oxygen and carbon dioxide in the marine intertidal environment: diurnal and tidal changes in rockpools. Resp. Physiol., 39:241-254.
- Turko, A.J., and P.A. Wright. 2015. Evolution, ecology and physiology of amphibious killifishes (Cyprinodontiformes). J. Fish Biol., 87(4):815-835.
- Warner, R.R., and R.K. Harlan. 1982. Sperm competition and sperm storage as determinants of sexual dimorphism in the dwarf surfperch, *Micrometrus minimus*. Evolution 36:44-55.
- Watters, J.V., and J.J. Cech Jr. 2003. Behavioral responses of mosshead and woolly sculpins to increasing environmental hypoxia. Copeia, 2003:397-401.
- Wells, A.W. 1986. Aspects of ecology and life history of the woolly sculpin, *Clinocottus analis*, from southern California. Calif. Fish Game, 72:213-226.

- White, G.E., and C. Brown. 2013. Site fidelity and homing behaviour in intertidal fishes. *Mar. Biol.*, 160(6):1365-1372.
- Williams, G.C. 1954. Differential vertical distribution of the sexes in *Gibbonsia elegans* with remarks on two nominal subspecies of this fish. *Copeia*, 1954:267-273.
- Williams, G.C. 1957. Homing behavior of California rocky shore fishes. *Univ Calif. Publ. Zool.*, 59:249-284.
- Wright, W.G., and J.A. Raymond. 1978. Air-breathing in a California sculpin. *J. Exp. Zool.*, 203:171-176.
- Yoshiyama, R.M. 1981. Distribution and abundance patterns of rocky intertidal fishes in central California. *Environ. Biol. Fish.*, 6:315-332.
- Yoshiyama, R.M., and J.J. Cech Jr. 1994. Aerial respiration by rocky intertidal fishes of California and Oregon. *Copeia*, 1994:153-158.
- Yoshiyama, R.M., and J.D.S. Darling. 1982. Grazing by the intertidal fish *Anoplarchus purpurescens* upon a distasteful polychaete worm. *Environ. Biol. Fish.*, 7:39-45.
- Yoshiyama, R.M., K.B. Gaylord, M.T. Philippart, T.R. Moore, J.R. Jordan, C.C. Coon, L.L. Schalk, C.J. Valpey, and I. Tosques. 1992. Homing behaviour and site fidelity in intertidal sculpins (Pisces: Cottidae). *J. Exp. Mar. Biol. Ecol.*, 160:155-130.
- Yoshiyama, R.M., A.L. Knowlton, J.R. Welter, S. Comfort, B.J. Hopka, and W.D. Wallace. 1996b. Laboratory behaviour of mosshead sculpins *Clinocottus globiceps* toward their sea anemone prey. *J Mar. Biol. Ass. UK*, 76:793-809.
- Yoshiyama, R.M., C. Sassaman, and R.N. Lea. 1986. Rocky intertidal fish communities of California: temporal and spatial variation. *Environ. Biol. Fish.*, 17:23-40.
- Yoshiyama, R.M., C.J. Valev, L.L. Schalk, N.M. Oswald, K.K. Vaness, D. Lauritzen, and M. Limm. 1995. Differential propensities for aerial emergence in intertidal sculpins (Teleostei; Cottidae). *J. Exp. Mar. Biol. Ecol.*, 191:195-207.
- Yoshiyama, R.M., W.D. Wallace, J.L. Burns, A.L. Knowlton, and J.R. Welter. 1996a. Laboratory food choice by the mosshead sculpin, *Clinocottus globiceps* (Girard) (Teleostei; Cottidae), a predator of sea anemones. *J. Exp. Mar. Biol. Ecol.*, 204:23-42.
- Zander, C.D. 1972. Beziehungen zwischen Korperbau und Lebensweise bei Blenniidae (Pisces) aus dem Roten Meer. I. Aubere Morphologie. *Mar. Biol.*, 13:238-246.
- Zander, C.D., J. Nieder, and K.L. Martin. 1999. Vertical distribution patterns. Pp. 26-53. In: intertidal fishes: life in two worlds. M.H. Horn, K.L.M. Martin, and M.A. Chotkowski (eds.). Academic Press.
- Zarzecny, K.M., M. Rius, S.T. Williams, and P.B. Fenberg. 2024. The ecological and evolutionary consequences of tropicalisation. *Trends Ecol. Evol.*, 39(3):267-279.